



Role of Natural Preservatives and their Effect on Food's Shelf Life

Raja Kumar ^a and Reshu Rajput ^{a*}

^a University Institute of Agricultural Sciences, Chandigarh University, Gharuan-140413, Punjab, India.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/CJAST/2023/v42i474312

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/110394>

Review Article

Received: 03/10/2023

Accepted: 09/12/2023

Published: 13/12/2023

ABSTRACT

All living creatures need food to survive. Food has a limited period to keep it safe for consumption and fresh. To keep the food product safe to consume for an extended period of time, and maintain the quality in terms of its taste and odor. Preservatives are incorporated in the modern food industries. There are two types of preservative present in nowadays market. Nature preservative are safe to consume, but their shelf life is less than the synthetic one. Natural additives, derived from plants, herbs, and other natural sources, can have positive effects on human health. These additives are often rich in nutrients, antioxidants, and bioactive compounds that may contribute to overall well-being. It's important to incorporate a variety of natural additives into a balanced and diverse diet to maximize their potential health benefits. Synthetic preservatives are cost-effective in contributing to a longer shelf life for the food product, so many manufactures use sodium benzoate, potassium sorbate, sorbic acid, and benzoic acid during the processing. This evaluation focuses on natural preservatives which have been used for increasing shelf-life, their mechanisms of action and its uses forms (i.e., powder, fruit, essential oils, extract) to food has been conserved. The amounts of phenolic compound in herbs and spices play an essential role on significant part in their antibacterial properties, However, it is also dependent on a variety of intrinsic and extrinsic variables that must be identified prior to using herbs and spices as additions. As a result, the current review provides a thorough overview of the numerous natural preservatives, focusing on their antioxidant

*Corresponding author: E-mail: reshu.rajput21@gmail.com;

and antimicrobial properties, mechanism of action, and application in the food industry. Natural additives often contain antioxidants that help neutralize free radicals in the body. Antioxidants play a role in protecting cells from oxidative stress and inflammation, potentially reducing the risk of chronic diseases. Many herbs and spices possess natural antimicrobial properties that can help inhibit the growth of bacteria and fungi. This can contribute to food preservation and support overall immune health.

Keywords: Natural preservative; antioxidants effects; shelf life; spices; herbs.

1. INTRODUCTION

Preservatives are substances added to food, pharmaceuticals, cosmetics, and industrial products to keep them from deteriorating and extend their shelf lives. The effects of light, air, humidity, and bacteria all contribute to degradation. Preservatives typically have chelating, antibacterial, and antioxidant properties. Depending on where they come from, preservatives might be labeled as natural or artificial. The growing awareness of the detrimental effects of synthetic preservatives on health has increased the value of natural preservatives. Plants are one of the most important natural preservatives [1]. Some natural preservatives are as: Vinegar, Salt, Spices, Edible oil, honey, Sugar, etc. The widespread issue plaguing the food supply chain is the substantial risk of contamination by harmful microorganisms responsible for both pathogenic infections as well as waste. Dairy products, in particular, are highly susceptible to contamination. Multiple sources of contamination have been identified in the dairy business, particularly in smaller-scale farm production. These sources include the utilization of unpasteurized milk, the hygiene of the dairy processing environment, and, in certain instances, even the use of vegetable coagulants [2]. To generate energy, promote growth, and maintain excellent health, humans and animals ingest food in the form of raw, processed, or specially designed components. Most cases in some instances, there are no restrictions on eating. Nevertheless, excessive use of some kinds of foods high in fat, sugar, and salt could be dangerous to health [3]. The food sector is extremely concerned about microbial food deterioration. According to estimates, microbial activity causes up to 25% of all food produced to be wasted after harvest. Although the development of spoilage microbiota in food poses no threat to human health, it has a detrimental effect on the finished products' overall quality, shelf life, and textural attributes. It also influences customer preferences and

causes considerable economic losses. Thus, for the current worldwide food production, preventing or inhibiting microbial growth in foods is of highest importance [4]. Preservatives are used to inhibit the growth of microorganisms that might cause food spoilage and food poisoning; and to extend the shelf-life of items so that they can be disseminated and sold to consumers with a longer shelf-life. Preservatives and antibacterial agents are critical components of today's food supply. The growing demand for convenience meals, along with the relatively extended shelf life of processed foods, necessitates the usage of chemical food preservatives. Some regularly used preservatives, such as nitrate and salt, have been employed in processed meats and wine for ages. As a result, preservative works well as an antibacterial in high-acid meals, fruit drinks, cider, carbonated beverages, and pickles [5].

Natural preservatives are commonly employed in the food industry to inhibit the growth of undesirable bacteria. Plant antimicrobials can be incorporated into a product's formulation, applied to the surface of food, or incorporated into packaging material. It does, however, have a distinct mode of action. Plant-based essential oils, animal-derived enzymes, microbial-derived bacteriocins, organic acids, and naturally occurring polymers [chitosan] have all been used in the food industry [6]. The scientific community could explore novel chemical techniques to research and develop new compounds, building upon existing natural molecules. By modifying these familiar molecules, there is potential to enhance their performance, achieving dual benefits such as improved coloring and preservation properties. This avenue of research offers a promising way to overcome challenges and capitalize on the utilization of these additive compounds within the food industry [7].

2. NATURAL PRESERVATIVES: MECHANISMS OF ACTIONS

Polyphenols are well-known for their antibacterial and antioxidant properties. Their bioactivity

originates from their ability to carry out a number of critical functions, including:

1. Polyphenols have the ability to neutralize and eliminate free radicals and reactive oxygen/nitrogen species (ROS/RNS), which are unstable molecules that can cause cellular harm.
2. Reducing Oxidized Intermediates: Polyphenols can help reduce and stabilize oxidized intermediates, preventing the harmful effects of oxidative stress on cells and tissues.
3. Inducing Metal Chelation: Some polyphenols can bind to metal ions, a process known as chelation. This can be beneficial in preventing metal-catalyzed oxidative reactions that generate free radicals.
4. Inhibiting Enzymes Responsible for Free Radical Formation: Polyphenols can inhibit enzymes involved in the formation of free radicals, further reducing oxidative damage.

Different types of polyphenols, such as flavonoids and phenolic acids, exhibit antioxidant properties that help prevent deterioration due to oxidation. Here are examples of food products that can benefit from the application of polyphenols: Spinach, carrots, broccoli, poultry, apples, lemons, almonds, tea etc.

Overall, polyphenols are valuable substances for promoting health and avoiding disease by countering oxidative stress and microbial risks in the body [8].

In recent years, there has been an increase in consumer demand for higher-quality, preservative-free, and less processed foods with a longer shelf life. This tendency has fueled interest in employing natural substances, particularly preservatives produced from herbs and plant products, as alternatives to synthetic chemical compounds currently in use [9].

3. CHEMISTRY AND CLASSIFICATION OF POLYPHENOLS

Antimicrobials produced from plants hold significant promise as natural preservatives due to their reputation for safety, health benefits, and the added value they bring through their bioactive and nutritional properties. These natural preservatives have a rich history of traditional consumption by humans over decades.

Polyphenols are secondary metabolites generated from either the shikimate or polyketide pathways in plants, are particularly noteworthy. They exhibit a wide range of variations among different plant species and serve multiple essential functions, including:

1. Pigment Formation: Polyphenols play a vital role in the formation of pigments, contributing to the coloration of plants.
2. Resistance to Environmental Stresses: They help plants endure environmental challenges by providing defense mechanisms against diseases and UV radiation protection.
3. Chemical Messengers: Polyphenols also act as chemical messengers within plants, regulating various physiological processes.

4. OTHER TYPES OF BIOACTIVE COMPOUNDS AS THE NATURAL PRESERVATIVES

These compounds are often made up of one or more aromatic rings that have hydroxyl groups connected to them. Their various chemical structures and functions make polyphenols valuable for various applications, including as natural preservatives in food products [10].

Methanolic and ethyl acetate root extracts had 85.18 and 62.96% activity against *E. coli*, respectively. Furthermore, methanolic (84.61%), chloroform (61.53%), and ethyl acetate root extracts (69.23%) significantly inhibited *S. aureus*. Methanolic leaf extracts had high activity against *A. bumanni* (78.26%), *E. coli* (77.78%), and *S. aureus* (73.07%). They also discovered that methanolic bark extracts had modest effectiveness against *A. bumanni* (65.21%) and *E. coli* (62.96%). Furthermore, the extracts demonstrated strong antioxidant activity, with EC₅₀ values ranging from 157.82 to 361.61g/ml. The antibacterial effects of *Azadirachta indica* (*A. indica*) phenolic extracts were investigated on the shelf-life stability of uncooked beef patties stored at 4°C for 11 days [11].

Plants high in bioactive chemicals can be used as preservatives and antioxidants to naturally protect and enhance the quality of various products (as shown Table 1). These bioactive-rich plants contain natural substances that can help preventing spoilage and oxidation. Instead of relying on synthetic chemicals, which can have health and environmental concerns, bioactive-rich plants offer a more sustainable and

potentially healthier approach to preserving and safeguarding products. Synthetic preservatives have long been used in the food industry to enhance shelf life and prevent spoilage. However, an increasing body of evidence suggests that these artificial additives may come at a cost to human health. Synthetic preservatives such as sulfites and benzoates have been linked to allergic reactions in susceptible individuals. Symptoms may include skin rashes, itching, and, in severe cases, anaphylaxis. Some synthetic preservatives release volatile organic compounds (VOCs) that can trigger respiratory problems. Individuals, especially those with asthma, may experience difficulty breathing, coughing, or wheezing. Certain synthetic preservatives, such as nitrites and nitrates commonly used in processed meats, have been associated with headaches and dizziness. Some studies suggest a potential link between the consumption of certain synthetic preservatives and neurological issues. These may include hyperactivity in children and, in extreme cases, contribute to neurodegenerative conditions.

In Table 1, there is a list of plant to use as natural preservative and also mentioned the used plant part, amount which is added, the effect and its major component in extending the shelf-life of a food product.

5. FACTORS INFLUENCING POSTHARVEST SPOILAGE BY: MOISTURE ACTIVITY

Moisture activity, abbreviated "aw," is a measure of the ratio between a food's equilibrium water vapor pressure (P_w , in kilopascals, kPa) and the saturated vapor pressure of pure water (P_{wo} , in kPa) at the same temperature. This metric is an important measure of how water is coupled to the food, showing the product's moisture availability. It is influenced by a variety of factors, including:

1. **Lipid Oxidation:** The presence of lipids in the food can influence moisture activity, as interactions between water and lipids can affect the food's moisture-holding capacity.
2. **Microbial Growth:** Microorganisms require water to grow and proliferate. Moisture activity plays a significant role in determining whether the water content is sufficient to support microbial growth or if it inhibits it.
3. **Enzymatic and Non-enzymatic Activity:** Moisture activity can impact enzymatic

reactions and other chemical processes within the food, as the availability of water can influence the rate and extent of these reactions.

4. **Changes in Texture Profile:** The moisture content and distribution within a food product can affect its texture and consistency. Moisture activity is a key factor in understanding and controlling these textural attributes.

Overall, moisture activity is a critical parameter in food science and technology as it helps assess the stability, safety, and quality of food products, particularly with regard to water-related reactions and microbial growth [24].

6. FACTORS INFLUENCING POSTHARVEST SPOILAGE BY: OXIDATION ACTIVITY

Water, fat, protein, and carbs make up the primary building blocks of food resources produced from plants and animals. Fats and oils are the most ubiquitous lipid compounds in nature. Oil is liquid at room temperature while fat is solid, indicating a difference in its physical characteristics and consistency. The melting point of lipids (polyunsaturated fatty acids, PUFA) is affected by the amount of double bonds, the length of the carbon chain, and whether the unsaturated fatty acids are cis or trans [25].

7. FACTORS INFLUENCING POSTHARVEST SPOILAGE BY: ANTIOXIDANTS

There are numerous synthetic antioxidant preservatives, such as butylated hydroxyl toluene (BHT), butylated hydroxy anisole (BHA), tertbutyl hydro-quinone (TBHQ), and propyl gallate (PG), are commonly used in meat products to avoid lipid oxidation deterioration. The use of antioxidants in food is governed by international standards or national regulatory regulations. The use of synthetic antioxidants is generally restricted due to the possibility of cancer-causing effects [26].

7.1 Plant-Based Preservatives

Natural herbs and spices have preservation properties based on the kind, nature, and concentration of the organism. Spices and herbs have volatile compounds in them. These compounds are extracted using a variety of techniques and utilized to make preservatives. As self-defense mechanisms, the substances

present in herbs and spices help the plant defend itself against infectious organisms. Plant oils and extracts have long been used as food preservatives and medicinal compounds in medicine, according to sources [27].

Table 1. Use of bioactive-rich plants as preservatives and antioxidants

Plant	Part Used	Amount added in Meat (%)	Meats type	Effect on Food product	Major Component	Reference
Curry	Leaf	2.5	Meat homogenate	Lower level of Thiobarbituric acid reactive substances	Cinnamic, chlorogenic, ferulic, vanillic acids, Tannic, gallic, and caffeic	[12]
Black Pepper	Fruit	1	Fresh pork sausage	Inhibition of lipid oxidation	Capsanthin and capsorubin	[13]
Black Cumin	Seed	>0.5	Fresh minced beef	Inhibited of lipid oxidation	Anthocyanins	[14]
Lotus	Leaf	6	Chicken patties	Lower level of Thiobarbituric acid reactive substances	Citric acid, tannin, malic acid, succinic acid, and tartaric acid	[15]
Garlic	Bulbs	1.5	Chicken patties	inhibitor on both gram-positive and gram-negative bacteria.	Allicin	[16]
Ginger	Rhizome	1.5	Minced meat	antimicrobial activity	Gingerol and shogaols	[17]
Amla	Fruit, seed	1	Goat patties	Inhibition of lipid oxidation	Quercetin, gallic acid, kaempferol, and ellagic acid	[18]
Clove	Flower buds	1	Beef patties	Inhibits Gram-negative and Gram-positive bacteria and yeast	Eugenol	[19]
Broccoli	Leaf	>0.1	Ground beef patties	Lower level of Thiobarbituric acid reactive substances	Phenolic acids and flavonoids	[20]
Lemon	Fruit	0.001	Chicken patties	E. coli O157:H7 was reduced by 1.7 logs	Limonene, Sabinene Citronella, linalool	[21]
Thyme	Leaves	0.8-1.2	Chicken meat	improved both the microbial and oxidative stability	Thymol, terpinene, pcymene, carvacrol, and linalool	[22]
Turmeric	Rhizome	3.5	Rabbit patties	Inhibition of lipid oxidation	Curcumin	[23]

7.2 Preservative Classification

There are primarily two groups of preservatives:

- I. Class I: The food preservatives derived from nature, such as: sugar, salt, vinegar, honey, spices, edible oils, etc., were included in this class.
- II. Class II: Benzoates, sorbates, potassium nitrites and nitrates, sulfites, glutamates, and glycerides are examples of chemical, semi-synthetic, and synthetic food preservatives.

Humans and animals who consume or use foods or pharmaceuticals containing more than one preservative face a significant risk of chemical exposure. Chemical preservatives are classified into three types: antimicrobial, antioxidant, and anti-enzymatic [28].

Food preservatives are classified into two classes: Class I and Class II. As stated in Fig. 1, natural preservatives are classed as Class I, whereas synthetic or chemical preservatives are designated as Class II.

However, only one type of Class II preservative should be used in a specific food product because individuals who ingest too much of them, such as sulfites, benzoates, sorbates, and so on, may be harmed. The classification of both types of preservatives is depicted in Fig. 1.

7.3 Function of Preservatives

1. Enhancement of nutritional value of food.

Traditional food preservation methods are inefficient in preventing the spread of food-borne pathogens. The increased desire for chemical-free food has cleared the path for antimicrobials to be used in the food industry. Antimicrobials are a revolutionary technology utilized by the food industry to increase food shelf life while also addressing quality and safety problems [30].

2. To improve quality and lower waste.

Food preservation is the treatment and handling of food to prevent or considerably slow deterioration (loss of quality, edibility, or nutritious value) caused or accelerated by microorganisms. Some approaches, on the other hand, use harmless bacteria, yeasts, or fungi to add specific attributes and preserve food (e.g., cheese, wine). It is critical to maintain or create

nutritional value, texture, and flavor in order to maintain its worth as a food. This is culturally dependent, as what qualifies as human nourishment in one society may not qualify in another. Preventing the growth of bacteria, fungi, and other microorganisms, as well as slowing the oxidation of fats that causes rancidity, are common methods of preservation. It also covers methods that are used to prevent natural aging and discoloration that can occur during food preparation, such as the enzymatic browning reaction that occurs when apples are sliced [31].

3. To increase consumer acceptance.

The safety of natural substances is being questioned, and customers are increasingly requesting them. As a result of their inherent qualities, spices are a powerful tool for the food sector. Spices do have antioxidant properties, owing to the presence of phenolic chemicals. They have antioxidant properties because they scavenge free radicals, chelate transition metals, quench singlet oxygen, and increase the activity of antioxidant enzymes. At 37 and 80 degrees Celsius, the CO₂ extract of ginger demonstrated in vitro efficacy comparable to that of BHT in preventing lipid peroxidation. Furthermore, in a model mixture replicating a heated potato matrix (180°C for 20 minutes), pimento and black pepper extracts reduced acrylamide production by up to 75 and 50%, respectively [32].

4. Inhibition of microbial growth.

Natural food preservatives do not yield negative side effects like synthetic preservatives. Natural food preservatives are usually sourced from plants, animals, or microorganisms. Natural preservatives are generally employed in the food industry to prevent the growth of undesirable bacteria. Plant antimicrobials can be used in product formulation, as a coating on the surface of meals, or as a component of packaging material. Its manner of action, however, is distinct. Plant-derived essential oils, animal-derived enzymes, microbe-derived bacteriocins, organic acids, and naturally occurring polymers (chitosan) have all been used in the food sector [33].

5. They help to extend the shelf life of processed goods.

The current study aims to increase the shelf life of fresh prepared foods by employing plant-based natural preservatives. This study makes

use of lemon juice, basil leaf extract, mint leaf extract, and lemongrass essential oil. The goal is to compare the various natural preservatives based on enzymatic and microbiological inhibition as well as sensory acceptance. The formulation of a natural preservative was optimized using a multi-objective genetic algorithm. Furthermore, principal component analysis is used to visualize changes in several quality metrics after preservative-added storage [34].

8. EXAMPLES OF SOME NATURAL PRESERVATIVES ADDED FOR FOOD PRESERVATION

8.1 Salt

Salt acts as a preservative because it can attract water, and as a result, the material's water activity reduces, preventing microbe growth and reproduction. Salt preferentially inhibits polluting bacteria such as spores and proteolytic microbes [35]. Salting food can also produce osmotic stress in microbial cells, which causes the cell to lose water and eventually results in cell death or reduced growth [36]. Furthermore, salt may limit the solubility of oxygen for some microorganisms, interfere with biological enzymes, or make cells work harder to discharge sodium ions from the cell, all of which can inhibit growth [37]. Salting or curing eliminates moisture

from meat via osmosis. Curing meat involves the use of salt, sugar, or a mixture of the two. Furthermore, nitrates and nitrites are widely used to cure meat, giving it a unique pink tint and limiting the growth of *Clostridium botulinum* [38]. Salt, in moderation, plays a crucial role in maintaining various physiological functions in the human body. Here are some of the benefits of salt on human health: Hydration, Electrolyte Balance, Nerve Function, Digestive health, Blood pressure regulation, etc.

8.2 Sugar

Fruit and some vegetable products use sugar to enhance flavor and extend shelf life. Whole and cut-in-half fruit, relishes, pickles, jams, and jellies are examples of these items. Except for vegetables, sweeteners are regarded an essential element in the majority of canning products. They operate as preservatives, preserving the desired appearance, flavor, color, and body of the items. These qualities will change if the type and amount of sugar in standardized preservation recipes are changed. Sugar (sucrose) and corn syrup are the two main sweeteners used in canning. Saccharin, a noncaloric sweetener, is found in dietetic meals. Noncaloric sweeteners aspartame and acesulfame K are allowed for usage in a restricted variety of foods [39].




Fig. 1. Categories of food preservatives [29]

At 10% concentrations, the relative sweetness of invert sugar compared to sucrose. In terms of sweetness, sucrose is sweeter in concentrations less than 10%, while invert is sweeter at concentrations greater than 10%. However, due to the sweetness-enhancing property of sucrose, a partially inverted sucrose solution will be sweeter than a wholly inverted solution. The sweetness of the different sugars varies. Sucrose sweetness is instantly sensed and achieves a maximum intensity, but dextrose sweetness stimulates the taste organs more slowly and reaches a maximum intensity later. The fundamental flavor of glucose is sweet, but the secondary flavors are bitter, sour, or tart, whereas secondary flavors are absent in sucrose [40]. While sugar is a source of quick energy for the body, it's important to note that excessive consumption of added sugars can have negative effects on health. However, there are some benefits of sugar when consumed in moderation: Quick energy, preservation of food, weight gain, brain function, etc.

8.3 Vinegar

In the past, raw ingredients containing starch and sugar were used to make vinegar. In a two-stage fermentation, ethanol is produced first, followed by acetic acid. Traditional vinegar manufacturing requires a longer fermentation period of roughly 30 days, and it serves as a starter culture for vinegar manufacture at commercial platforms, which is finished in 24 hours. Traditional vinegar is made from apple, coconut, plum, tomato, grape, and potato juices [41]. Wine and beer are the most likely alcoholic beverages that are used to make vinegar when it is diluted. In addition, many additional basic materials, including as rice, sugarcane, and malt, are already present in the system as alternatives to both beer and wine. These materials undergo fermentation processes to produce alcoholic beverages. Before beginning the manufacturing of vinegar, certain treatments are given after the collection of alcoholic liquid, including pasteurization and filtering [42].

Herbs and fruits with a variety of specialized functions are integrated into the vinegar production method to produce flavor-infused vinegar. Implemented are the herb categories of tarragon, garlic, and basil. Lemon, pomegranate, apple, and cherries are all included [43]. Spices can demonstrate antimicrobial effects through two distinct mechanisms: firstly, by impeding the proliferation of spoilage microorganisms, thus

aiding in food preservation; and secondly, by limiting or regulating the growth of harmful bacteria and thereby improving food safety [44]. Vinegar, particularly varieties like apple cider vinegar and white vinegar, has been studied for its potential health benefits. It's important to note that while some studies suggest certain advantages, more research is needed, and the effects can vary among individuals. Here are some potential health benefits associated with the consumption or use of vinegar: Blood sugar control, weight management, improved digestion, antimicrobial properties, antioxidant effects, etc.

8.4 Spices

The phenolic chemicals found in spices give them their antioxidant properties. They exhibit antioxidant properties through scavenging free radicals, chelating transition metals, quenching singlet oxygen, and increasing the activity of antioxidant enzymes [45]. Ginger's CO₂ extract demonstrated in vitro efficacy that was comparable to BHT's in preventing lipid peroxidation at both 37°C and 80°C [46]. Lipid oxidation is a major cause of food rotting. As a result, the food industry employs antioxidants such as butylated hydroxytoluene (BHT) and butylated hydroxyanisole (BHA) to prevent spoilage [47]. Customers, on the other hand, are increasingly preferring natural substances, and their safety is questioned. Because of their inherent qualities, spices are a powerful tool for the food sector [48]. Spices not only add flavor and aroma to food but also come with a variety of potential health benefits. Here are some of the advantages associated with the consumption of spices: Anti-Inflammatory effects, antioxidant properties, improved digestion, etc. Vinegar, particularly varieties like apple cider vinegar and white vinegar, has been studied for its potential health benefits. It's important to note that while some studies suggest certain advantages, more research is needed, and the effects can vary among individuals. Here are some potential health benefits associated with the consumption or use of vinegar: Blood sugar control, weight management, improved digestion, antimicrobial properties, antioxidant effects, etc.

8.5 Honey

Honey's antimicrobial properties can be divided into peroxide and non-peroxide components. Honey's main antibacterial component is hydrogen peroxide, which is created when the enzyme glucose oxidase, present in the

hypopharyngeal glands of honey bees, oxidizes glucose and other monosaccharides [49]. Honey is frequently used as a preservative in foods including milk, cheese, and turkey slices. Honey was utilized to improve the shelf life of kunun zaki, a Nigerian non-alcoholic beverage, by up to five days at room temperature 25°C [50]. Honey is a natural sweetener that has been used for centuries for its potential health benefits. Here are some of the advantages associated with the consumption of honey: Antioxidant properties, wound healing, cough suppressant, skin health, etc.

8.6 Edible Oils

Essential oils are aromatic and volatile liquids extracted from numerous parts of the plant, including the flowers, roots, bark, leaves, seeds, peel, fruits, wood, and whole plant. Essential oils have been used in gastronomy as spices or herbs, as well as in fragrance, cosmetics, and medicine since antiquity [51]. The bulk of essential oils are composed of low-molecular-weight volatile chemical molecules. The ingredients of essential oils can be splitted into two categories: terpenoids and non-terpenoid hydrocarbons. Terpenoids, which are generated by combining two (monoterpene), three (sesquiterpene), or four (diterpene) isoprene units, and phenylpropanoids (non-terpenoid hydrocarbons) are the two structural families of phenolic chemicals [52].

The effect of an edible gelatin covering containing orange leaf essential oil on the shelf life of chilled pink shrimp. Adding 2% orange leaf essential oil to the gelatin solution increased the shelf life of shrimp by 10 days as compared to uncoated samples [53]. Phenolic compounds in EOs have a strong interaction with the lipids in cell membranes, making them more permeable. This causes the cellular structure to be destroyed or disrupted, as well as the release of intracellular chemicals. In the formulation of EO-edible coatings, a number of microbial species and strains were tested [54]. Carvacrol, eugenol, and thymol are common phenolic components found in high amounts in EOs with the strongest antibacterial activity against foodborne pathogens. The phenolic chemical family, which has a wide range of structural modifications, is one of the most diverse families of secondary metabolites. Because the hydroxyl (-OH) groups in phenolic compounds can interact with bacterial cell membranes, causing membrane structures to rupture and cellular components to leak, it is

thought that these groups have an inhibitory impact [55]. Edible oils are a crucial component of a balanced diet and contribute to various health benefits when consumed in moderation. Different oils offer different nutritional profiles, and their benefits can vary. Here are some general advantages associated with the consumption of various edible oils: Sources of healthy fats, heart health, vitamin E content, improved nutrient absorption, etc.

8.7 Herbs

Plants and plant materials are rich in bioactive compounds (such as natural antioxidants) that have the potential to improve health, natural antioxidants may be a solution to the problem of preventing lipid oxidation processes. For generations, people have utilized herbs and spices to make food. They are used in a number of recipes to enhance flavor and scent or to extend food shelf life [56]. The herb thyme is a raw material used in a variety of food industries and adds flavor and scent to cooked foods (meat, sausages). Thyme is a fragrant shrub that is used as a spice and for therapeutic purposes. Additionally, it gives food scent and functions as a preservative, antioxidant, antifungal, and antibacterial. Oregano, a popular spice, contains essential oils such as thymol and carvacrol [57].

Natural antimicrobials can come from a variety of sources, including plants, animals, bacteria, algae, and fungi. Plant extracts have the advantage of being used by humans for thousands of years. Many plants are employed in traditional medicine, functional foods, nutritional supplements, and the creation of recombinant proteins, in addition to antimicrobial. Their ability to inhibit/control the growth of germs, particularly harmful bacteria, as well as to manage natural deteriorating processes (food preservation), lends them to their role as antimicrobial [58]. Herbs have been used for centuries not only to enhance the flavor of food but also for their potential health benefits. Many herbs contain a variety of compounds with antioxidant, anti-inflammatory, and antimicrobial properties. Here are some general health benefits associated with the consumption of herbs: Rich in antioxidants, anti-inflammatory properties, Digestive health, improves immune function, antimicrobial properties, etc.

9. CONCLUSIONS

The present paper presents that a number of natural preservatives developed from different

sources with different preservative properties. Preservatives are simply an option for extending the life of the product by blocking the growth of spoilage bacteria and preventing oxidative reactions. We can get the taste of non-seasonal fruit and vegetable at every season. If possible, we should only use natural preservatives. Natural preservatives may have a smaller environmental impact than synthetic alternatives because they are frequently produced from renewable plant sources. Many consumers prefer products incorporated with natural preservatives, reflecting an increasing demand of natural processed product than synthetic processed products. Herbs and spices, for example, are natural preservatives that can improve the flavor and general quality of food products. Natural preservatives decrease the need for synthetic chemical additives, which some consumers may be concerned about due to potential health hazards and allergies. Many natural preservatives come from plants and have a long history of being used safely in traditional cuisines. Using them properly in a moderate amount can give healthier and more alternatives to synthetic preservatives for consumers.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Busra KP, and Seyhan GV. Natural Preservatives. International Journal of Traditional and Complementary Medicine Research. 2021;2(3):184-192. DOI: 10.53811/ijtcmr.987607.
2. Mena R, Pamela M. Natural Preservatives from Plant in Cheese Making. National Library of Medicine. 2020;10(4):749. DOI:10.3390/ani10040749
3. Sharif ZIM, Mustapha FA, Jai J, Mohd N, Yusof Zaki NAM. Methods for preservation and natural preservatives for extending the food longevity. Chemical Engineering Research Bulletin. 2017; 19:145-153. DOI:10.3329/CERB.V19I0.33809
4. Moreno B, Andrea L, Simona de N, Patrizia M, and Chrissanthy P. Natural Preservatives to Improve Food Quality and Safety. Hindawi Journal of Food Quality. 2017;1090932. Available:https://doi.org/10.1155/2017/1090932
5. Mirza SK, Asema UK, Sayyad SK. To study the harmful effects of food preservatives on human health. Journal of Medicinal Chemistry and Drug Discovery. 2017;2(2):610-616.
6. Ejigayehu T, Sirawdink FF, Vasantha HP Rupasinghe, and Ebisa, OK. Potentials of Natural Preservatives to Enhance Food Safety and Shelf Life: A Review. Hindawie Scientific World Journal. 2022;9901018:11. Available:https://doi.org/10.1155/2022/9901018
7. Claudia Novais, Adriana K Molina, Rui MV Abreu, Celestino SB, Isabel CFRF, Carla Pereira, and Lillian Barros. Natural Food Colorants and Preservatives: A Review, a Demand, and a Challenge. Journal of Agricultural and Food Chemistry. journal of agricultural and food chemistry. 2022;70:2789-2805. Available:https://doi.org/10.1021/acs.jafc.1c07533.
8. Michel M. Beya, Michael E. Netzel, Yasmina Sultanbawa, Heather Smyth and Louwrens C Hoffman. Plant-Based Phenolic Molecules as Natural Preservatives in Comminuted Meats. A Review. Antioxidants. 2021;10(2):263. Available:https://doi.org/10.3390/antiox10020263.
9. Bag BB, Panigrahi C, Gupta S, and Mishra, HN. Efficacy of plant-based natural preservatives in extending shelf life of sugarcane juice: Formulation optimization by MOGA and correlation study by principal component analysis. Applied Food Research. 2022;2(2):100164. Available:https://doi.org/10.1016/j.afres.2022.100164
10. Ofosu FK, Banan MDE, Fazle E, Ramachandran C, Lee BH, and Oh DH. New Insights on the Use of Polyphenols as Natural Preservatives and Their Emerging Safety Concerns. Frontiers in Sustainable Food Systems. 2020;4. Available:https://doi.org/10.3389/fsufs.2020.525810
11. Ouerfelli M, Villasante J, Kaâb LBB, and Almajano MP. Effect of neem (*Azadirachta indica* L.) on lipid oxidation in raw chilled beef patties. Antioxidants. 2019;8(8):305. DOI: 10.3390/antiox8080305
12. Yogesh K, Jha SN, Yadav DN. Antioxidant Activities of *Murraya koenigii* (L.) Spreng Berry Extract: Application in Refrigerated (4 ± 1 °C) Stored Meat Homogenates. Agricultural Research. 2012;1(2):183-189.

13. Luis M, Irene C, A Jose, Bel and Pedro R. Effect of *Capsicum annum* (Red Sweet and Cayenne) and *Piper nigrum* (Black and White) Pepper Powders on the Shelf Life of Fresh Pork Sausages Packaged in Modified Atmosphere. *Journal of Food Science*. 2006;71:48–53.
14. Jia N, Kong B Liu, Q Diao, X Xia. Antioxidant activity of black currant (*Ribes nigrum* L.) extract and its inhibitory effect on lipid and protein oxidation of pork patties during chilled storage. *Meat Science*. 2012;91(4):533–539. Available:<https://doi.org/10.1016/j.meatsci.2012.03.010>
15. Yun-Sang C, Ji-Hun C, Hack-Youn K, Hyun-Wook K, Mi-Ai L, Hai-Jung C, Sung K L, Cheon-J. K. Effect of Lotus (*Nelumbo nucifera*) Leaf Powder on the Quality Characteristics of Chicken Patties in Refrigerated Storage. *Food Science of Animal Resources*. 2011;31(1):9–18.
16. Rana SV, Pal R, Vaiphei K, Sanjeev K Sharma and RP Ola. Garlic in health and disease. *Nutrition Research Reviews*. 2011;24(1):60-71. Available:<https://doi.org/10.1017/S0954422410000338>
17. Draszanowska A, Karpinska TM, Olszewska MA. The effect of ginger rhizome and refrigerated storage time on the quality of pasteurized canned meat. *Food Science Technology International*. 2020;26(4):300–310. Available:<https://doi.org/10.1177/1082013219889439>
18. Bariya AR, Patel AS, Gamit VV, Bhedi KR and Parmar RB. Assessment of Antioxidant and Sensory Properties of Amla (*Embolica officinalis*) Fruit and Seed Coat Powder Incorporated Cooked Goat Meat Patties. *International Journal of Current Microbiology and Applied Sciences* Sci. 2018;7(7):3306–3318
19. Nunez L, and Aquino MD. Microbicide activity of clove essential oil (*Eugenia caryophyllata*). *Springer Brazilian Journal of Microbiology*. 2012;43(4):1255–1260. Available:<https://doi.org/10.1590%2F51517-83822012000400003>
20. Kim SJ, Cho AR, and Han J. Antioxidant and antimicrobial activities of leafy green vegetable extracts and their applications to meat product preservation. *Food Control*. 2013;29(1):112–120. Available:<https://doi.org/10.1016/j.foodcont.2012.05.060>
21. Kotzekidou P, Giannakidis P, and Boulamatsis A. Antimicrobial activity of some plant extracts and essential oils against foodborne pathogens in vitro and on the fate of inoculated pathogens in chocolate. *LWT--Food Science and Technology*. 2018;41(1):119–127.
22. Pellegrino L, and Tirelli A. A sensitive HPLC method to detect hen's egg white lysozyme in milk and dairy products. *International Dairy Journal*. 2000;10(7): 435–442.
23. Mancini S, Preziuso G, Paci G. Effect of turmeric powder (*Curcuma longa* L.) and ascorbic acid on antioxidant capacity and oxidative status in rabbit burgers after cooking. *World Rabbit Science*. 2016;24(2):121–127 PMC7223362.
24. Yuxi G, Xuefeng C, Pin G, Ruotong W, Aoyang H, Zhenfang D, Zhuoya Q, Hui L, Jiating W, Wenbo Y, Wenjuan Y, Jing W, and Nan L. Advances in the Role and Mechanisms of Essential Oils and Plant Extracts as Natural Preservatives to Extend the Postharvest Shelf Life of Edible Mushrooms. *Foods*. 2023;12(4). Available:<https://doi.org/10.3390/foods12040801>.
25. Nur J, Safri I, Achmad Z, Pham VH, and Dikdik K. Potential of *Syzygium polyanthum* as Natural Food Preservative. *A Review Foods*. 2023;12(12):2275. Available:<https://doi.org/10.3390/foods12122275>
26. Celia j Hugo and Arno Hugo. Current trends in natural preservatives for fresh sausage products. *Trends in Food Science and Technology*. 2015;45(1):12-23. Available:<http://dx.doi.org/10.1016/j.tifs.2015.05.003>
27. Nalan G. Novel natural food preservatives and applications in seafood preservation: a review. *Journal of the Science of Food and Agriculture*. 2018;99(5):2068–2077. DOI 10.1002/jsfa.9416.
28. Sangeeta D, Palash P, Narendra V, Sapna M, Anil K. A Review on Food Preservation: Methods, harmful effects and better alternatives. *Asian Journal of Pharmacy and Pharmacology*. 2017; 3(6):193-199.
29. Anand SP and Sati N. Artificial preservatives and their harmful effects: looking toward nature for safer alternatives. *International Journal of Pharmaceutical Sciences and Research*. 2013;4(7): 2496-2501.

30. Teshome E, Forsido SF, Rupasinghe HP, and Keyata EO. Potentials of Natural Preservatives to Enhance Food Safety and Shelf Life: A Review. *The Scientific World Journal*. 2022;11:9901018.
31. Abdulumumeen HA, Risikat AN, and Sururah, AR. Food: Its preservatives, additives and applications. *International Journal of Chemical and Biochemical Sciences*. 2012;(1):36-47. DOI:10.13140/2.1.1623.5208
32. Anamaria P, Muste S, paucean A, Chis S, Man S, Salanta, L, Marc R, Muresan A, and Martis G. Herbs and spices in terms of food preservation and shelf life. *Hop and Medicinal Plants*. 2019; 1(2):2360-0179.
33. Ejigayehu Teshome, Sirawdink Fikreyesus Forsido, HP Vasantha Rupasinghe, and Ebisa Olika Keyata. Potentials of Natural Preservatives to Enhance Food Safety and Shelf Life: A Review. *The Scientific World Journal*. 2022;22(1): Available:https://doi.org/10.1155/2022/9901018
34. Bijaya BB, Chirasmita P, Sakshi and Mishra HN. Efficacy of plant-based natural preservatives in extending shelf life of sugarcane juice: Formulation optimization by MOGA and correlation study by principal component analysis. *Applied Food Research*. 2022;2(2):100164. Available:https://doi.org/10.1016/j.afres.2022.100164
35. Rossi I, Georgina J, Hilman M, Namira AY. A Mini-Review of Salting Techniques to Improve Food Quality. *International Journal of Scientific and Technology Research*. 2021;10(1):285-289. Available:https://www.researchgate.net/publication/348759004
36. Davidson PM. Chemical preservatives and natural antimicrobial compounds. *Food microbiology: Fundamentals and frontiers*. 2001;4(6).
37. Shelef LA, and Seiter J. Indirect and miscellaneous antimicrobials. Taylor and Francis Group an informa business. 2005;3:573-598.
38. Hamid AA, Ahmed R, and Sururah AR. Food: Its preservatives, additives and applications. *International Scientific Organization*. 2012;1:36-47.
39. Georgia CL. Reduced Sugar and Sugar-free Food Preservation. *Utahstate University Extension FN*. 1992;209 (Book).
40. Cotton HR, Rebers AP, Maudru EJ, and Rorabaugh G. The Role of Sugar in the Food Industry. *Advances in chemistry series publications*. 2023;1(1):3-20.
41. Bhat SV, Akhtar R, and Amin T. An overview on the biological production of vinegar. *International Journal of Fermented Foods*. 2014;3(2):139. Available:https://www.researchgate.net/publication/340816450
42. Hailu S, Admassu S, and Jha YK. Vinegar production technology—An overview. *Beverage and Food World*. 2012;3(2):139-155.
43. Hemke JE, Prasad R, Sawinder K, Kumbhar PT, and Jyoti S. Vinegar: a traditional functional food. *THINK INDIA JOURNAL*. 2020;22(34): ISSN:0971-1260
44. Tajkarimi MM, Ibrahim SA, Cliver DO. Antimicrobial herb and spice compounds in food. *Food control*. 2010;21(9): Available:https://doi.org/10.1016/j.foodcont.2010.02.003
45. Rubio L, Motilva MJ, and Romero MP. Recent advances in biologically active compounds in herbs and spices: a review of the most effective antioxidant and anti-inflammatory active principles. *Critical Reviews in Food Science and Nutrition*. 2013;53(9):943–953. DOI: 10.1080/10408398.2011.574802
46. Stoilova I, Krastanov A, Stoyanova A, Denev P, and Gargova S. Antioxidant activity of a ginger extract (*Zingiber officinale*). *Food Chemistry*. 2007;102(3):764–770. DOI: 10.1016/j.foodchem.2006.06.023
47. Stoilova I, A. Krastanov, A. Stoyanova, P. Denev, and S. Gargova. Antioxidant activity of a ginger extract (*Zingiber officinale*). *Food Chemistry*. 2007; 102(3):764-770. Available:https://doi.org/10.1016/j.foodchem.2006.06.023
48. Davide G, Danka B, Sahdeo P, and Amit KT. Beneficial Effects of Spices in Food Preservation and Safety. *Frontiers in Microbiology*. 2016;7. DOI: 10.3389/fmicb.2016.01394
49. Krushna NSA, Kowsalta A, Radha S, and RB Narayanan. Honey as a natural preservative of milk. *Indian Journal of Experimental Biology*. 2007;45(5):459-464. Available:https://www.researchgate.net/publication/34376926_Honey_as_a_natural_preservative_of_milk
50. Nadja WJ, TS Azilah A, and Aishath N. The antibacterial properties of stingless bee

- honey in Malaysia and its effects as food preservative. The Maldives National Journal of Research. 2023; 10(2):63-72. Available:<https://www.researchgate.net/publication/371011158>
51. Morten Hyldgaard, Tina Mygind, and Rikke Louise Meyer. Essential oils in food preservation: mode of action, synergies, and interactions with food matrix components. *Frontiers in Microbiology*. 2012;3(1): Available:<https://doi.org/10.3389/fmicb.2012.00012>.
52. Srutee R, Srushti T, Ram KD, Suraj M, Jorddy C, Prem Prakash S, Purnima D Amin, Kirtiraj KG, Eloisa H de AA, Mozaniel, S de O. Recent trends in the application of essential oils: The next generation of food preservation and food packaging. *Trends in Food Science and Technology*. 2022;129:421-439. Available:<https://doi.org/10.1016/j.tifs.2022.10.012>
53. Vinay KP, Rayees UI, Rafeeya S, Aamir and H Dar. A comprehensive review on the application of essential oils as bioactive compounds in Nano-emulsion based edible coatings of fruits and vegetables. *Applied Food Research*. 2022;2(1):100042. Available:<https://doi.org/10.1016/j.afres.2022.100042>
54. Jian J, Yunfei X, Yahui G, Yuliang C, He Q. and Weirong Y. Application of edible coating with essential oil in food preservation. *Critical Reviews in Food Science and Nutrition*. 2018;59(15): 2467-2480. DOI: 10.1080/10408398.2018.1456402
55. Laranjo M, AM Fernandez-Leon, Potes ME, AC Agulheiro-Santos and Elias M. Use of essential oils in food preservation. *Antimicrobial research: Novel bioknowledge and educational programs*. 2017;177-188. Available:<https://www.researchgate.net/publication/320729063>
56. Velasco V, and Williams P. Improving meat quality through natural antioxidants. *Chilean Journal of Agricultural Research*. 2011;71(2):313–322. Available:<http://dx.doi.org/10.4067/S0718-58392011000200017>
57. Awahd HA, Al-Sirrag MAAW, Abbas ZH, Mahmmod RM and Wahed SA. Improving the quality of beef burger by adding thyme and rosemary powder. *Journal of Global Pharma Technology*. 2018;10:949–954. ISSN: 0975 -8542.
58. Gracia CM, Bermúdez CAG, Valcarcel AM C, Pascual M S, and Saseta, CF. Use of herbs and spices for food preservation: advantages and limitations. *Current Opinion in Food Science*. 2015; 6:38-43. Available:<https://doi.org/10.1016/j.cofs.2015.11.011>

© 2023 Kumar and Rajput; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/110394>