

# **Analysis of GIS Based Morphometric Parameters and Hydrological Changes in Indrawati River Sub-Basin of Godawari Basin, Chhattisgarh, India**

**Adikant Pradhan<sup>1\*</sup>, T. Chandrakar<sup>1</sup>, A. Kerketta<sup>1</sup>, S. K. Nag<sup>1</sup>  
and S. C. Mukherjee<sup>1</sup>**

<sup>1</sup>S.G. College of Agriculture and Research Station, IGKV, Kumhrawand, Jagdalpur (C.G.)  
494 005, India.

## **Authors' contributions**

*This research work was carried out in collaboration of all authors. Author AP framed research work and data analysis with writing manuscript, wrote the protocol and wrote the first draft of the manuscript. Authors TC and AK managed the analyses of the study. Author AK managed the literature searches. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/CJAST/2020/v39i1330682

### Editor(s):

(1) Dr. Vyacheslav O. Vakhnenko, Institute of Geophysics National Academy of Sciences of Ukrainian, Ukraine.

### Reviewers:

(1) Hamdi S. Aldharab, Aligarh Muslim University, India.

(2) Ibrahim Sufiyan, Federal Polytechnic Nasarawa, Nigeria.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/56773>

**Original Research Article**

**Received 10 March 2020**

**Accepted 14 May 2020**

**Published 04 June 2020**

## **ABSTRACT**

Basin is main carrying surface of rainwater collected from basin area and moves to main drain as river system. The basin surface characteristic governs the movement, storage and drainage in basin due to rainwater. Morphometric analysis is a quantitative description of a basin with important aspect of the basin character. Godawari basin divided into five sub-basins. *Indrawati* is main tributary river of Godawari originated from Mardiguda, Dandakarnya range of Kalahandi district, Odisha. The basin includes Bastar Craton covering a large area (>2105 km<sup>2</sup>) in Central India, bounded by two Mesoproterozoic mobile belts, the Eastern Ghats Mobile Belt (EGMB) to the east, Satpura Mobile Belt to the northwest, and two major Palaeozoic tectonic features the Godavari Graben to the southwest and the Mahanadi Graben in the northeast. The basin has been extracted using Advanced Space borne Thermal Emission and Reflection Radiometer (ASTER GDEM) 30 m (USGS, 2012) images by giving pour points and it has been projected on WGS 1984 UTM zone 45 for the morphometric analysis was used. Morphometric analysis expresses various aspects of basin characters which are 8 stream order (u), 85636 stream number, 42259.27 Stream length (Lu)

\*Corresponding author: E-mail: [adi19agro@gmail.com](mailto:adi19agro@gmail.com);

(km), 0.493 Mean Stream length (Lsm), 3.144 Bifurcation ratio (Rb), 0.002 Length of overland flow (Lg), 0.110 drainage density (Dd), 3.088 Stream frequency (Fs), 1.220 Form factor (Ff), 2.634 Circulatory ratio (Rc), 0.0040 Elongation ratio (Re), 6.348 Lemniscate method (k) and 1.758 Relief ratio of *Indrawati* basin.

**Keywords:** *Morphometric analysis; gis tools; Indrawati River; drainage pattern; basin characteristics; land use covers.*

## 1. INTRODUCTION

The major resources of water on earth is rainfall which directly falls on earth surface and further movement of the water preferably depends on surface morphology either through free movement or obstacles on the way of flow. How much quantity of rainfall received on surface is not so important but the movement of rainwater decides the behaviour of water with land surface. These all aspects make vital the study of morphology. The drainage basin is a basic unit in fluvial geomorphology expresses relationships between landforms and the processes which modify surface. The study of the geometry on the basin through the changes shows response to processes becoming a major part of geomorphology. Morphometric analysis is a quantitative description of a basin with important aspect of the basin characteras given by Strahler [1]. The term 'morphometry' indicates the meaning 'measurement of form' derived from morpho (form) and metry measurement [2]. The morphometric analysis involves linear, relief, aerial and gradient of channel networks and slopes of the basin [3,4,5]. The remote sensing technique is an authentic way for basin analysis with the satellite images provides the opportunities to get a realistic view on large area. Digital Elevation Model (DEM) in the field of GIS provides the dimensional evaluation of the earth surface. Advanced Space borne Thermal Emission and Reflection Radiometer (ASTER) provide the data of DEM and for hydrological analysis and extraction of stream network of a drainage basin [6]. The DEM is used assuming that the water will flow from higher to lower elevation using steep differences, will also produce a stream extraction model with a thematic layer of aspect, slope, relief, and drainage density, stream frequency [7]. GIS is used for the assessment of various basin parameters, is a reliable and authentic technique for the interpretation and analysis of spatial parameters of the drainage basin [8].

Trend detection of stream flow and rainfall is an active area of interest for both hydrology and

climatology in consideration of climate scenario. Stream movement is known to reflect on integrated response of the entire river basin, while rainfall serves as one of the major inputs into the stream processes. Hydro-climatic condition at larger scale may lead to loss of spatial information of the variable levels. Therefore, it seems to be logical to analyse hydro-climate variables at a small scale such as river basin or sub river basin scales [6]. The trend of stream flow and their association with rainfall and other influencing factor at river basin scale. Therefore, this study was carried out to investigate the stream trend in *Indrawati* river sub basin and possible linkages for the observed changes in stream flow with rainfall. Detailed knowledge in hydro-climatic trends is essential in proper planning of water management and thereby the agricultural planning in regions of basin.

### 1.1 Description of the Study Area

Godawari basin divided into five sub-basins namely Godawari lower, Godawari upper, Godawari middle, *Indrawati*, Manjra, Pranhita and others, Wardha and Weinganga which are jointly responsible for total basin flow, the highest and lowest area covers under Weinganga (49695.40 km<sup>2</sup>) and Godawari upper (22443.20 km<sup>2</sup>) sub-basin, respectively. Among these sub-basins, *Indrawati* sub basin covers utmost 38306.10 km<sup>2</sup>, with 21 dam and 15 reservoirs constructed on the way of flow, it includes 60 watersheds having maximum, minimum and average area of 992.92 km<sup>2</sup>, 648.95 km<sup>2</sup> and 343.14 km<sup>2</sup> are, respectively. *Indrawati* is main tributary river of Godawari originated from Mardiguda, Dandakarnya range of Kalahandi district (Odisha state) and flow through states Odisha, Chhattisgarh and Maharashtra covering 535 km while travelling from origin to confluence point with Godawari. *Indrawati* forms beautiful water fall at Chitrakot near Jagdalpur (Chhattisgarh) and hydro-power project in upper *Indrawati* at Mukhiguda, Kalahandi. It flows east to west of southern Chhattisgarh state crossing Koraput district of Odisha from origin point as

perennial river and forms large sub basin area of Godawari basin in the Bastar plateau at traverses in a east-west direction and assumed as life line of Bastar. Indarwati river runs zigzag manner throughout its flow covering six districts of Chhattisgarh except Sukma district, 2 districts of Odisha in the sub basin of Godawari into west, while travelling 306.75 km from its origin to joining point of the Godawari River, the whole basin extends between 20°35'59.30" N to 18°26'10.744" N latitude and 83°5'44.349" to 80°4'34.328" E longitude (Fig. 1). It is one of the important tributaries of the Godawari River in eastern part serving large rain fed area for prosperity and meets with the Godawari at Sukma district. Elevation variation of the basin is from 122 to 688 metres above mean sea level from origin to confluence point and drain area covered 38306.10 km<sup>2</sup>. The river generally static in nature but during rainy season, flood flow covers large area of basin due to common drainage of stream into river. Undulating land surface induce erosion and sedimentation in the river. The average bed slope of *Indrawati* River is 26.33 m/km, but its zigzag flow forms small and sharp tracks in medium size hillocks. The relief in the basin is formed due to the morphometric origin, and it is the predominance of shale, sandstone, Arcosic (Abujhmar group) ferruginous formations, which induces erosion with different magnitude of erodability. There is presence of some thrusts in the form of recumbent foldlike Chitrakot waterfall. This whole area is part of Eastern Ghats and central India.

## 2. GEOLOGY OF BASTAR

The Bastar Craton covers a large area (>2105 km<sup>2</sup>) in Central India, bounded by two Mesoproterozoic mobile belts, the Eastern Ghats Mobile Belt (EGMB) to the east, Satpura Mobile Belt to the northwest, and two major Palaeozoic tectonic features the Godavari Graben to the southwest and the Mahanadi Graben in the northeast. The craton comprises a number of linear supra crustal belts such as the Bengpal-Sukma Belt in the south, Kotri-Dongargarh Belt in the centre and north, the Amgaon Belt in the west and the Sasur-Chilipi Belt in the north and the Sonakhan Belt in the east, near Bhopalpatnam, located on the northern shoulders of the Godavari Rift. The other important granulite belts in the region include: the Kondagaon Belt (70.11 km) at the northern margin of the Indravati Basin and the Konta granulite Belt in the south of the Sukma Basin, sub-parallel to the Eastern Ghats Mobile Belt

(EGMB) [9,10,11,12,13,14]. A distinct North-South trending linear belt of banded iron formation (Bailadila) is well developed in the centre of the region overlying rocks of the Bengpal Group [15]. It was also reported similar formations by Mishra et al. [16] in lithological association and tectono-metamorphic history with those of the iron formations of other belts in the region. In the southern part of Bastar Craton, several episodes of mafic magmatism include mafic dykes and mafic volcanic rocks trending in a NW-SE direction, that occur mostly in the west of the Indravati, Sukma basins and around the Bailadila Iron Formation. Some of the shear zones cut across the sedimentary basins of Indravati, Sukma, Abujhmar, Khariar and Amphani. Karimnagar-Godavari Shear Zone (KGSZ) This zone, trending NW-SE extends for more than 200 km in length and a few kilometres wide passing through the Pakhal Basin sediments and older metamorphic rocks shows a dextral sense. These formations north of the Abujhmar Basin are strongly deformed and show a clear-cut north-south sinistral displacement suggesting major continental extensional tectonics during the Palaeoproterozoic, which was probably responsible for the evolution of the sedimentary basins. Several mafic dykes are also observed to the west and south of the *Indravati* Basin (Fig. 2).

## 3. METHODS AND MATERIALS FOR THE STUDY

The basin has been extracted using ASTERGDEM (USGS, 2012 [17]) images and by giving pour points and it has been projected on WGS 1984 UTM zone 45 for the morphometric analysis was used. The freely available Advanced Spaceborne Thermal Emission and Reflection Radiometer (ASTER) 30 m GDEM (version 2) of one arcsecond (arcsec) grid was downloaded for the study area in GeoTIFF format with geographic coordinates (USGS, 2012). The slope and relief of the basin were studied using digital elevation model data (DEM) available at USGS Earth Explorer for river basin and construct the stream analysis. The law of Strahler's method of stream ordering has been adopted for ordering the streams [18]. The result of the stream order has been converted from raster to feature. A manual correction has been done by merging the stream of the same order and assigning these as same order streams [19]. The stream order, number of stream, stream length, mean stream length, bifurcation ratios, relief ratio, drainage density, stream frequency,

form factor, circulatory ratio and elongation ratio were estimated using the mathematical formula given in Table 1 with the ASTERGDEM satellite images. ArcGIS software has been used for the entire image processing and calculation of all parameters of morphometry study [20,21,22,23].

To show the changes in the hydrology of the Indrawati river basin, the trend of water flow discharge has been analysed. The fluctuations in the trend exert the peak flow of the river. The data has been collected from several research paper and available reports.





Fig. 1. Location map of study area



Geological map of the Bastar Craton. Inset in the upper part shows the location of the Bastar and other cratons in India, modified after Ramachandra et al. (2001); (1) Chilpi Belt, (2) Sausar Belt, (3) Amgaon Belt, (4) Kotri Linear Belt, (5) Bengal Belt, (6) Sukma Belt, (7) Sonakhan Belt, Chh = Chhattisgarh Basin, Kha = Kharar Basin, Amp = Amphani Basin, Ind = Indravati Basin, Abu= Abujhmar Basin, Suk = Sukma Basin, Pak = Pakhal Basin. Figure 3. SRTM-DEM image of southern Bastar Craton (studied area).

Fig. 2. Geological map of the region  
Courtesy: Yellappa et al., (2011)

**Table 1. Mathematical formulae for analysis of basin characteristics**

S. No.	Parameters	Formulae	References	Results
1	Stream order (u)	Hierarchical rank	Strahler [18]	8
2	Stream number	Based on stream order	Strahler [1]	85636
3	Stream length (Lu) (km)	Length of stream	Horton (1945)	42259.27
4	Mean Stream length (Lsm)	$Lu=SL/Nu$	Strahler [1]	0.493
5	Bifurcation ratio (Rb)	$Rb=Nu/Nu+1$	Schum [24]	3.144
6	Length of overland flow (Lg)	$Lg=1/Dx2$	Horton (1945)	0.002
7	Drainage density (Dd)	$Dd=\sum Lu/Au$	Horton (1945)	1.041
8	Stream frequency (Fs)	$D=\sum Nu/Au$	Horton (1945)	3.088
9	Form factor (Ff)	$Ff=A/LP^2$	Horton (1945)	1.220
10	Circulatory ratio (Rc)	$Rc=4\sqrt{A}/P^2$	Strahler [1]	2.634
11	Elongation ratio (Re)	$Re=D/Lb$	Schum [24]	0.901
12	Leminscate method (k)	$K=L2/4A$	Chortey et al. (1957)	6.348
13	Relief ratio	$Rh=H/Lb$	Schum (1963)	1.758

**4. RESULTS AND DISCUSSION**

**4.1 Indrawati River Basin: Linear Aspects of Morphometry**


**4.1.1 Stream order (u)**

Stream order is defined as a measure of the position of a stream in the hierarchy of tributaries [25,26,18], have developed stream ordering system. The *Indrawati* river, which is a main stream of the study area, is the eight stream orders (Based on Strahler’s method of stream order). The variation extends order and size of the tributary basin is due to physiographic and structural conditions prevailing in the basin. The river basin develops over the shale, sandstone, gneiss and slate as major rock and thereby the number and order of streams are more. Total stream number is 85636 showing value 31.023 and 8179.61 as minimum and maximum, respectively from total mean length of 0.493 km


and standard deviation of 692.83 while study of the *Indrawati* basin. The highest number of first order streams are 600150 followed by second order stream of 308424 and lowest was 3035 expressing initiation and formation of new stream apparently.

**4.1.2 Stream number**


The total stream segments in each order are known as stream number. R. E. Horton states that the number of stream segments of each order from an inverse geometric sequence with the order number. The number of streams orders has been counted with the help of ArcGIS software. A total number of streams in the basin are 85636. The underlying tender rocks and undulating plain favour a large number of streams further more initiation. In the *Indrawati* basin, the first order of the streams is associated with the highest number of streams and remaining are gradually decreased.



**a. Basin coverage of Indrawati River**



**b. River basin of Indrawati**



**Fig. 3. Basin characteristic map of Indrawati River**

#### 4.1.3 Stream lengths (Lu)

The total streams lengths of various orders have been measured with the help of calculate geometry of GIS tools. It is indicative of the chronological development of streams and derived from source to mouth of the streams [8]. The total length of the *Indrawati* river is about 535 kms, and the length of the total streams of the *Indrawati* river basin is 42259.27 km as length of streams increase flow coverage automatically increased.

#### 4.1.4 Mean stream length (Lsm)

The total length of stream measured and total stream length is divided by the total number of segments in that order. There is the relationship between mean stream length and basin order *i.e.* mean stream length increases with successive increasing orders. It is related to the size of the drainage network [1]. Mean Stream Length of the *Indrawati* River basin is 0.493 km and is quite small while flowing takes many things away.

#### 4.1.5 Bifurcation ratio (Rb)

It is the ratio between the numbers of stream segments of any given order to the number the next higher order. The bifurcation ratio ranges in between 3 to 5. The irregularities of bifurcation ratio depend on the geological and the lithological development of the drainage basin. The lower value of the bifurcation ratio reveals that the basin has suffered less structural disturbances and the higher values of the bifurcation ratio indicate strong structural control on the drainage pattern [1]. Bifurcation ratio having >10 indicates the drainage basin developed over the easily erodible rocks [27].

The average value of bifurcation is 3.144 and values of the bifurcation ratio of different order ranges from 2 to 5.26 due to large area coverage and topography of plateau as district demarcation lines by hillocks and hills favours less structural disturbances.

#### 4.1.6 Length of overland flow (Lg)

Normally, a higher value of Lg represents low relief and whereas a low value of Lg is an indicative of high relief. The length of the overland flow of *Indrawati* River is 0.002 km/km<sup>2</sup>. The value is usually low that indicates the basin is consists of high relief and slope, and thus rainwater enters the stream very quickly, meanwhile entry of rain water in stream way, the flow washes a lot of materials on the way.

### 4.2 Aerial Aspects of Morphometry

#### 4.2.1 Drainage density (Dd)

Drainage density (Dd) is the ratio of the total length of the streams in a given drainage basin and the area of that drainage basin [26] and envisaged that the value of the drainage density ranges from 0.93 km/km<sup>2</sup> to 1.24 km/km<sup>2</sup> in the steep impervious area of the high precipitation region and zero for the permeable basin with high infiltration rate. Langbien considered that the drainage 4 density is ranged from 0.55 to 2.09 for the humid region with the average drainage density of 1.03. Low drainage density indicates permeable subsurface and coarse drainage [8]. Drainage density of the *Indrawati* River Basin is 1.041 km/km<sup>2</sup>, and it means weak subsurface material and belongs to medium texture of the basin with steep impervious.



#### 4.2.2 Stream frequency (Fs)

It is expressed as the ratio of a total number of channels cumulated for all orders within a given drainage basin and the area of the drainage basin [26]. It indicates the index of various stages of landscape development and depends on the nature and amount of rainfall, the nature of rock and soil permeability of the region. Stream frequency of the *Indrawati* River basin is  $3.088/\text{km}^2$ , which indicates high relief and high infiltration of the basin because of compartmentation. Hence, both the drainage density and the stream frequency are the parameters for measuring the spacing of stream channel in the basin.

#### 4.2.3 Form factor (Ff)

The value of the form factor varies from 0 (highly elongated shape) to 1 (perfect circular shape). The value of form factor of the *Indrawati* River basin is 1.22 which depicts that the basin form belongs to perfectly circular shape and cause flood situation in the district in rainy season, and it is the indication of high peak flow of the basin.

#### 4.2.4 Circulatory ratio (Rc)

The Circulatory ratio is to provide a quantitative index of the shape of the basin. Circulatory ratio is the ratio of basin area to the area of the circle having the same parameter as the basin. A circular basin has a maximum efficiency of the movement of runoff, where as an elongated basin has the least frequency. This information is very significant for forecast of drainage discharge, particularly in a time of the flood [28]. The value of the circulatory ratio is 2.364 which explain high circular shape and high run off flow of discharge of the *Indrawati* River basin while heavy down pour.

#### 4.2.5 Elongation ratio (Re)

This ratio usually runs from 0.6 to 1.0 over a broad range of climatic and geological type. Values near 1.0 are a typical region of low relief while values from 0.51 to 0.94 are associated with strong relief and steep ground slope. The ratio is a meaningful index for classifying drainage basins into varying shapes as circular (above 0.9), oval (0.8-0.9), less elongated (0.7-0.8) and elongated (below 0.7). The value of elongation ratio of the *Indrawati* river basin is 0.901. So, the basin has a circular in shape and

associating strong relief of surface and low peak flow.

#### 4.2.6 Lemniscate method (K)

It is the comparison of the basin with the lemniscate curve. The high value of K indicates elongated shape and on the contrary, a little value indicates the circular shape of the basin. The K value of *Indrawati* river basin is 6.348 that depicts that the river is a perfectly circular and it indicates high peak flow of discharge of the basin. *Indrawati* basin has higher K means high order stream is present and further it will increase with progress.

### 4.3 Relief Aspects of Morphometry

#### 4.3.1 Relief ratio

The Relief ratio is the ratio between the total relief of a basin and the longest dimension of the basin parallel to the main drainage line [24] and a dimensionless height-length ratio equal to the tangent of the angle formed by two planes intersecting at the mouth of the basin, one representing the horizontal, the other passing through the highest point of the basin. It denotes the overall steepness of a drainage basin and is an indicator of the intensity of degradational processes operating on slopes of that basin. The relief ratio of the river basin is 1.758 that indicates this basin is composed of more resistant in flow by hindrance of hills and hillocks of basin, and the basin is under intense relief and steep slope.

#### 4.3.2 Drainage patterns of the *Indrawati* river basin

In the *Indrawati* river basin, Tree-like dendritic type of drainage pattern mainly develops over the existing rocks apart from the dendritic pattern also exhibits the Herringbone, parallel and centrifugal type of drainage pattern. The parallel drainage pattern, where the control of the regional slope, parallel faults and lineaments predominate is discernible too. In this drainage pattern, parents and tributaries flow in right angle to meet. In some peak of the ridges, Centrifugal drainage pattern, where streams radiate in all directions from a common centre, is significant feature of the hillocks.

#### 4.3.3 Long profile of the river

It is generally concave to the sky, but the slope is decreasing rapidly at first and then gradually

[29,25]. The River originates at 688 m above mean sea level and runs 535 km. The long profile of a river is dependent upon the lithology of the underlying structure, stream flow, discharge, the amount and texture of the channel load, flow resistance, flow velocity, width and depth of the channel, and regional slope [27]. The *Indrawati* River has a different channel slope and concavity. The regular concave profile is interrupted by the convex profile at about 20-30kms from the origin. The convex slope is between 1500-3000 metres due to plateau shortly change in movement on structural differences and concave in between 1000-2000 metres above meaning sea level having sharp turn due to landforms. The irregularities are due to the presence of the more resistance hillocks and change in slope (Fig. 3).

**4.3.4 Cross profiles of the river**

The cross profile varies from the narrow, steep-sided trenches to gentle open form, ultimately depends on the resistance offered by the undulating plateau slopes and the erosive capacity of the water during rain. Cross profile of the *Indrawati* River in different regimes exerts the processes and dominant controlling factor. Most of the cross profiles are ‘U’ shaped in nature and it represents lateral cutting rather than down erosion. Near the source, the dominance of hills and river has more stream power for lateral cutting, thus flat ‘U’ shaped profile is significant (Fig. 4).


**4.3.5 Recent hydrological changes in *indrawati* river basin**

The significant changes have been observed in the *Indrawati* River Basin mainly due to climatic and anthropogenic causes. The climatic


variability leads to heavy and erratic rainfall of the region induces soil loss and thereby retreat of sedimentation of the river basin is discernible and consequences of the events are shallow depth, diversion, flood with small quantum of rainfall washing out top agriculturally important soil. The source region of the *Indrawati* river, is eastern ghats experienced with abrupt changes. Deforestation in the basin area also increases the erosion capacity of the rivers along with tributaries. As a result, basin might be more prone to soil and water loss in coming years (Fig. 3).

**4.3.6 Temporal changes in discharge of *indrawati* river**

A tendency of decrease in average annual and seasonal discharge of the *Indrawati* River has been experienced. It is significant from the several studies that the decline of discharge is also experienced. The flow characteristic is controlled by variability of rainfall. The gradual decrease of discharge in winter on wards is noticed due to less flow and uneven distribution between states like Odisha and Chhattisgarh during lean periods. Simultaneously, the irregularity of the Monsoon disrupts the average discharge of the river in a lesser extent. The present study on drainage basin has been done on the basis of geo-morphological perspective. Study of the morphometry is concerned with the genesis, development and distribution of landforms [30]. An important property of the drainage basin is its hierarchical in nature, each tributary in a drainage system has its basin area. In any drainage basin, there are several variables which always interacting with other variables and give birth to particular types of landforms.



**Fig. 4. Long profile of Indrawati basin**



**Fig. 5. Cross profile of Indrawati basin**



## 5. CONCLUSION

The study reveals that the drainage basin is associated with the early stage of drainage development, and high order of stream dominates the river basin. The circular shape of the basin reveals high runoff and low infiltration capacity and uncontrolled flow of river water with disturbing dependent factors of basin. Recently, several anthropogenic activities like house construction on banks, fast deforestation, siltation and climatic variability (erratic rainfall and rising temperature) caused the several hydrological changes in the *Indrawati* River basin. Recent changes in discharge can affect the aggradational and degradational processes that also affect the overall welfare of society.

## ACKNOWLEDGEMENT

The authors are highly thankful to All India Coordinated Research Project on Dry land Agriculture, S.G. CARS IGKV, Raipur for their necessary inputs and guidance while study and analysis of observed data.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Strahler AN. Quantitative geomorphology of drainage basins and channel networks In V T Chow Edition. Handbook of Applied Hydrology. 1964:439-476.
2. Pandey PK, Das SS. Morphometric analysis of Usri River basin, Chhotanagpur Plateau, India, using remote sensing and GIS. Arab Jour Geosci. 2016; 9(3):240.
3. Nag SK, Chakraborty S. Influence of rock types and structures in the development of drainage network in hard rock area. Journal of the Indian Society of Remote Sensing. 2003;31:25-35.
4. Avinash K, Jayappa KS, Deepika B. Prioritization of subbasins based on geomorphology and morphometric analysis using remote sensing and geographic information system (GIS) techniques. Geocarto Int. 2011;26(7):569-592.
5. Magesh NS, Jitheshlal KV, Chandrasekar N, Jini KV. GIS based morphometric evaluation of Chimmini and Mupily watersheds parts of Western Ghats Thrissur, India. Earth Science Informatics. 2012;5:111-121.
6. Bindu G, Neelkantan R, Ragunath R. Assessment of morphometric characteristics of Chittar river basin, Triruvantpuram district, Kerala: A remote sensing and GIS based study. Indian J Geomorphol. 2012; 1:127-132.
7. Chandrasekar N, Magesh. GIS model-based morphometric evaluation of Tamiraparani sub basin Tirunelveli, India. Arab Journal Geo Science. 2014;7:131-141.
8. Magesh NS, Jitheshlal KV, Chandrasekar N, Jini KV. Geographical information system-based morphometric analysis of Bharathapuzha river basin Kerala, India. Applied Water Science. 2013;3:467-477.
9. Mishra VP, Dutta NK, Kanchan VK, Vasta US, Guha K. Archaean granulite and granite gneiss complexes of Kondagaon area, Bastar District, M.P. Records of Geological Survey of India. 1984;113(6):150-158.
10. Mishra VP, Dutta NK, Guha KK, Kumaran K, Ramachandra HM. Geology of Pakahanjur area, Bastar and Rajnandgaon district, Madhya Pradesh, Raipur, India. Geological Survey of India (unpub. report). 1981;76.
11. Bali R, Agarwal KK, Ali SN, Rastogi SK, Krishna K. Drainage morphometry of Himalayan Glacio-fluvial basin, India: hydrologic and neotectonic implications. Environ Earth Sci. 2012;66(4):1163-1174.
12. Panda B, Venkatesh M, Kumar B, Anshumali. A GIS-based approach in drainage and morphometric analysis of Ken River basin and sub-basins, Central India. J Geol Soc India. 2018;93(1):75-84.
13. Santosh M, Yokoyama K, Acharya SK. Geochronology of tectonic evolution of Karimnagar-Bhopalpatnam granulite belts, Central India. Gondwana Research. 2004; 7(2):501-518.
14. Chorley RJ. Illustrating the laws of morphometry. Geol Mag. 1957;94(2):140-150.
15. Crookshank H. Geology of southern Bastar and Jeypore from the Bailadila range to Eastern Ghats. Geological Survey of India, Calcutta, India, Memoir. 1963;87:1-150.
16. Mishra VP, Singh P, Dutta NK. Stratigraphy, structure and metamorphic history of Bastar Craton. Records of Geological Survey of India. 1988;117(3-5):1-26.
17. USGS. Global data explorer, 2012. Available: <http://gdex.cr.usgs.gov/> (Accessed 11.12.12)

18. Strahler AN, Dynamic basis of geomorphology Bull. Geological Science America. 1952;63:923-938.
19. Dave M. ASTER Global Digital Elevation Model Version 2-Summary of Validation Results. US Geological Survey. 2011;28.
20. Vittala SS, Govindaiah S, Gowda HH. Morphometric analysis of subwatersheds in the pavagada area of Tumkur district South India using remote sensing and gis techniques. Journal of the Indian Society of Remote Sensing. 2004;32:351-362.
21. Angillieri MYE. Morphometric characterization of the Carrizal basin applied to the evaluation of flash floods hazard, San Juan, Argentina. Quat Int. 2012;253:74-79.
22. Patel PP, Sarkar A. Terrain characterization using SRTM Data. Journal of Indian Society of Remote Sensing. 2010;38:11-24.
23. Biswas SS. Role of human induced factors Soil erosion and Climatic variability in changing the fluvial system: A case study of the Barakar River Basin India Unpublished M Phil Dissertation. Jawaharlal Nehru University New Delhi; 2014.
24. Schumm SA. The evolution of drainage system and slopes in badlands at Perth Amboy, New Jersey. Geological Science America Bull. 1956;67: 597-646.
25. Leopold LB, Wolman MG, Miller JP. Fluvial Processes in Geomorphology New York: 1964; Dover Publication.
26. Horton RE. Drainage basin characteristics Trans Am Geophys Union. 1932;13:350-361.
27. Kale VS, Gupta A. Introduction to Geomorphology (First impression edition). Hyderabad: University Press; 2010.
28. Chitra C, Alaguraja P, Ganeshkumari K, Yuvaraj D, Manivel M. Watershed characteristics of Kundah sub basin using remote sensing and GIS techniques. Int J Geomat Geosci. 2011;2(1):311.
29. Knighton D. Fluvial Forms and Processes: A New Perspective London: Arnold; 1998.
30. Kumar B, Venkatesh M, Tripathi A, Anshumali. A GIS-based approach in drainage morphometric analysis of Rihand River Basin, Central India. Sustain Water ResourManag. 2017;4(1):45-54.

© 2020 Pradhan et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:  
<http://www.sdiarticle4.com/review-history/56773>*