



Efficacy of 2,4-D Choline as Influenced by Weed Size in the Texas High Plains

Misha R. Manuchehri^{1*}, Peter A. Dotray² and J. Wayne Keeling²

¹Department of Plant and Soil Sciences, Oklahoma State University, 371 Agricultural Hall, Stillwater, OK 74078, US State.

²Texas A&M Agri Life Research & Extension Center, Lubbock, TX 79403, US State.

Authors' contributions

This work was carried out in collaboration among all authors. Author MRM established the trials, collected data, performed the statistical analysis and wrote the first draft of the manuscript. Author PAD designed the study, wrote the protocol and co-wrote the manuscript. Author JWK contributed to the protocol assisted with data collection and reviewed the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2019/v35i230199

Editor(s):

(1) Dr. Renata Guimarães Moreira-Whitton, Professor, Departamento de Fisiologia - Instituto de Biociências-USP, Cidade Universitária, Brazil.

Reviewers:

(1) Renata Kieloch, Institute of Soil Science and Plant Cultivation, Poland.

(2) Young Jasper, Rivers State University, Nigeria.

(3) Toungos, Mohammed Dahiru, Adamawa State University, Nigeria.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/48301>

Received 07 February 2019

Accepted 13 April 2019

Published 27 April 2019

Original Research Article

ABSTRACT

Aim: Postemergence timing trials based on weed size were conducted near Lubbock, TX to assess the effectiveness of 2,4-D choline + glyphosate on control of Palmer amaranth (*Amaranthus palmeri* S. Wats.), Russian-thistle (*Salsola tragus* L.), and kochia (*Kochia scoparia* L.) at three growth stages (3 to 5 cm, 10 to 15 cm, and 20 to 30 cm).

Study Design: All trials were arranged in a randomized complete block design with four replications.

Place and Duration of Study: Field experiments were conducted in 2013, 2014, and 2015 in Lubbock, TX at the Texas A&M AgriLife Research and Extension Center near Lubbock, TX.

Methodology: Herbicide treatments consisted of a single postemergence application of 2,4-D choline + glyphosate at two rates, 2,4-D choline + glyphosate at two rates + glufosinate, 2,4-D choline + glyphosate + S-metolachlor, 2,4-D choline + glyphosate + acetochlor, 2,4-D choline + glufosinate, glyphosate, or glufosinate.

*Corresponding author: E-mail: misha.manuchehri@okstate.edu;

Results: The greatest level of weed control for all three weed species was achieved at the 3 to 5 cm timing; however, weed size was most critical for Palmer amaranth and Russian-thistle compared to kochia. Averaged over all three years, Palmer amaranth control decreased from 93 to 74% when evaluated 21 days after treatment following applications that included 2,4-D choline when applied to plants 3 to 5 and 10 to 30 cm, respectively. For Russian-thistle, control decreased from 98 to 78% when evaluated 21 days after treatment following treatments that included 2,4-D choline when applied to plants 3 to 5 and 10 to 30 cm, respectively. For kochia, control decreased from 98 to 84% when evaluated 21 days after treatment following treatments that included 2,4-D choline when applied to plant 3 to 5 and 10 to 30 cm, respectively.

Keywords: Application timing; growth stage; tank mixtures.

1. INTRODUCTION

Effective, economical, and sustainable weed management is crucial to a profitable cotton production system. Weeds decrease cotton lint yield and quality by competing for nutrients, water, and light [1]. Palmer amaranth (*Amaranthus palmeri* S. Wats.), Russian-thistle (*Salsola tragus* L.), and kochia (*Kochia scoparia* L.) are among the most difficult-to-control weeds in Texas High Plains cotton. Palmer amaranth was ranked as the most troublesome cotton weed in the southern United States in 2009, occurring in nine of ten states surveyed [2]. It also has become one of the most economically damaging glyphosate-resistant weed species in the United States [3]. Russian-thistle, a C4 summer annual broadleaf weed that is prevalent in the western United States, is extremely competitive due in part to its aggressive root system [4,5]. Early seedling emergence, tolerance to drought, heat, and salinity, hermaphroditic flowers that are out-crossed and self-fertile, and wind-mediated pollen dispersal also contribute to its competitiveness [6-8]. The competitiveness of kochia, also a troublesome C4 summer annual broadleaf weed in croplands and non-croplands over the Great Plains of North America, is attributed to its early seedling emergence, rapid growth rate, heat and salt tolerance, prolific seed production, and long-distance seed dispersal by tumbling [9-15].

Additionally, all three of these weeds have developed resistance to critical herbicides modes of action. In the United States, Palmer amaranth has even evolved resistance to multiple herbicide modes of action such as EPSP synthase inhibitors (Group 9), ALS inhibitors (Group 2), HPPD inhibitors (Group 27), PPO inhibitors (Group 14), microtubule assembly inhibitors (Group 3), and photosystem II inhibitors (Groups 5-7) [16]. Russian-thistle and kochia populations resistant to ALS and/or EPSP synthase inhibitors have been documented and kochia populations resistant to synthetic auxins and photosystem II inhibitors also have developed [17-

18]. Therefore, the list of available modes of action to control these weed species in cotton is limited; however, an additional option became available with the release of Enlist™ technology in cotton in 2016. Enlist™ technology utilizes cotton tolerance to 2,4-D choline, glyphosate, and glufosinate. Cotton tolerant to 2,4-D choline was conferred by the insertion of a gene (AAD-12) that codes for an aryloxyalkanoate dioxygenase enzyme [19]. Plants transformed to include this gene can metabolize certain auxin herbicides, including 2,4-D, to a nonlethal form [20]. Enlist™ cotton provides growers with a new tool to effectively manage Palmer amaranth, Russian-thistle, kochia, and other difficult-to-control weeds in Texas High Plains cotton.

Weed size at the time of application [21-24] and tank-mix combinations [25-27] are two factors that often impact the success of a herbicide. The importance of weed size at the time of 2,4-D application has been well-documented (Everitt and Keeling 2007; Siebert et al. 2004). Therefore, weed size should be considered when making 2,4-D choline applications. The objective of this research was to evaluate the effectiveness of mixtures of 2,4-D choline with glyphosate, glufosinate, S-metolachlor, and/or acetochlor on control of Palmer amaranth, Russian-thistle, and kochia at various growth stages.

2. MATERIALS AND METHODS

2.1 Experimental Site

Non-crop, field experiments were conducted in 2013, 2014, and 2015 in Lubbock, TX at the Texas A&M AgriLife Research and Extension Center (33.415319°N, -101.483274°W, elevation 1,001 m). The soil type was an Amarillo fine sandy loam (fine-loamy, mixed, superactive, thermic Aridic Paleustalfs) with less than 1% organic matter and pH of 7.5. All studies were arranged in a randomized complete block design with four replications. Individual plots were 3.0 m

Table 1. Dates of palmer amaranth, Russian-thistle, and Kochia applications at several weed sizes near Lubbock, TX

Weed size (cm)	Palmer amaranth			Russian-thistle/kochia		
	2013	2014	2015	2013	2014	2015
3 to 5	June 14	-	June 3	April 13	May 15	-
10 to 15	June 27	July 23	June 18	May 14	-	April 21
20 to 30	July 8	August 19	June 24	June 13	June 3	May 4

Table 2. Herbicide treatments and application rates for 2013, 2014, and 2015 application timing trials near Lubbock, TX

Herbicide common names	Brand names or designations	Application rates	Manufacturer
2,4-D choline + glyphosate	Enlist Duo™	1.64 or 2.19 kg ae ha ⁻¹	Dow AgroSciences, Indianapolis, IN, http://www.dowagro.com
Glufosinate	Liberty® 280 SL	0.59 kg ai ha ⁻¹	Bayer CropScience, Research Triangle Park, NC, https://www.cropscience.bayer.com
S-metolachlor	Dual MAGNUM®	1.09 kg ai ha ⁻¹	Syngenta Crop Protection, Greensboro, NC, https://www.syngenta.com
Acetochlor	Warrant®	1.26 kg ai ha ⁻¹	Monsanto Company, St. Louis, MO, http://www.monsanto.com
2,4-D choline	Enlist One™	1.07 kg ae ha ⁻¹	Dow AgroSciences
Glyphosate	Roundup PowerMAX®	1.12 kg ae ha ⁻¹	Monsanto Company

wide by 6.1 m in length. Annual rainfall was 292 mm in 2013, 460 mm in 2014, and 354 mm in 2015. No supplemental irrigation was provided. On average over all three years, there were approximately 1,200 Palmer amaranth, 30 Russian-thistle, and 10 kochia plants per plot (18.3 m²).

2.2 Experimental Design and Data Collection

In 2013, postemergence applications were made to 3 to 5, 10 to 15, and 20 to 30 cm Palmer amaranth, Russian-thistle, and kochia (Table 1). Weeds were susceptible to all herbicides (no resistance had developed); however, the Palmer amaranth population was in the initial stages of developing glyphosate resistance. In 2014, applications were made to 10 to 15 and 20 to 30 cm Palmer amaranth and 3 to 5 and 20 to 30 cm Russian-thistle and kochia. In 2015, applications were made to 3 to 5, 10 to 15, and 20 to 30 cm Palmer amaranth and 10 to 15 and 20 to 30 cm Russian-thistle. Kochia was not evaluated in 2015 as a late freeze eliminated most of the populations at this location. The nontreated control did not receive a herbicide application. All applications were made at 4.8 km per hour with a CO₂-pressurized backpack sprayer equipped with AIXR11002 spray tips (TeeJet® Technologies, Glendale Heights, IL) calibrated to

deliver 140 L ha⁻¹ at 205 kPa. No adjuvants were included with any application.

Treatments consisted of a single postemergence application of 2,4-D choline + glyphosate at two rates, 2,4-D choline + glyphosate at two rates + glufosinate, 2,4-D choline + glyphosate + S-metolachlor, 2,4-D choline + glyphosate + acetochlor, 2,4-D choline + glufosinate, glyphosate alone, or glufosinate alone. Herbicides and application rates are listed in Table 2. Visual control estimates were recorded 14, 21, and 28 days after treatment (DAT) using a scale of 0 to 100 percent, where 0 was no weed control and 100 was complete control. Foliar chlorosis, necrosis, tissue distortion, and plant stunting were considered when making visual control estimates.

2.3 Statistical Analysis

A univariate analysis was performed on all responses in order to test for stable variance. No data sets were transformed as transformation did not increase stabilization. Data sets were analyzed using PROC MIXED with pdmix 800 macro included [28] and treatments were separated by Fisher's Protected LSD at an alpha level of 0.05 using SAS 9.4 software (SAS Institute Inc., SAS Campus Drive, Cary, North Carolina 27513).

3. RESULTS AND DISCUSSION

For Palmer amaranth, Russian-thistle, and kochia control, trials were analyzed independently due to a significant year effect ($P < 0.05$) across all possible year combinations. Within year (2013, 2014, and 2015), 10 to 15 and 20 to 30 cm Palmer amaranth ratings were combined due to no difference in control based on weed height at application ($P > 0.05$). In 2013, 10 to 15 and 20 to 30 cm Russian-thistle and kochia ratings were combined due to no difference in control based on weed height at application ($P > 0.05$). All other control ratings were analyzed independently due to a significant weed height effect ($P < 0.05$).

3.1 Palmer Amaranth Control

In 2013, 2,4-D choline + glyphosate at 1.64 kg ae ha⁻¹, 2,4-D choline + glyphosate at 2.19 kg ae ha⁻¹, 2,4-D choline + glyphosate + S-metolachlor, and 2,4-D choline + glyphosate + acetochlor controlled 3 to 5 cm Palmer amaranth 95 to 98% 21 DAT while glufosinate alone controlled Palmer amaranth 58% (Table 3). For 10 to 30 cm Palmer amaranth, 2,4-D choline + glyphosate at 2.19 kg ha⁻¹, 2,4-D choline + glyphosate + S-metolachlor, and 2,4-D choline + glyphosate + acetochlor controlled Palmer amaranth the greatest (71 to 77%) while glufosinate alone provided the least control (5%).

In 2014, 2,4-D choline + glyphosate at 1.64 kg ha⁻¹ + glufosinate, 2,4-D choline + glyphosate at 2.19 kg ha⁻¹ + glufosinate, and 2,4-D choline + glyphosate + acetochlor controlled 10 to 30 cm Palmer amaranth 88 to 90% 21 DAT while glufosinate alone achieved the least control (54%) (Table 3). In 2015, 2,4-D choline + glyphosate at 2.19 kg ha⁻¹, 2,4-D choline + glyphosate + S-metolachlor, and 2,4-D choline + glyphosate + acetochlor achieved the greatest Palmer amaranth control (97 to 98%) while glyphosate alone achieved the least control (82%) (Table 3). 2,4-D choline + glyphosate at 2.19 kg ha⁻¹ and 2,4-D choline + glyphosate + S-metolachlor achieved the greatest control (86 to 87%) of 10 to 30 cm Palmer amaranth while glufosinate alone achieved the least control (44%).

3.2 Russian-thistle Control

In 2013 at 21 DAT, all treatments controlled 3 to 5 cm Russian-thistle 96 to 99% with the exception of glufosinate alone, which controlled this weed 75% (Table 4). All treatments achieved similar control (81 to 85%) of 10 to 30 cm

Russian-thistle with the exception of 2,4-D choline + glyphosate at 1.64 kg ha⁻¹ alone (70%), glyphosate alone (34%), and glufosinate alone (28%).

In 2014 at 21 DAT, all treatments controlled 3 to 5 cm Russian-thistle 95 to 100% and 20 to 30 cm Russian-thistle 71 to 76% with the exception of glyphosate alone (61%) and glufosinate alone (23%) (Table 4). Glyphosate alone controlled 3 to 5 and 20 to 30 cm Russian-thistle 69 and 61%, respectively, while glufosinate alone controlled 3 to 5 and 20 to 30 cm Russian-thistle 0 and 23%, respectively. In 2015, 2,4-D choline + glyphosate at 2.19 kg ha⁻¹ and 2,4-D choline + glyphosate + S-metolachlor achieved the greatest 10 to 15 cm Russian-thistle control (81 to 88%) 21 DAT while glufosinate alone achieved the least control (16%) (Table 4).

3.3 Kochia Control

In 2013 at 21 DAT, all treatments controlled 3 to 5 cm kochia 95 to 100% with the exception of glufosinate alone, which controlled this weed 79% (Table 5). All treatments achieved 76 to 90% control of 10 to 30 cm kochia with the exception of glufosinate alone (49%). In 2014, all treatments controlled 3 to 5 cm kochia 97 to 99% with the exception of glufosinate alone, which only controlled this weed 3% (Table 5). 2,4-D choline + glyphosate at 2.19 kg ha⁻¹, 2,4-D choline + glyphosate at 2.19 kg ha⁻¹ + glufosinate, 2,4-D choline + glyphosate + S-metolachlor, and 2,4-D choline + glyphosate + acetochlor achieved the greatest 20 to 30 cm kochia control (84 to 90%) while glufosinate alone controlled this weed the least (53%).

Similarly, Everitt and Keeling [29] found that 2,4-D at 0.56 and 1.12 kg ha⁻¹ controlled 3 to 8 cm horseweed at least 92% 28 DAT; however, reduced horseweed control was observed with these same rates of 2,4-D when applied to 10 to 15 cm and 25 to 46 cm-tall horseweed. A comparable response to 2,4-D also has been reported with other weed species such as red morningglory (*Ipomoea coccinea* L.) and dogfennel [*Eupatorium capillifolium* (Lam.) Small] [30]. Siebert et al. observed 100% control of 30 cm red morningglory; however, a 6 to 19% reduction in control was observed when 2,4-D was applied to 60 cm plants. Dogfennel control was reduced from 85 to 70 to 6% when applications of 2,4-D and dicamba were applied to plants 36, 72, and 154 cm in height, respectively [31].

Table 3. Influence of weed height and herbicide treatment on Palmer amaranth control 21 days after treatment in 2013, 2014, and 2015 near Lubbock, TXa

Treatments	Rate kg ae or ai ha ⁻¹	2013		2014	2015	
		3 to 5 cm	10 to 30 cm	10 to 30 cm	3 to 5 cm	10 to 30 cm
2,4-D choline + glyphosate	1.64	95 ab	66 b	80 bc	94 c	79 b
2,4-D choline + glyphosate	2.19	96 ab	77 a	82 b	97 ab	87 a
2,4-D choline + glyphosate + glufosinate	1.64 + 0.59	80 cd	48 cd	90 a	95 bc	66 c
2,4-D choline + glyphosate + glufosinate	2.19 + 0.59	78 d	54 c	90 a	95 bc	69 c
2,4-D choline + glyphosate + S-metolachlor	2.19 + 1.09	98 a	71 ab	79 bc	98 a	86 a
2,4-D choline + glyphosate + acetochlor	2.19 + 1.26	95 ab	72 ab	75 c	98 a	81 b
2,4-D choline + glufosinate	1.07 + 0.59	86 cd	44 d	88 a	94 c	64 c
Glyphosate	1.12	88 bc	63 b	74 c	82 d	59 d
Glufosinate	0.59	58 e	5 e	54 d	93 c	44 e

*a*Means within a column followed by the same letter are not significantly different according to Fisher's Protected LSD at $P < 0.05$. Data pooled for 10 to 15 cm and 20 to 30 cm Palmer amaranth control ratings within each year

Table 4. Influence of weed height at application and herbicide treatment on Russian-thistle control 21 days after treatment in 2013, 2014, and 2015 near Lubbock, TX.^a

Treatments	Rate kg ae or ai ha ⁻¹	Russian-thistle control					
		2013		2014		2015	
		3 to 5 cm	10 to 30 cm	3 to 5 cm	20 to 30 cm	10 to 15 cm	20 to 30 cm
2,4-D choline + glyphosate	1.64	96 ab	70 b	97 ab	71 a	100 a	70 cd
2,4-D choline + glyphosate	2.19	99 ab	85 a	100 a	75 a	100 a	88 a
2,4-D choline + glyphosate + glufosinate	1.64 + 0.59	96 b	84 a	95 b	73 a	99 ab	68 d
2,4-D choline + glyphosate + glufosinate	2.19 + 0.59	97 ab	85 a	99 a	73 a	100 a	74 cd
2,4-D choline + glyphosate + S-metolachlor	2.19 + 1.09	99 ab	83 a	100 a	75 a	100 a	81 ab
2,4-D choline + glyphosate + acetochlor	2.19 + 1.26	98 ab	81 a	98 a	76 a	100 a	70 cd
2,4-D choline + glufosinate	1.07 + 0.59	99 a	81 a	100 a	73 a	100 a	70 cd
Glyphosate	1.12	98 ab	34 c	69 c	61 b	99 a	75 bc
Glufosinate	0.59	75 c	28 c	0 d	23 c	98 b	16 e

^aMeans within a column followed by the same letter are not significantly different according to Fisher's Protected LSD at $P < 0.05$. In 2013, 10 to 15 and 20 to 30 cm Russian-thistle control ratings were combined due to no weed height effect ($P > 0.05$)

Table 5. Influence of weed height at time of application and herbicide treatment on kochia control 21 days after treatment in 2013, 2014, and 2015 near Lubbock, TX^a

Treatment	Rate	Kochia control			
		2013		2014	
		3 to 5 cm	10 to 30 cm	3 to 5 cm	20 to 30 cm
	kg ae or ai ha ⁻¹	%			
2,4-D choline + glyphosate	1.64	98 a	76 b	98 a	76 d
2,4-D choline + glyphosate	2.19	98 a	90 a	98 a	90 abc
2,4-D choline + glyphosate + glufosinate	1.64 + 0.59	95 a	84 ab	97 a	81 d
2,4-D choline + glyphosate + glufosinate	2.19 + 0.59	95 a	88 ab	99 a	84 a-d
2,4-D choline + glyphosate + S-metolachlor	2.19 + 1.09	100 a	85 ab	98 a	90 ab
2,4-D choline + glyphosate + acetochlor	2.19 + 1.26	98 a	86 ab	98 a	91 a
2,4-D choline + glufosinate	1.07 + 0.59	98 a	77 ab	98 a	83 bcd
Glyphosate	1.12	100 a	79 ab	88 b	82 cd
Glufosinate	0.59	79 b	49 c	3 c	53 e

^aMeans within a column followed by the same letter are not significantly different according to Fisher's Protected LSD at $P < 0.05$. In 2013, 10 to 15 and 20 to 30 cm kochia control ratings were combined due to no weed height effect ($P > 0.05$). Kochia was not evaluated in 2015

Regardless of weed size, treatments that included 2,4-D choline were the most successful. Among these treatments, 2,4-D choline + glufosinate and 2,4-D choline + glyphosate + glufosinate achieved the greatest levels of weed control. Glyphosate alone applications were inconsistent, especially for larger weeds and glufosinate alone performed poorly across weed species with the exception of 3 to 5 cm Palmer amaranth in 2015 and 10 to 15 cm Russian-thistle in 2015.

4. CONCLUSION

The greatest level of weed control for all three weed species was achieved at the 3 to 5 cm timing; however, weed size was most critical for Palmer amaranth and Russian-thistle compared to kochia. Averaged over all three years, Palmer amaranth control decreased from 93 to 74% following treatments that included 2,4-D choline when applied to plants 3 to 5 and 10 to 30 cm, respectively. For Russian-thistle, control decreased from 98 to 78% following treatments that included 2,4-D choline when applied to plants 3 to 5 and 10 to 30 cm, respectively. For kochia, control decreased from 98 to 84% following treatments that included 2,4-D choline when applied to plant 3 to 5 and 10 to 30 cm, respectively.

ACKNOWLEDGEMENTS

The collaborators of this research are grateful for funding provided by Dow AgroSciences and the Texas State Support Committee, Cotton Incorporated. Work on this project also has been enhanced by the technical support and expertise of the following people: Shay Morris, Seth Taylor, Rand Merchant, and Mike Lovelace.

COMPETING INTERESTS

Authors have declared that no competing interests exist. The products used for this research are commonly used products in our area of research and country. There is no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge.

REFERENCES

1. Stuart BL, Harrison SK, Abernathy JR, Krieg DR, Wendt CW. The response of

- cotton (*Gossypium hirsutum*) water relations to smooth pigweed (*Amaranthus hybridus*) competition. *Weed Sci.* 1984;32: 126-132.
2. Webster TM, Nichols RL. Changes in the prevalence of weed species in the major agronomic crops of the Southern United States: 1994/1995 to 2008/2009. *Weed Sci.* 2012;60:145-157.
3. Beckie JH. Herbicide resistant weeds: Management tactics and practices. *Weed Technol.* 2006;20:793-814.
4. Pan WL, Young FL, Bolton RP. Monitoring Russian thistle (*Salsola iberica*) root growth using a scanner-based, portable mesohizotron. *Weed Technol.* 2001;15: 762-766.
5. Young FL. Effect of Russian thistle (*Salsola iberica*) interference on spring wheat (*Triticum aestivum*). *Weed Sci.* 1988;36:594-598.
6. Beckie HJ, Francis A. The biology of Canadian weeds. 65. *Salsola tragus* L. (Updated). *Can J Plant Sci.* 2009;89:775-789.
7. Schillinger WF. Ecology and control of Russian thistle (*Salsola iberica*) after spring wheat harvest. *Weed Sci.* 2007;55: 381-385.
8. Warwick SI, Sauder CA, Beckie HJ. Acetolactate synthase (ALS) target-site mutations in ALS inhibitor-resistant Russian thistle (*Salsola tragus*). *Weed Sci.* 2010;58:244-251.
9. Baker DV, Withrow JR, Brown CS, Beck KG. Tumbling: Use of diffuse knapweed (*Centaurea diffusa*) to examine an understudied dispersal mechanism. *Invasive Plant Sci Manage.* 2010;3:301-309.
10. Christoffoleti PJ, Westra PB, Moore F. Growth and analysis of sulfonylurea-resistant and -susceptible kochia (*Kochia scoparia*). *Weed Sci.* 1997;45:691-695.
11. Eberlin CV, Fore ZA. Kochia biology. *Weeds Today.* 1984;15:5-6.
12. Forcella F. Spread of Kochia in the northwestern United States. *Weeds Today.* 1985;16:4-6.
13. Friesen LF, Beckie HJ, Warwick SI, Van Acker RC. The biology of Canadian weeds. 138. *Kochia scoparia* (L.) Schrad. *Can J Plant Sci.* 2009;89:141-167.
14. Schwinghamer TC, Van Acker RC. Emergence timing and persistence of kochia (*Kochia scoparia*). *Weed Sci.* 2008;56:37-41.

15. Wicks GA, Martin AR, Haack AE, Mahnken GW. Control of triazine-resistant kochia (*Kochia scoparia*) in sorghum (*Sorghum bicolor*). Weed Technol. 1994;8:748- 753.
16. Heap I. Herbicide resistant Palmer amaranth globally (*Amaranthus palmeri*); 2019. [Accessed 28 February 2019]
17. Heap I. Herbicide resistant Russian-thistle globally (*Salsola tragus*); 2019. [Accessed 28 February 2019]
18. Heap I. Herbicide resistant kochia globally (*Kochia scoparia*); 2019. [Accessed 28 February 2019]
19. Wright TR, Shan G, Walsh TA, Lira JM, Cui C, Song P, Zhuang M, Arnold NL, Lin G, Yau K, Russell SM, Cicchillo RM, Peterson MA, Simpson DM, Zhou N, Ponsamuel J, Zhang Z. Robust crop resistance to broadleaf and grass herbicides provided by arloxyalkanoate dioxygenase transgenes. Proc Natl Acad Sci USA. 2010;107:20240-20245.
20. Richburg JS, Wright JR, Braxton LB, Robinson AE. Inventors: Dow agro sciences, assignee 12 July. Increased tolerance of DHT-enabled plants to auxinic herbicides resulting from MOIETY differences in auxinic molecule structures. US Patent. 2012;13,345,236.
21. Bellinder RR, Arsenovic M, Shah DA, Rauch BJ. Effect of weed growth stage and adjuvant on the efficacy of fomesafen and bentazon. Weed Technol. 2003;51: 1016- 1021.
22. Craigmyle BD, Ellis JM, Bradley KW. Influence of weed height and glufosinate plus 2,4-D combinations on weed control in soybean with resistance to 2,4-D. Weed Technol. 2013;27:271-280.
23. Mellendorf TG, Young JM, Matthews JL, Young BG. Influence of plant height and glyphosate on saflufenacil efficacy on glyphosate-resistant horseweed (*Conyza canadensis*). Weed Technol. 2013;27:463-467.
24. Schuster CL, Shoup DE, Al-Khatib K. Response of common lambsquarters (*Chenopodium album*) to glyphosate as affected by growth stage. Weed Sci. 2007; 55:147-151.
25. Everman WJ, Burke IC, Allen JR, Collin J, Wilcut JW. Weed control and yield with glufosinate-resistant cotton weed management systems. Weed Technol. 2007;21:695-701.
26. Merchant RM, Culpepper AS, Eure PM, Richburg JS, Braxton LB. Controlling glyphosate-resistant Palmer amaranth (*Amaranthus palmeri*) in cotton with resistance to glyphosate, 2,4-D, and glufosinate. Weed Technol. 2014;28:291-297.
27. Richardson RJ, Wilson HP, Armel GR, Hines TE. Trifloxysulfuron plus pyriithiobac mixtures for broadleaf weed control in cotton (*Gossypium hirsutum*). Weed Technol. 2006;20:130-136.
28. Saxton AM. A macro for converting mean separation output to letter groupings in Proc Mixed. Pages 1243-1246 in Proceedings of the 23rd SAS Users Group International. Cary, NC: SAS Institute; 1994.
29. Everitt JD, Keeling JW. Weed control and cotton (*Gossypium hirsutum*) response to preplant applications of dicamba, 2,4-D, and diflufenzopyr plus dicamba. Weed Technol. 2007;21:506-510.
30. Siebert JD, Griffin JL, Jones CA. Red morningglory (*Ipomoea coccinea*) control with 2,4-D and alternative herbicides. Weed Technol. 2004;18:38-44.
31. Sellers BA, Ferrell JA, MacDonald GE, Kline WN. Dogfennel (*Eupatorium capillifolium*) size at application affects herbicide efficacy. Weed Technol. 2009; 23:247- 250.

© 2019 Manuchehri et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/48301>