



Effect of Dates of Sowing on Grain Yield of *Rabi* Rice (*Oryza sativa* L.)

**Boppidi Manasa Reddy^{a++*}, S. G. Mahadevappa^{b#},
K. Indudhar Reddy^{c†} and K. Pavan Chandra Reddy^{d‡}**

^a Department of Agronomy, College of Agriculture, Rajendranagar, PJTSAU, 500030, India.

^b ACRC, ARI, Rajendranagar, Hyderabad, PJTSAU, 500030, India.

^c AICRP on Women in Agriculture, PG&RC, PJTSAU, 500030, India.

^d AICRP on STCR, ARI, Rajendranagar, PJTSAU, 500030, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2023/v13i113212

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107767>

Original Research Article

Received: 29/07/2023

Accepted: 05/10/2023

Published: 12/10/2023

ABSTRACT

Aim: To determine the optimum date of sowing to realize higher grain yield in *rabi* rice.

Study Design: Split plot.

Place and Duration of Study: Krishi Vigyan Kendra, Rudrur, Nizamabad during *rabi*, 2022.

Methodology: The experiment consisted of a total of fifteen treatments which were laid out in split plot design with three replications. Treatments included were five dates of sowing in the main plot M₁-20th November, M₂-1st December, M₃-10th December, M₄-20th December, and M₅-30th December and three treatments of varieties in subplots S₁-JGL 24423 (long slender variety), S₂-Ganga Kaveri (medium slender variety) and S₃-RNR 15048 (short slender variety) randomly placed in subplots of the main plot.

⁺⁺ P.G Scholar;

[#] Senior Scientist;

[†] Scientist;

[‡] Principal Scientist;

*Corresponding author: E-mail: boppidimanasareddy@gmail.com;

Results: Maximum plant height (100.3 cm), number of tillers m^{-2} (432), dry matter accumulation (15869 $kg\ ha^{-1}$), grain yield (7219 $kg\ ha^{-1}$), straw yield (8387 $kg\ ha^{-1}$) and HI (46.1) was noticed in crop Sown on 30th December. Significantly maximum number of tillers m^{-2} , dry matter accumulation (14263 $kg\ ha^{-1}$), grain yield (6127 $kg\ ha^{-1}$), straw yield (8650 $kg\ ha^{-1}$) was observed in S₂-Ganga Kaveri while significantly higher plant height (100.6 cm) and HI (45.0) was found in S₃-RNR 15048. Lowest height was recorded in S₂-Ganga Kaveri (92.6 cm), and above parameters were recorded lowest in crop sown on 20th November, while lowest number of tillers m^{-2} and straw yield (8122 $kg\ ha^{-1}$) was observed in crop sown on 1st December. Significantly less number of tillers m^{-2} (326), dry matter accumulation (13981 $kg\ ha^{-1}$), grain yield (6166 $kg\ ha^{-1}$), straw yield (7526 $kg\ ha^{-1}$) was recorded in S₃-RNR 15048, while lowest HI (43.9) was recorded in S₁-JGL 24423.

Conclusion: Crop sown on 30th December revealed better performance in terms of grain yield under the present study during Rabi conditions. Among varieties, Ganga Kaveri (S₂) performed better in terms of grain yield.

Keywords: Sowing dates; varieties; yield.

1. INTRODUCTION

Rice (*Oryza sativa* L.) is one of the most important food crops in the world, forms the staple diet of billions of people, and provides 70% of direct employment to rural India. In Asia, its cultivation is crucial to maintain food security as more than 90% of rice production is consumed in this continent. Rice contributes 32-59% of the dietary energy and 25-44% of the dietary protein in 39 countries. Globally, rice is grown in an area of 165.3 million hectares with an annual production of 502.98 million metric tons and 4.63 million tons ha^{-1} of productivity [1]. The projected demand for rice by 2050 is estimated to be 197.40 million tons for a population of 1.65 billion almost 80% higher than the current demand [2]. India is the second largest producer of rice after China and it is the staple food for more than two-thirds of the Indian population contributing to 40% of the total foodgrain production. Rice plays a pivotal role in the food and livelihood security of the Asian people as the continent wise data shows, more than 90% of rice production and consumption takes place in Asia and more than two billion people are getting 60-70% of their energy requirement from rice and its derived products [3].

In India, rice crop is grown in an area of 45.8 million hectares occupies prime position in the world with a production of 124.4 million metric tons and productivity of 2717 $kg\ ha^{-1}$ [4]. In Telangana, rice is the major crop extensively cultivated in all the districts in an area of 97.98 lakh acres with the production of 202.18 lakh tons during 2021–22 [5].

The time of sowing in rice is very crucial to ensure adequate vegetative growth which influence the grain yield during a period of optimum temperature with high solar radiation. Selecting the appropriate sowing date for each cultivar guarantees that the vulnerable cold sensitive phase coincides with historically warmer minimum night temperatures, thus ensuring that grain filling takes place under milder temperature conditions. sowing too early or too late diminishes yields in a manner that cannot be offset by any other input or agricultural practice [6]. Most of the world rice is grown in the tropics and the critical determinant for the growth to be temperature. Rice is being a tropical and sub-tropical plant that requires a fairly high temperature ranging from 20° to 40°C. Crop is adversely affected by high temperature in the lower elevation of the tropics and by lower temperature in the temperate regions. The critical temperature differs with variety, duration of critical temperature, diurnal changes and physiological changes of plant. Higher maximum and minimum temperature during ear initiation depress the yield. In general, high temperature accelerates the floral initiation. Rice is most sensitive to high temperatures at heading and next most sensitive at about 9 days before heading. One or two hours of high temperature at anthesis has a decisive effect on the incidence of sterility. High temperatures before or after anthesis have much less effect on sterility. Optimum temperature for ripening is 20-25°C. Too high and too low temperature may cause adverse effect on flowering and prevent pollen shedding leading to high infertility and production of chaffy grains resulting in yield loss. In order to ensure a determined yield level, avoid damage due to too high and low temperature. It is

necessary to optimize time of sowing to attain maximum yield.

There is large diurnal variation in *rabi* season in Telangana region with colder nights and bright sunny days and temperature gradually increases from march onwards with very low relative humidity which is very favourable for higher yields. The crop is very much subjected to wide environmental conditions. Some research on long slender rice grain varieties was carried out in *rabi* season. The research on evaluation of fine grain varieties in *rabi* season is meagre. Keeping the above factors in mind, the present study is proposed during *rabi* 2022.

2. METHODS AND MATERIALS

Field experiment was conducted at Krishi Vigyan Kendra, Rudrur, Nizamabadduring *Rabi* (2022). The experimental site is geographically located in Northern Telangana agro climatic zone of Telangana. The soil of experimental site is silty clay in texture with nearly neutral pH in reaction (8.1), E.C(1.94 dsm^{-1}), low in organic carbon (0.42%), medium in available N(231.1 kg ha^{-1}) & Phosphorous (29.8 kg ha^{-1}) and high in available potassium (350.5 kg ha^{-1}).

The experiment consisted of totally fifteen treatments which was laid out in split plot design with three replications. Treatments included were five dates of sowing in main plot M₁-20th November, M₂-1st December, M₃-10th December, M₄-20th December and M₅-30th December and three treatments of varieties in sub plots S₁-JGL 24423 (long slender variety), S₂-Ganga Kaveri (medium slender variety) and S₃-RNR 15048 (short slender variety) randomly placed in subplots of the main plot. JGL 24423 is a long slender, non-lodging, less shattering and resistant to brown plant hopper with medium duration of 135-140 days with a yield potential of 7500 to 8000 kg ha^{-1} . Ganga Kaveri is a medium slender grain type with medium duration of 135 days duration. It is highly resistant to blast, brown plant hopper with a yield potential of 8750 to 9500 kg ha^{-1} . RNR 15048 is a short slender grain type with medium duration (135 days) and highly photosensitive in nature. It is highly resistant to blast and gives higher yields (6500 to 7500 kg ha^{-1}).

The treatments are sown with a spacing of 15×15 cm. Recommended dose of fertilizers 120-60-40 NPK ha^{-1} , urea, single super phosphate, murate of potash are the fertilisers

used in this experiment. Adequate plant protection measures were taken as per requirement. Randomly five plants were selected and tagged from each plot for recording various growth and yield parameters periodically and at harvest through destructive and non-destructive sampling. The rainfall of 22mm was received during 2 rainy days during the entire period of crop growth. The mean maximum temperature and minimum temperature recorded was 33.4°C and 19.3°C respectively. The mean weekly bright sunshine hours varied from 4.2 to 9.1 hr day^{-1} with an average of 7.3 hr day^{-1} . All the data recorded in the study were conducted statistical analysis of variance technique for split plot design.

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Plant height

Varied sowing dates showed significant influence on plant height at different growth stages of rice.

Rice crop sown on 30th December recorded significantly higher plant height (38.7 cm) at Maximum tillering stage over crop sown on 20th November (34.6 cm) and 1st December (34.2 cm). However, plant height of crop sown on 30th December was comparable with plant height of the crop sown on 20th December (37.3 cm) and 10th December (37.0 cm).

At panicle initiation stage, significantly higher plant height was observed in crop sown on 30th December (75.2 cm) and which was comparable with the plant height of the 20th December (70.2 cm) sown crop. Significantly lower plant height was recorded in crop sown on 1st December (59.1 cm), which was on par with that of plant height recorded with 20th November sowing (59.8 cm). Crop sown on 10th December recorded plant height of 64.7 cm, which was comparable with the plant height recorded in crops sown on 20th November, 1st December and 20th December.

At 50% flowering stage, crop sown on 30th December recorded significantly higher plant height (91.4 cm) and which was on par with 20th December (90.6 cm) and 10th December (87.9 cm) sown crop plant height. Significantly lower plant height was observed in crop sown on 20th November (78.9 cm) over the rest of the treatments.

At physiological maturity stage, significantly higher plant height was observed in crop sown on 30th December (100.3 cm) and which was on par with that of plant height recorded with 20th December (98.9 cm) and 10th December (94.6 cm) sown crop. Significantly lower plant height was recorded in M₁-Sowing on 20th November (92.4 cm) which was on par with crop sown on 1st December (92.6 cm). Significantly higher plant height at all stages of the crop growth were observed with delayed sowing i.e., on 30th December. This might be due to optimum weather conditions favoured in rapid cell division and elongation of plants. The vegetative growth period had a positive effect on the plant growth. Similar findings were reported by Bindu and Bhanumurthy [7]. These results align with the research conducted by Anil and Siddi [8] and Bindu [9], who reported that plant height is influenced by genotypes, various management factors and the environment. The increased plant height observed in delayed sowings in the season could be attributed to greater exposure to solar radiation and heightened photosynthetic activity. These findings are consistent with those reported by Roy [10], who reported that the initial stunted growth of the *borocrop* may be attributed

to an extended period of cold weather during the vegetative growth stage. However, as the air temperature increased, the plant height gradually improved.

Different varieties of rice crop shown significant influence on plant height at different growth stages in different sowing times. Plant height gradually increases as the crop growth stages advance until it reaches the maturity stage. Plant height was increasing gradually from tillering, thereafter increased linearly up to 50% flowering and then it continued to increase till maturity but at a diminishing rate.

At maximum tillering stage, significantly higher plant height (37.2 cm) was observed with RNR 15048 and which was statistically on par with Ganga Kaveri (36.4 cm). Significantly lower plant height (35.5 cm) was noticed with JGL 24423 and which was comparable with Ganga Kaveri. At panicle initiation stage, significantly higher plant height (66.4 cm) was recorded with RNR 15048 which was comparable with that of plant height of Ganga Kaveri (66.2 cm) and JGL 24423 (64.9 cm).

Table 1. Plant height(cm) at different growth stages of rice as influenced by sowing dates and varieties

Treatments	Maximum tillering	Panicle initiation	50% Flowering	Physiological maturity
Mainplot: Sowings dates				
M ₁ - 20 th November	34.6	59.8	78.9	92.4
M ₂ - 1 st December	34.2	59.1	84.2	92.6
M ₃ - 10 th December	37.0	64.7	87.9	94.6
M ₄ - 20 th December	37.3	70.2	90.6	98.9
M ₅ - 30 th December	38.7	75.2	91.4	100.3
SEm(±)	0.67	2.00	1.24	2.60
CD (p=0.05)	2.23	6.60	4.11	6.50
Subplot: Varieties				
S ₁ - JGL 24423	35.5	64.9	84.6	94.1
S ₂ - Ganga Kaveri	36.4	66.2	84.2	92.6
S ₃ - RNR 15048	37.2	66.4	91.1	100.6
SEm(±)	0.41	0.77	1.30	1.67
CD (p=0.05)	1.22	2.40	3.85	4.95
Interaction				
M×S				
SEm(±)	1.17	3.46	2.25	6.20
CD (p=0.05)	NS	NS	NS	NS
S×M				
SEm(±)	1.01	2.45	2.77	4.70
CD (p=0.05)	NS	NS	NS	NS




Fig. 1. Plant height(cm) at different growth stages of rice as influenced by sowings dates and varieties

At 50% flowering stage, significantly higher plant height (91.1 cm) was recorded with RNR 15048 over other varieties. Significantly lower plant height (84.2 cm) was noticed with Ganga Kaveri which was on par with JGL 24423 (84.6 cm). At physiological maturity, significantly higher plant height (100.6 cm) was recorded with RNR 15048 over other varieties and JGL 24423 (94.1 cm) and Ganga Kaveri (92.6 cm) were comparable with each other. Higher plant height was observed with RNR 15048 compared to other varieties during the different crop growth stages. It might be due to the genetic character of the variety and higher photosynthetic efficiency. These results are similar to those obtained by Anil and Siddi [8], Nizamani et al. [11] and Suleiman et al. [12] who observed that plant height differed significantly among the varieties.

3.1.2 Number of tillers m⁻²

Varied sowing dates showed significant influence on number of tillers m⁻² in rice at different growth stages.

At maximum tillering stage, crop sown on 30th December recorded significantly higher number of tillers m⁻² (459) over the advanced sowings and followed by crop sown on 20th December (430) which was on par with the other treatments. Significantly lower number of tillers

m⁻² (403) was observed in crop sown on 1st December which was comparable with that of tillers (411) in crop sown on 20th November and 10th December. At panicle initiation stage, significantly higher number of tillers m⁻² (453) was observed in crop sown on 30th December over other sowing times and followed by crop sown on 20th December (417) which was on par with the rest of the treatments and crop sown on 20th November produced lowest number of tillers m⁻² (396).

At 50% flowering stage, crop sown on 30th December recorded significantly higher number of tillers m⁻² (445) over the rest of the treatments and followed by crop sown on 20th December recorded 398 number of tillers which was on par with other dates of sowings. At physiological maturity stage, crop sown on 30th December recorded significantly higher number of tillers m⁻² (432) over the other treatments and followed by 20th December sown crop (399) which was on par with 10th December (383) and 20th November (375) sowings. Significantly lower number of tillers m⁻² (371) was observed with crop sown on 1st December. This might be due to availability of favourable weather conditions during crop growth cycle. The tillering ability of the crop sown on 20th November was affected severely as minimum temperatures dropped to below 10°C (9.6°C during the standard week 49). The rice crop

exhibits optimal tillering rates within a temperature range of 25-31°C, while it performs poorly at critical low temperatures ranging from 9-16°C. The variation in the number of total and effective tillers per hill can be attributed to differences in cultivar genetics, temperature and sunlight levels among the various cultivars and transplanting dates as reported by Yoshida. [13], Sarwar et al. [14] and Roy et al. [15].

Different varieties of rice shown significant influence on number of tillers m⁻² at different growth stages of rice crop. Number of tillers m⁻² were increasing from transplanting and reached maximum at maximum tillering and thereafter it decreased as the stage progress.

At maximum tillering stage, medium slender variety Ganga Kaveri recorded significantly higher number of tillers m⁻² (538) over other varieties followed by long bold variety JGL 24423 (367), which was on par with short slender variety RNR 24423 (365). At Panicle initiation, 50% flowering and physiological maturity stage, significantly higher number of tillers m⁻² (540, 531 and 512) were observed in Ganga Kaveri over other varieties followed by JGL 24423 (355, 346 and 338), which was on par with RNR 24423 (346, 330 and 326). The number of tillers m⁻² varied among the cultivars [16]. It might be due to differences in genetic makeup of these rice varieties. These results are similar to those obtained by Anil and Siddi [8], Mali and Choudhary [17] and sarwar et al. [14].

3.1.3 Dry matter accumulation

Varied sowing dates showed significant influence on dry matter accumulation in rice at different growth stages.

At maximum tillering stage, crop sown on 30th December recorded significantly higher dry matter accumulation (4877 kg ha⁻¹) over the other treatments and followed by 20th December (4607 kg ha⁻¹) which was on par crop sown on 10th December (4461 kg ha⁻¹) and 1st December (4368 kg ha⁻¹). Significantly lower dry matter accumulation was recorded in crop sown on 20th November (4130 kg ha⁻¹) over the rest of the treatments. At panicle initiation stage, significantly higher dry matter accumulation (9877 kg ha⁻¹) was recorded with crop sown on 20th December and which was on par with 30th December (9874 kg ha⁻¹) and 10th December (9691 kg ha⁻¹) sowings. Significantly lower dry

matter accumulation (9137 kg ha⁻¹) was recorded with crop sown on 1st December and which was comparable with 20th November sown crop dry matter accumulation (9186 kg ha⁻¹).

At 50% flowering stage, crop sown on 30th December recorded significantly higher dry matter accumulation (14474 kg ha⁻¹) and which was on par with crop sown on 20th December (14022 kg ha⁻¹) and 10th December (13948 kg ha⁻¹). Significantly lower dry matter accumulation (13219 kg ha⁻¹) was recorded with crop sown on 20th November and which was comparable with that of crop sown on 1st December (13387 kg ha⁻¹). Similarly, at physiological maturity stage, crop sown on 30th December recorded significantly higher dry matter accumulation (15869 kg ha⁻¹) and which was on par with crop sown on 20th December (15242 kg ha⁻¹) and 10th December (15283 kg ha⁻¹). Significantly lower dry matter accumulation (14448 kg ha⁻¹) was recorded with crop sown on 20th November and which was comparable with that of crop sown on 1st December (14675 kg ha⁻¹).

Dry matter production of early-planted crop was lower and it increased with the advancement of sowing dates due to the favourable weather conditions. The results are in line with the findings of Mannan et al. [18]. Taller plants, higher LAI and increased tiller production might have led to higher dry matter accumulation with crop sown on 30th December. The results are in line with the findings of Bindu and Bhanumurthy [7].

Among the varieties, significantly higher dry matter accumulation was found in Ganga Kaveri from the maximum tillering to physiological maturity stage. At maximum tillering, panicle initiation, 50% flowering and at physiological maturity, significantly higher dry matter accumulation was observed in Ganga Kaveri (4564, 9924, 14445 and 15800 kg ha⁻¹, respectively) and which was on par with JGL 24423 (4549, 9751, 14199 and 15529 kg ha⁻¹, respectively). Significantly lower dry matter accumulation (4353, 8985, 12786 and 13981 kg ha⁻¹, respectively) was observed with RNR 15048. Variations in dry matter accumulation among the different varieties could be attributed to their genetic potential as well as differences in height and growth rates. The results are in line with the findings of Anil and Siddi [8] and Dileep et al. [19].

Table 2. Number of tillers m² at different growth stages of rice as influenced by sowings dates and varieties

Treatments	Maximum tillering	Panicle initiation	50% flowering	Physiological maturity
Mainplot:Sowingsdates				
M₁ - 20 th November	411	396	387	375
M₂ - 1 st December	403	397	384	371
M₃ - 10 th December	411	404	397	383
M₄ - 20 th December	430	417	398	399
M₅ - 30 th December	459	453	445	432
SEm(±)	7.30	7.54	10.39	7.28
CD (p=0.05)	24.17	24.97	34.40	24.10
Subplot:Varieties				
S₁ - JGL 24423	367	355	346	338
S₂ - Ganga Kaveri	538	540	531	512
S₃ - RNR 15048	365	346	330	326
SEm(±)	6.01	6.21	7.77	6.00
CD (p=0.05)	17.85	18.45	23.10	17.84
Interaction				
M×S				
SEm(±)	12.64	13.10	17.99	12.60
CD (p=0.05)	NS	NS	NS	NS
S×M				
SEm(±)	13.18	13.61	17.58	13.16
CD (p=0.05)	NS	NS	NS	NS




Fig. 2. Number of tillers m² at different growth stages of rice as influenced by sowings dates and varieties

Table 3. Dry matter accumulation (kg ha⁻¹) at different growth stages of rice as influenced by sowings dates and varieties

Treatments	Maximum tillering	Panicle initiation	50% Flowering	Physiological maturity
Mainplot:Sowingsdates				
M₁ - 20 th November	4130	9186	13219	14448
M₂ - 1 st December	4368	9137	13387	14675
M₃ - 10 th December	4461	9691	13948	15283
M₄ - 20 th	4607	9877	14022	15242

Treatments	Maximum tillering	Panicle initiation	50% Flowering	Physiological maturity
December				
M ₅ - 30 th December	4877	9874	14474	15869
SEm(±)	67.67	148.42	217.50	238.13
CD (p=0.05)	224.10	491.54	720.31	788.62
Subplot: Varieties				
S ₁ - JGL 24423	4549	9751	14199	15529
S ₂ - Ganga Kaveri	4564	9924	14445	15800
S ₃ - RNR 15048	4353	8985	12786	13981
SEm(±)	50.45	107.15	154.91	169.22
CD (p=0.05)	149.86	318.33	460.22	502.73
Interaction				
M×S				
SEm(±)	117.20	257.08	376.72	412.45
CD (p=0.05)	NS	NS	NS	NS
S×M				
SEm(±)	114.29	245.57	356.80	390.10
CD (p=0.05)	NS	NS	NS	NS




Fig. 3. Dry matter accumulation (kg ha⁻¹) at different growth stages of rice as influenced by sowings dates and varieties

3.2 Yield Parameters

3.2.1 Grain yield

The effect of sowing dates was found significant for grain yield in rice crop. Crop sown on 30th December recorded significantly higher grain yield (7219 kg ha⁻¹) over other dates of sowing followed by 20th December (6758 kg ha⁻¹) which was on par with 10th December (6648 kg ha⁻¹). Significantly lower grain yield (6067 kg ha⁻¹) was recorded with crop sown on 20th November which was on par with 1st December (6289 kg ha⁻¹). The increase in grain yield during delayed sowing dates can be attributed to the

optimization of growth parameters and yield attributing factors like higher number of panicles m⁻², an increased number of filled grains panicle⁻¹, and test weight, all of which positively contribute to the overall grain yield. These results corroborate with the findings of Bindu [9]. The enhanced yield could be attributed to the optimal growth and development parameters, as well as the yield-contributing factors associated with favourable weather conditions. These conditions led to increased growth and development, resulting in a higher accumulation of photosynthates in the grains. These results are in line with the findings of Chendgeet al. [20] and Walia et al. [21]. It was observed that the grain

yield increased by 16.8% with delayed sowing from 20th November to 30th December. The yield potentiality of the crop depends on the type of genotype and the environmental condition especially the air temperature and solar radiation of a particular area [22]. Cultivars sown on December 30th benefited from consistently favourable temperature conditions throughout their growth cycle, experiencing minimal stress during both the seedling and reproductive growth stages. Similar findings on sowing dates were also reported by Singh and Singh [23].

The effect of varieties was found significant for grain yield in rice crop. Among different varieties tested, medium slender variety Ganga Kaveri recorded significantly higher grain yield (6879 kg ha⁻¹) which was on par with the long bold variety JGL 24423 (6744 kg ha⁻¹). Short slender variety RNR 15048 recorded significantly lower grain yield (6166 kg ha⁻¹). The superiority of Ganga Kaveri and JGL 24423 might be due to its better growth, dry matter accumulation and yield attributes *i.e.*, a greater number of filled grains panicle⁻¹ and test weight. These results are in line with the findings of Bindu [9]. Significantly lower grain yield was found in short slender variety RNR 15048 (6166 kg ha⁻¹) which was 10.9% and 8.6% lesser than Ganga Kaveri and JGL 24423, respectively might be due to poor yield attributing characters like lower test weight and higher number of unfilled grains panicle⁻¹

resulting in higher sterility%. The combined impact of growth and yield characteristics ultimately manifested in the form of grain production. Similar findings were reported by Hussain et al. [24] and Anil and Siddi [8]. The superiority of variety Ganga Kaveri might be due to its superior growth

characteristics like its ability to accumulate higher dry matter. These results are in accordance with Bindu [9].

3.2.2 Straw yield

Varied sowing dates showed significant influence on straw yield of rice. Crop sown on 30th December recorded significantly higher straw yield (8387 kg ha⁻¹) and which was on par with 10th December (8365 kg ha⁻¹) and 20th December (8226 kg ha⁻¹). Crop sown on 1st December recorded significantly lower straw yield (8122 kg ha⁻¹) which was on par with crop sown on 20th November (8123 kg ha⁻¹). The later three delayed sowings resulted in an increased straw yield primarily due to the higher dry matter accumulation. This can be attributed to the extended vegetative phase and favourable climatic conditions, which positively influenced dry matter production and subsequently augmented the straw yield. These observations align with the findings of previous research conducted by Assaduzzaman [25] and Bindu [9].

Table 4. Yield of rice as influenced by sowing dates and varieties

Treatments	Grain yield(kgha ⁻¹)	Straw yield(kgha ⁻¹)	Harvest index
Mainplot: Sowingsdates			
M ₁ - 20 th November	6067	8123	42.7
M ₂ - 1 st December	6289	8122	43.6
M ₃ - 10 th December	6648	8365	44.3
M ₄ - 20 th December	6758	8226	45.1
M ₅ - 30 th December	7219	8387	46.1
SEm(±)	104.5	70.4	0.90
CD (p=0.05)	340.7	229.4	NS
Subplot: Varieties			
S ₁ - JGL 24423	6744	8558	43.9
S ₂ - Ganga Kaveri	6879	8650	44.2
S ₃ - RNR 15048	6166	7526	45.0
SEm(±)	81.1	84.2	0.70
CD (p=0.05)	240.0	248.2	NS
Interaction			
M×S			
SEm(±)	198.0	120.0	1.50
CD (p=0.05)	NS	NS	NS
S×M			
SEm(±)	185.5	175.0	1.50
CD (p=0.05)	NS	NS	NS




Fig. 4. Yield of rice as influenced by sowing dates and varieties

Different varieties of rice showed significant influence on straw yield. Among different varieties tested, S₂-Ganga Kaveri recorded significantly higher straw yield (8650 kg ha⁻¹) and was on par with S₁-JGL 24423 (8558 kg ha⁻¹). Significantly lower straw yield was found in S₃-RNR 15048 (7526 kg ha⁻¹). These findings are consistent with those reported by [15]. The cultivation of various rice varieties at different times significantly influences their potential.

3.2.3 Harvest index

The effect of sowing dates was found nonsignificant for harvest index in rice crop, while numerically higher harvest index (46.1%) was recorded in crop sown on 30th December and lowest harvest index (42.7%) was recorded in crop sown on 20th November. The significant increase in the harvest index can be primarily attributed to the higher grain yield achieved through improved performance in various yield related traits. These findings are consistent with those reported by Anil and Siddi [8]. The effect of varieties was found nonsignificant for harvest index in rice crop, while numerically higher harvest index was recorded in the cultivar RNR 15048 (45.0%) and lowest harvest index was recorded in the variety JGL 24423 (43.9%) and Ganga Kaveri (44.2%). Similar findings were observed by Salahuddin et al. [26].

4. CONCLUSION

Based on the research work, it can be concluded that among different sowing dates, crop sown on 30th December (M₅) performed better in terms of dry matter accumulation and grain yield and among the varieties, Ganga Kaveri (S₂) performed better in dry matter accumulation resulting in higher grain yield under *rabi* conditions.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. FAOSTAT; 2022. Available: <http://faostat.fao.org/site/567/DesktopDefault.aspx>
2. Joshi S. The future of rice: Shrinking production area, yield and productivity; 2021.
3. Outlook; 2022. Available: <https://www.fao.org/statistics>.
4. Indiastatistics 2020-21. Available: www.indiastat.com, 2020-21.
5. Available: <https://www.telangana.gov.in/PDFDocuments/Telangana-Socio-Economic-Outlook-2023.pdf>

6. Kapoor A, Singh I, Serawat M, Dhaiya S, Meena D, Meena BR. Effects of different dates of transplanting on yield attributes, yields and quality of non-scented varieties of rice (*Oryza sativa* L.). International Journal of Chemical Studies. 2017; 5(6):831-833.
7. Bindu GSM, Bhanumurthy VB. Study of growth parameters in rabi rice as influenced by sowing dates and varieties. Progressive Research. 2007;2(1/2):39-42.
8. Anil D, Siddi S. Performance of pre released rice (*Oryza sativa* L.) genotypes under different sowing windows in rabi season. International Journal of Environment and Climate Change. 2020;10(10):43-51.
9. Bindu GM. Effect of cold on the performance of rice varieties sown on different dates during rabi. Ph.D. Thesis. Acharya N. G. Ranga Agricultural University, Andhra Pradesh, India; 2004.
10. Roy BC. nitrogen fertilizer management and its effect on growth and yield of rice varieties in Bangladesh. Ph.D. Thesis. University of Hohenheim. Germany. 1999;1-17.
11. Nizamani GS, Imtiaz AK, Abdula K, Siddiqui MA, Nizamani MR, Khaskheli MI. Influence of different row spacing on agronomic traits in different wheat varieties. International Journal of Development Research. 2014;4(11):2207-2211.
12. Suleiman AA, Nganya JF, Ashraf MA. Effect of cultivar and sowing date on growth and yield of wheat (*Triticum aestivum* L.) in Sudan. Journal of Forest Products and Industries. 2014;3(4): 198-203.
13. Yoshida S. Fundamentals of rice crop science. In: Climate and rice. Los Banos (Philippines): International Rice Research Institute. 1981;87-88.
14. Sarwar AKMG, Sohail SA, Islam M, Ashrafuzzaman M. Pollen characteristics and yield performances of rice as influence by air temperature and relative humidity. Bangladesh Journal of Botany. 2017;46(3):947-954.
15. Roy TK, Paul SK, Sarkar MAR. Influence of date of transplanting on the growth and yield performance of high yielding varieties of boro rice: Yield of boro rice as affected by transplanting date. Journal of the Bangladesh Agricultural University. 2019; 17(3):301-308.
16. Gomosta AR, Quayyum HA, Mahub AA. Tillering duration and yielding ability of rice varieties in the winter rice season of Bangladesh. In Rice research for food security and poverty alleviation. Proceedings of the International Rice Research Conference. 2001;87-94.
17. Mali H, Choudhary J. Performance of bread wheat (*Triticum aestivum* L.) varieties under different row spacing. Journal of Cereal Research. 013;4(2): 55-57.
18. Mannan MA, Bhuiya MSU, Akhand MIM, Rana MM. Influence of date of planting on the growth and yield of locally popular traditional aromatic rice varieties in boro season. Journal of Science foundation. 2012;10(1):20-28.
19. Dileep K, Pasupalak S, Baliarsingh A. Effect of establishment methods and sowing time on growth and yield of rice varieties (*Oryza sativa* L.). The Pharma Innovation Journal. 2018;7(4): 904-907.
20. Chendge PD, Chavan SA, Patil A, Kumar S. Effect of sowing times on yield and economics of different rice genotypes under climatic condition of Konkan. Journal of Pharmacognosy and Phytochemistry. 2017;6(5):2462-2466.
21. Walia US, Walia SS, Sidhu AS, Nayyar S. Productivity of direct seeded rice in relation to different dates of sowing and varieties in Central Punjab. Journal of Crop and Weed. 2014;10(1):126-129.
22. Singh KM, Pal SK, Verma UN, Thakur R, Singh MK. Effect of time and methods of planting on performance of rice (*Oryza sativa* L.) cultivars under medium land of Bihar plateau. Indian Journal of Agronomy. 1997;42(3):443-445.
23. Singh AK, Singh L. Role of thermal time in rice phenology. Environment and Ecology. 2007;25(1):46.
24. Hussain A, Bhat MA, Ganai MA, Hussain T. Influence of planting dates and spacing schedules on performance of basmati rice Pusa Sugandh-3 under Kashmir valley conditions. Environment and Ecology. 2009;27(1A):396-398.
25. Assaduzzaman M. Effect of spacing and date of transplanting on the performance of boro rice cvb. BRRI dhan. M.Sc. (Agri.) Thesis. Bangladesh Agricultural University, Mymensingh, India; 2006.

26. Salahuddin KM, Chowdhury SH, Munira S, Islam MM, Parvin S. Response of nitrogen and plant spacing of transplanted aman rice. *Bangladesh Journal of Agricultural Research*. 2009; 34(2):279-285.

© 2023 Reddy et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107767>