



Growth Performance and Nutrients Digestibility of Weaner Rabbits FED Diets Containing Par-Boiled and Non-Parboiled Rice Milling By-Products

T. T. Anongo ^{a*}, D. T. Shaahu ^b, T. Ahemen ^c, C. D. Tuleun ^d,
and S. N. Carew ^b

^a Oracle Feed Mill, KM 3, Industrial Layout, Makurdi-Naka Road, Makurdi. Nigeria.

^b Department of Animal Production, Joseph Saawuan Tarkaa University, Makurdi, Nigeria.

^c Department of Animal Breeding and Physiology, Joseph Saawuan Tarkaa University, Makurdi, Nigeria.

^d Department of Animal Nutrition, Joseph Saawuan Tarkaa University, Makurdi, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JEAI/2023/v45i112236

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/107909>

Original Research Article

Received: 12/08/2023

Accepted: 17/10/2023

Published: 03/11/2023

ABSTRACT

A feeding trial was conducted for 84-days to investigate the feeding value of rice milling by-products on growth performance, digestibility and carcass characteristics of growing rabbits in Makurdi. Forty two (42) weaner rabbits weighing between 412 – 420 g were randomly allocated to 7 treatment diets with each treatment replicated 6 times in a completely randomize design (CRD). Each rabbit constituted a replicate and each treatment was made up of three males and three females. The treatments were, D₁, control, D₂ contained parboiled rice offal (PRO), D₃ contained parboiled rice

*Corresponding author: E-mail: phada01@gmail.com;

bran (PRB), D₄ contained parboiled grain rejects (PGR), D₅ contained non-parboiled rice polishing (NPRP), D₆ contained non-parboiled rice bran (NPRB) and D₇ contained non-parboiled grain rejects (NPGR). Result of growth performance revealed that Significant ($P < 0.05$) dietary influence were observed on average daily weight gain and feed conversion ratios. In both parameters (ADWG and FCR) performances were significantly ($P < 0.05$) poor (10.97g and 6.69 respectively) in rabbits fed PRO. Result of digestibility by the rabbits fed diets containing RMBPs were similar ($P > 0.05$) to the rabbits fed the control diet and this similarity was observed across the treatment groups. No significant ($P > 0.05$) dietary influence was observed on carcass parameters measured. It was concluded that, the combination of rice milling by-products (RMBPs) with other feed materials provided cheaper diets that are nutritionally adequate for enhance growth, good carcass development and yield of rabbits at levels equivalent to the reference diet. It is recommended therefore that, feed manufacturers and Rabbit farmers can incorporate up to 25% of rice offal, rice bran and rice grain rejects in the diets of rabbits.

Keywords: Rabbits; par-boiled; non-par-boiled; rice milling by-products (RMBPs); growth; digestibility.

1. INTRODUCTION

World population continues to grow in geometric progression and, thus, places ever-increasing pressure on arable land, water and the other resources required to provide the necessities of life such as healthy food, potable water and clean air [1]. "Consequently, there is insufficient supply of feedstuffs at economic prices. This has continued to limit the production and thus, availability of animal protein in the diets of humans in the developing countries of the world. All stakeholders in the animal production industry are interested in issues of availability and quality of feed. There is, therefore, great interest in the potential of any material to serve as feed ingredients" [2]. "Thus, the use of agricultural by-products in animal feed production is one major strategy that holds great potential for bringing down feed cost" [3]. "Since by-products are generally cheaper than the originating commodity, the use of agricultural by-products will lower the price of animal protein, by reducing feed cost" [4]. This will in turn raise animal protein consumption levels and ultimately improve the health and productivity of the general populace and children in particular. Rice milling by-products (such as rice offal, rice grain, rejects and rice bran) are some of the cheapest and readily available agro-industrial by products that could be used to feed both ruminants and monogastric animals such as rabbits. With the current intervention on industrial rice production in the country, the processing of paddy rice into edible products in Nigeria is increasing rapidly. Recently, rice mills have grown in number and sophistication, in response to increased production of paddy rice, resulting in higher volumes and varieties of by-products. In this study, an attempt was made to nutritionally

determine the feeding value of rice milling by-products (RMBPs) in the diet of growing rabbits.

2. MATERIALS AND METHODS

2.1 Location of the Study

The study was conducted at the Rabbit Unit of Oracle Farms, km 5, Makurdi-Naka Road, Makurdi, Benue State. Makurdi lies within the Southern Guinea Savannah Agro ecological zone of Nigeria.

2.2 Test Materials

The rice milling by-products (Rice offal) were obtained from small-scale rice mills in various locations in Makurdi. Parboiled Rice bran and parboiled grain rejects were purchased from a commercial rice mill; MIKAP rice processing plant, makers of MIVA rice located along kilometer 5 Gboko Road, Makurdi while non-parboiled rice bran, non-parboiled grain rejects and non-parboiled rice polishing was purchased from Oracle Rice Mill located at KM 3, Industrial layout, Makurdi-Naka road, Makurdi. Other ingredients; maize, brewers dried grain (BDG), Soybeans Meal (SBM), palm kernel cake (PKC) were purchased from the market within Makurdi metropolis. Bone charcoal, common salt, Synthetic Methionine, Synthetic Lysine, Vitamin/Mineral premix and drugs required for medications were purchased from a Veterinary and Livestock feed shop within Makurdi.

2.3 Experimental Animals, Design and Management

Forty-two (42) cross breed (Chinchilla x Newzealand white) weaner rabbits aged between

five to seven weeks ranging in live weights between 412.00 g to 420.00 g obtained from rabbit farms within Makurdi were used for the study. The physical appearances of the animals, the disease history and other management practices of the various farms were taken into consideration in choosing the animals to be purchased. This was to ensure that the rabbits were healthy and suitable for experimental purposes. The rabbits were housed in individual cages made of iron bars/wire netting measuring 60 cm x 60 cm x 90 cm raised 60 cm above the floor. The rabbits were allowed seven days to get acclimatised to the new environment and diets after which six rabbits were randomly allocated to each treatment, balancing for sex and minimising live weight differential. Each rabbit served as a replicate. The completely randomised design (CRD) model was adopted. Standard rabbit husbandry practices including feeding standards, hygiene, medications and external/ internal parasite control measures were strictly observed throughout the experimental period. Feed and water were provided *ad-libitum*.

2.4 Experimental Diets

Seven experimental diets coded D₁, D₂, D₃, D₄, D₅, D₆ and D₇ were formulated with D₁ having no rice milling by-product and served as the control, D₂, D₃, D₄, D₅, D₆ and D₇ containing parboiled rice offal (PRO), Parboiled rice bran (PRB), Parboiled grain reject (PGR), none parboiled rice bran (NPRB), none parboiled rice polishing (NPRP) and none parboiled grain reject (NPGR) at 25 % inclusion respectively. The ingredients and calculated nutrient and energy composition of the diets is presented in Table 1.

2.5 Parameters Measured

a. Weight gain: The rabbits were weighed at the beginning of the trial and weekly thereafter. The weight at the beginning of the week was termed initial weight (IW), and that at the end of the week was called present weight (PW), while the weight at the end of the final week was termed final weight (FW). Weight gain (WG) was calculated by difference: $WG = PW - IW$. Average daily weight gain (ADWG) over the entire study period was calculated as: $ADWG = [(FW - IW) \div 84]$ (84 days was the duration of the study).

b. Feed intake: A known quantity of feed termed feed supplied (FS) was set aside for each rabbit at the beginning of the week and the rabbit was

offered feed from it during the week. The left-over feed (LOF) at the end of the week was weighed, and feed consumed/intake (FI) calculated by difference. $FI = [FS - LOF]$. Daily feed intake was computed by dividing feed intake by number of days (84days). $ADFI = [(FS - LOF) \div 84]$ (84 days was the duration of the study).

c. Feed conversion ratio: The feed to gain ratio was computed by dividing feed intake (FI) by the weight gain, (WG) of the animals; as given in the formula:

$$FCR = FI \div WG.$$

Where; FCR= Feed Conversion Ratio; FI= Feed Intake; and WG= Weight Gain.

2.5.1 Nutrient digestibility

At the end of the 12th week of the study, a digestibility trial was carried out. From each treatment, four rabbits (two males and two females), with mean live weight closely approximated to the treatment mean, were selected for the trial. The rabbits were starved for eighteen (18) hours to clear the gut of previous feed residues. During the following five days, seventy-five per cent (75 %) of their daily feed intake was offered to the rabbits and the corresponding feces voided were collected by placing plastic netting under the cages. The weights of the fresh feces collected for each rabbit were recorded, oven dried to a constant weight. At the end of the trial the fecal sample for each rabbit was pooled milled and sampled for proximate analysis as outlined by [5]. Determination of apparent nutrient digestibility was calculated using the formula below:

$$\text{Apparent digestibility} = [(Nutrient\ intake - Nutrient\ voided) \div Nutrient\ intake] \times 100$$

2.5.2 Carcass and visceral organ characteristics

At the end of the feeding period, three rabbits (blocking for sex; 2 males and 1 female) whose live weight most closely approximated their treatment mean were randomly selected from each treatment, starved overnight and then slaughtered. The rabbits were weighed to obtain their live weight, and then stunned with a blow at the back of the head. Their jugular veins were severed with a sharp knife and allowed to bleed thoroughly under gravity. The carcasses were eviscerated, weighed, singed, and, then weighed

Table 1. Ingredients and Calculated Nutrient Composition of Experimental Diets

Ingredients (%)	Diet 1	Diet 2	Diet 3	Diet 4	Diet 5	Diet 6	Diet 7
	Control	25.00% PRO	25.00% PRB	25.00% PGR	25.00% NPRP	25.00% NPRB	25.00% NPGR
Maize	26.55	14.55	13.55	12.55	13.55	13.55	13.55
Soybean meal	12.13	11.13	11.13	13.13	10.10	11.13	12.13
palm kernel cake	20.43	16.43	16.43	15.43	16.43	16.43	15.43
Brewers dried gain	22.19	18.19	17.19	16.19	16.19	17.19	16.19
PRO	-	25.00	-	-	-	-	-
PRB	-	-	25.00	-	-	-	-
PGR	-	-	-	25.00	-	-	-
NPRP	-	-	-	-	25.00	-	-
NPRB	-	-	-	-	-	25.00	-
NPGR	-	-	-	-	-	-	25.00
Soybean straw	13.55	8.55	12.55	12.55	13.58	12.55	12.55
Palm oil	1.00	2.00	2.00	1.00	1.00	2.00	1.00
Bone charcoal	3.00	3.00	3.00	3.00	3.00	3.00	3.00
Methionine	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Table salt	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Vit. /Min. premix®	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Lysine	0.25	0.25	0.25	0.25	0.25	0.25	0.25
TOTAL	100.00	100.00	100.00	100.00	100.00	100.00	100.00
CALCULATED	ANALYSIS						
Crude Protein (%)	17.06	16.95	17.95	16.06	16.58	17.95	16.06
Energy (kcal/kg)	2637.05	2405.38	2500.34	2726.28	2686.18	2500.34	2726.28
Crude Fibre (%)	12.82	17.22	13.47	11.34	11.39	11.34	11.39
Ether extract (%)	4.63	4.87	7.11	4.98	5.81	7.11	5.81

®Vit-Min Premix (Nutrpoult) Vitamin A10,000,000IU, Vitamin D₃ 2,000,000IU, Vitamin E 40,000mg, Vitamin K₃ 2,000mg, Vitamin B₁ 500mg, Vitamin B₂ 5,000mg, Vitamin B₁₂ 20mg, Vitamin B₆ 40,000mg, Niacin 40, 000mg, Calpan 10,000mg, Folic acid 1, 000mg, Biotin 100mg, Antioxidant 100, 000mg, Cholin chloride 300,000mg, Cobalt 300mg, Selenium 200mg, Iodine 800mg, Iron 40,000mg, Manganese 80,000mg, Copper 80,00mg, Zinc 60,000mg. D₁-D₅=Treatments 1,2,3,4 and 5. RO= Rice Offal; RB=Rice Bran, GR=Grain Rejects, NPRB-none parboiled rice bran, NPRP-none parboiled rice polishings, NPGR- none parboiled grain rejects.

again to obtain the dressed weight using an electronic weighing balance. The dressed carcasses were reduced into the traditional cuts; head, neck, forelimbs, rack/ribs, back/loin and hind limbs and the weights of the various cuts were recorded except the tail. These carcass cuts were expressed as a percentage of the dressed weight as given by the formula;

$$\% \text{ Carcass cut} = [\text{Carcass cut} \div \text{Dressed weight}] \times 100$$

The visceral organs (the heart, lungs, kidneys, liver, intestines, stomach, oesophagus, caecum and the spleen) were each removed, trimmed free of fats and adhering connective tissues, weighed and expressed as percentage of live weight.

2.6 Statistical Analysis

The data collected was analyzed by one-way analysis of variance (ANOVA) using [6] and where significant differences occurred, treatment means were separated using the Duncan Multiple Range Test of the same statistical software.

3. RESULTS AND DISCUSSION

3.1 Growth Performance of Rabbits Fed Diets Containing RMBPs, 0-12 Weeks of Feeding

Results of growth performance of rabbits fed diets containing RMBPs revealed a significant ($P < 0.05$) dietary influence on average daily weight gain and feed conversion ratios. In both parameters (ADWG and FCR), performances were significantly ($P < 0.05$) poor in rabbits fed diet D₂ containing parboiled rice offal (PRO).

Rabbits fed diet D₂ recorded significantly ($p < 0.05$) lower average daily weight gain. Similar to the report of this study [7] also reported a significant difference in the daily weight gain of rabbits fed shea butter nut cake. Values obtained in this study ranged between 10.97 -16.64 g/rabbit/day. These values were however higher than 5.81 -12.13 g/rabbit/day reported by [7] but agreed with the values reported by [8] for rabbits fed diets containing rice milling by-products. The lower weight gain observed in rabbits on diet D₂ could be attributed to the low energy content of the diet and thus, poor conversion into meat. Energy is regarded as the main dietary requirement of all animals for both maintenance and other productive functions. The poor ADWG of rabbits on the PRO diet could also have

resulted from the higher CF and lower digestible carbohydrate content of the diet. Thus, this affirms an earlier report by [9] who noted that, the high digestibility of a corn-based diet was because of the high availability of digestible carbohydrates content. The higher the fiber level in the feed of the rabbit, the more quickly it passes through the digestive system and the lower the digestibility [10]. This implies poor efficiency of feed utilization by the animals.

It was expected that the high fibre and low metabolisable energy contents of diet (D₂) to have an influence on feed intake. This could have supported the findings of different scholars. While [11] reported that high fiber diets tend to increase feed intake in rabbits, [12] observed that, rabbits have the ability to adjust their feed intake voluntarily to meet their energy requirements [13] also reported that, various by-products with high fiber can affect voluntary feed intake depending on the nature of fibre. The present study however did not observe any significant difference in intake of feeds across treatments including rabbits on D₂. This report suggests that, both the rabbits on diets containing RMBPs and the control were similar in palatability /acceptability, rabbits were of similar age and the duration of the experiment was long enough to warrant adaptation to the feeds/diets provided. This finding was therefore consistent to the report of Alawa and [14] who had earlier reported that the ability of the rabbit to utilize highly fibrous diets may depend on the source of the fibre, age of the rabbits and length of adaptation of their digestive system to the fibre source. The values revealed in this study were similar to 67.94 – 79.42 g/rabbit/day reported by [8] who also reported similarities across treatment in ADFI of rabbits fed RMBPs. The similar feed intake observed across the treatment can also be explained on the basis of fibres ability to induce gut fill. Supporting the findings of [15] who reported that, “fibre is an intrinsic component of animal feeds and depending on the chemical structure and physiochemical properties, fibre may affect both gut physiology and feed intake”. In some species however, viscous soluble fibres are linked to an increased satiety caused by gastric distention and slower passage rate, and thereby a reduced feed intake. Significant ($P < 0.05$) dietary influence was observed on feed conversion ratio. This report contradicts earlier reports of [16], 17 and [8]. Values obtained in the study range between 4.28-6.69 and significantly ($P < 0.05$) lower in

rabbits fed diet D₂. These values were better than 4.72-9.14 reported by [7] but, were in line with the reports of [18] when fresh and succulent stems of *Ficus thonningi* were fed to growing rabbits. The high fibre and low energy contents of the diet D₂ could be implicated in the poor FCR observed in this study as reflected in the daily weight gain of the rabbits. In addition to the energetic inefficiency characterising the utilization of fibre and low metabolisable energy in D₂, the low weight gain of the rabbits on D₂ could be responsible for the poor value recorded for feed conversion ratio observed in the study. Aside the effect of feed characteristics/quality on voluntary feed intake and utilization, the animal's environment has also been implicated on voluntary feed intake and utilization. This observation was in line with that of [19] who observed that, one of the factors affecting feed conversion is the temperature of an animal's environment. Animals perform optimally when there are minimal variations in environmental temperature over a 24-hour period of time [19]. According to them, on one hand, during cool weather conditions, animals will eat more feed but, much of the energy they obtain from the consumption of the feed is used to maintain body temperature and thus not converted into meat (weight) nor other productive functions [19]. On the other hand, animals consume less feed and convert less efficiently at high environmental temperatures [19]. This biological cooling mechanisms that animals use during hot conditions require energy just as the warming mechanisms do during cool weather conditions. It therefore suggests that, the variations in environmental temperature during the days and nights within the period of experimentation can be implicated in the differences observed in the present study with previous findings.

Overtly, the positive growth characteristics of rabbits on the diets containing both the parboiled and non-parboiled RBMPs compared with the rabbits on the control, observed in this study gives a good indication that the experimental diets (save D₂) sufficiently supplied the nutrients that supported the growth and good performance of the rabbits.

3.2 Digestibility Coefficient of Rabbits Fed Diets Containing RBMPs, 0-12 Weeks of Feeding

It was observed in this study that there was a considerable reduction in the digestibility coefficient of the rabbits in the 12th week of the

study. At week 12, the result of digestibility by the rabbits fed diets containing RBMPs were similar ($P>0.05$) to the rabbits fed the control diet and this similarity was observed across the treatment groups (Table 3).

The values of DM digestibility reported in this study were slightly lower than 74.14 to 83.35 % reported by [20] but higher than 36.66 to 66.02 %, reported by [21]. Digestibility can be influenced by many factors, notably, age and quantity of feed consumed by the animals at the time of metabolic trial. There appears also to be age differences in the ability of the rabbits to utilize the feed consumed. This observation validates an earlier finding of [19] that, animals of higher live weight due to age differences required a higher amount of feed, even to attain a constant weight gain. Obviously, young and fast-growing animals have a far more favorable FCR in the early fattening stage than when near slaughter weight. Moreover, the maintenance requirements become high and are also responsible for the quick increase in FCR [22]; which is a total reflection of the amount of nutrients made available to the rabbits' body for maintenance and production (growth) functions.

Conversely, with the other nutrients, there was a remarkable improvement in the numerical value for the digestibility of crude fiber by the experimental animals. Implying that, the digestibility of crude fiber could be dependent on the age of the rabbit. The development of the caecum and of course its microbial population responsible for fermentation of plant cell wall components can therefore be said to correspond to the age of the rabbit. This aligns to an earlier report of [23] in an experiment to determine the functional, anatomical and histological development of caecum in rabbits. They observed that, the complete physiological developments of the rabbit caecum were evidenced at 12 weeks of age. Implying therefore that, the caecum of the rabbit is fully developed and will be more efficient in the utilization of plant cell wall as the rabbit's age progresses [23] further observed that, increasing feed intake will increase hindgut fermentation and activate microbial metabolism which may stimulate the development of caeca and colonic weights. As feed intake is also reported to correspond with age and body mass, the above observation can also be implicated in this study. Aside the above observation, the longer period of the rabbits on these diets allowed them opportunity to adopt fully to the diets and

Table 2. Growth Performance of Rabbits Fed Diets Containing RMBPs, 0-12 weeks of Feeding

Parameters	D1 (Cont.)	D ₂ PRO	D ₃ PRB	D ₄ PGR	D ₅ NPRP	D ₆ NPRB	D ₇ NPGR	SEM	P
IWt, (g)	413.00	420.00	412.00	418.67	419.00	414.50	414.67	24.84	1.00
FWt (g)	1577.33	1341.67	1773.67	1816.67	1674.17	1634.67	1597.83	45.62	0.12
ADFI (g)	70.38	69.23	77.47	73.10	70.33	76.91	58.76	1.97	0.19
ADWG (g)	13.86 ^a	10.97 ^b	16.21 ^a	16.64 ^a	14.79 ^a	14.52 ^a	14.09 ^a	0.44	0.01
FCR	5.10 ^{ab}	6.69 ^b	4.79 ^a	4.43 ^a	4.77 ^a	5.79 ^{ab}	4.28 ^a	0.22	0.04

ab=Means on the same row with different superscripts are significantly ($P<0.05$) different, NPRB-non parboiled rice bran, NPGR-non parboiled grain rejects, NPRP-non parboiled rice polishing, PRB-parboiled rice bran, PGR-parboiled grain rejects, PRO-parboiled rice offal. IW-initial weight, FW-final weight, ADWG-average daily weight gain, ADFI-average daily feed intake, FCR- feed conversion ratio

Table 3. Nutrients Digestibility by Grower Rabbits Fed Diets Containing RMBPs

Parameters (%)	D1 (Cont.)	D ₂ PRO	D ₃ PRB	D ₄ PGR	D ₅ NPRP	D ₆ NPRB	D ₇ NPGR	SEM	P
DM	65.36	65.80	67.62	79.37	62.91	75.24	62.33	1.99	0.16
NFE	57.17	45.73	53.16	74.25	47.74	53.43	45.92	3.04	0.15
CP	84.38	78.64	84.29	87.13	77.66	87.41	78.28	1.2	0.07
EE	80.59	88.24	91.42	86.27	81.11	91.06	89.27	1.58	0.36
CF	77.41	80.21	73.83	86.53	78.95	83.24	74.80	1.39	0.15

abcd=Means on the same row with different superscripts are significantly ($P<0.05$) different, NPRB-non parboiled rice bran, NPGR-non parboiled grain rejects, NPRP-non parboiled rice polishing, PRB-parboiled rice bran, PGR-parboiled grain rejects, PRO-parboiled rice offal. DM- dry matter, NFE-nitrogen-free extract, EE- ether extract, CF-crude fibre.

Table 4. Carcass Characteristics of Rabbits Fed Diets Containing RMBPs

Parameters (%)	D1 (Cont.)	D ₂ PRO	D ₃ PRB	D ₄ PGR	D ₅ NPRP	D ₆ NPRB	D ₇ NPGR	SEM	P
FLW(g)	1521.00	1389.00	1717.00	1688.00	1655.67	1571.67	1456.33	44.63	0.35
DW(g)	920.00	842.00	1088.33	1036.67	1024.00	1011.33	953.40	32.39	0.51
DP	61.21	60.51	63.31	61.34	61.81	64.27	65.55	0.55	0.15
Head	12.70	12.53	10.94	11.72	11.83	11.16	11.92	0.23	0.38
Neck	3.78	3.40	3.41	3.73	3.81	4.10	3.59	0.10	0.63
Ribs/Rack	15.03	14.56	14.67	14.38	13.70	14.92	16.19	0.37	0.76
Back/Loin	14.58	13.97	15.89	15.81	14.63	15.54	13.43	0.33	0.35
Forelimbs	14.32	13.81	13.01	14.62	13.80	14.00	13.42	0.19	0.43
H. limbs	30.28	30.80	30.43	29.97	29.79	29.59	29.42	0.17	0.39

FLW-fasted live weight, *DW*-dressed weight, *DP*-dressing percentage, NPRB-non parboiled rice bran, NPGR-non parboiled grain rejects, NPRP-non parboiled rice polishing, PRB-parboiled rice bran, PGR-parboiled grain rejects, PRO-parboiled rice offal, SI-small intestine, LI-large intestine, H.limbs-hind limbs. carcass parameters were expressed as a percentage of dressed weight.

perhaps the conditions of the experiments within the study duration therefore leading to no difference in fibre digestion as well as the other nutrients.

3.3 Carcass Characteristics of Rabbits fed Diets Containing RMBPs

Result of carcass characteristics is presented in Table 4. There was no significant ($P>0.05$) dietary influence on carcass parameters measured. Rabbits fed diets containing RMBPs showed values for dressed weight (953.40 g - 1088.33 g) and dressing percentage (61.31 % - 65.55 %) save for rabbits on diet D₂ (842.00 g and 60.51 % respectively).

The influence of feeding diets containing rice milling by-products to growing rabbits on carcasses has been established. It is shown in this study that the use of RMBPs in the diets of rabbits lead to similar ($P>0.05$) responses in final live weight gain across treatment groups. This result is consistent with an earlier report of [24] who also reported similarities in most of the parameters measured with the exception is shown of the forelimbs. This observation indicates that rabbits fed diets containing RMBPs had a fast rate of metabolism and thus growth similar to those fed on control hence, similarities in fasted live weight, dressed weight, dressing percentage and all parameters measured. This implies that, the use of rice milling by-products in rabbit nutrition did not negatively result to the modification of body parts that would affect consumer demand and preferences. Values for dressed weight and dressing percentage were in the range of 920.00 to 1088.33 % and 60.51 to 65.55 % lower than 1008.67 to 1121.67 g and 64.85 and 68.74 % reported by [24] respectively. As reported by [25], the development rate of different body tissues and meat quality features can be modified through an adequate feeding regime based on synchronizing different protein levels in connection to associated dietary energy levels. The body parts were however not modified in this study. The diets therefore, proved their adequacy in meeting the needs of the rabbits for optimal development of body tissues. Also, supporting this observation, [26] explained that, all factors which can influence growth potentials, by changing the relative growth of tissue and organs, lead to modifications of the carcass. The loin and legs are the most economically important portions of the carcass and also provide the greatest portion of edible meat in rabbits [27]. The inclusion of

RMBPs in the diets of the rabbits did not show any negative influence on the relative weight of these economic parts (loin and legs). Age and weight at slaughter and experimental procedures could partly explain differences observed in the present study with previous findings.

4. CONCLUSION

- i. The result of this study revealed that, the inclusion of 25% PRB, PGR, NPRB, NPRP and NPGR in the diets of growing rabbits improved the growth performance of the rabbits at levels equivalent to conventional ingredients/ reference diet.
- ii. The digestibility by rabbits the proximate fractions of RMBPs for growth and maintenance was not adversely affected at 25% inclusion.
- iii. The result of this study also revealed a reduction in the cost of producing a kg live weight of the rabbits and total production cost. Rabbits can therefore be reared at a cheaper cost, better rate of return and high profit made with the inclusion of RMBPs at 25%.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kendall H W and Pimentel D. Constraints on the Expansion of the Global Food Supply. *Ambio*. 1994;23(3): 198-205.
2. Khieu B, Brain O and Jan E L. Methods and techniques for the determination of amino acid digestibility: A Review. *Livestock Research for Agricultural Development*. 2002; 14 (6) 2002
3. Orayaga K T, Ayoade J A and Dose J J. Effects of three sources of Groundnut (*Arachis hypogea*) Meals on Carcass yield, internal organs and Meat quality of Rabbits. *Journal of Animal Advances*.2016; 6 (9): 991-998
4. Yashim SM, Gadzana IU and Anene DO. Nutrient utilization and Carcass Characteristics of Weaner Rabbit as influenced by differently processed Rattle Box (*Crotalaria refusa*) Leaf Meal. *Journal of Animal Production Research*. 2017; 29 (1): 171-179.
5. AOAC. Official methods of analysis. Association of Official Analytical Chemists.

- Official Methods of Analysis 17th edition Washington, D.C. 2002; 20-27.
6. SPSS 21. Statistical Package for Scientific Solutions, Software, 2016; 21 version
 7. Shaahu D. T, Dzungwe E N, Ahemen T and Tuleun C D. Replacement value of shea nut meal for dietary maize on growth performance and economics of production of rabbits. Nigerian Journal of Animal Production 2020; 47(3): 132-139.
 8. Carew SN, Shaahu DT, Anongo TT. Performance, Apparent nutrient digestibility responses and economics of feeding Nigerian Rice milling by-products to weaner Rabbits. Nigerian Journal of Animal Science. 2020;2(3):184-190.
 9. De Blas C and Wiseman J. The Nutrition of the Rabbit. Second edition. CABI Publishing, Oxon, UK. 2003; 334.
 10. Fielding D. The Tropical Agriculturalist – Rabbits. Macmillan Publishers Limited, Malaysia. 1991;12-29.
 11. Wafar R J, Ndubuisi D I, Tarimbuka L I, Sini T and Oda E. Growth performance and blood profile of weaner rabbits fed different legume haulms supplemented with concentrate. Nigerian Journal of Animal Science 2019;21(3):377-384.
 12. NRC. Nutrients Requirements of Domestic Animals. Nutrients Requirements of Rabbits. 2nd Edition. National Research Council, National Academy of Science. Washington, DC. USA. 1977;2-9.
 13. Bawa GS, Ajide S O, Adeyinka I A. and Ajala M K. Effects of Varying Levels of Groundnut Haulms and Cowpea Shells on Performance of Weaner Rabbits. Asian Journal of Animal and Veterinary Advances. 2008;3:54-61.
 14. Alawa JP. and Amadi. Voluntary feed intake and digestibility of diets containing corn cobs, brewers dried grains and wheat bran by rabbits. Journal of Animal Production Resource. 1991; 11: 9-20.
 15. Syihus B and Hervik A K. The influence of Fibre on Gut Physiology and Feed Regulation. 2020; (<http://internationalfibre.com/>).
 16. Ahemen T, Shaahu D T and Kwaghve N N. Growth performance, Genital organ Dimensions, Visceral organ Weights and Haematology of Male Rabbits Fed Graded Levels of *Gmelina Arborea* Leaf Meal. IOSR Journal of Pharmacy and Biological Sciences (IOSR-JPBS), 2016; 11 (3): 28-32.
 17. Shaahu D T, Ikyuor S A and Carew S. Effect of decortivating and cooking lablab seeds on performance and cost of producing table rabbits. International Journal of Biotechnology and Food Science, 2017; 5(2), pp. 18-22,
 18. Dzungwe, EN, Shaahu, DT, Carew SN, Tuleun, CD. Nutrient Composition and Feed Value of the Fresh Leaves and Succulent Stems of *Ficus thonnini* for Growing Rabbit. Journal of Animal Production Research. 2017;29 (1):175-183.
 19. Sultana N, Huque K S, Rahman M Z and Das N G. Effect of Age on Feed Efficiency and Carcass Yield Characteristics of Indigenous Bull. Bangladih Journal of Animal Science. Journal homepage:<http://www.banglajol.info/index.php/BJAS>. 2003;17-23.
 20. Omole T A, Ajasin F O, Olukun J A, Tiamiyu A K. Feed and feeding. Rabbit Farming without tears. Back to Agriculture (7). Green choice Agricultural publication. 2007;15-18.
 21. Oso A O, Bamgbose A M, Isah O A, Olatunji J E N, Mabadeje A T, Alade A A *et al.*, Performance of weaner rabbits fed rice millings waste-based diets. Journal of Animal and Veterinary Advances, 2006; 5(10): 836-838.
 22. Abdel-Khalek A E, Kalaba Z M and El-Gogary M R. Functional, Anatomical and Histological Development of Caecum in Rabbits. Current Research in Poultry Sciences. 2011;1(2):54-65.
 23. Maertens L and Gidenne T. Feed efficiency in Rabbit Production.: nutritional, Tecnico-economic and Environmental aspects. Feed and Feeding. 11th World Rabbit Congress. 2016, Qingdao, China
 24. Anongo TT. The Effect of Feeding Rice Milling By-Products on Performance, Economics of Production and Blood Profile of Weaner Rabbits. Thesis submitted to the Department of Animal Production, Federal University of Agriculture, Makurdi in partial fulfilment of the requirements for the award of Master of Science in Animal Production and Management; 2017.
 25. Asan N Carcass traits and meat quality properties as affected by nutritional programs and feeding patterns in a rabbit enterprise. Scientific Journal of Animal Science. 2017;6(11):459-470.
 26. Dalle Zotte A. Main factors influencing the rabbit carcass and meat quality.

- Proceedings of the 7th world rabbit congress, 4-7 July, 2000, Valencia, Spain. 2000;507-537.
27. Kagya-Agyemang J K, A Nelson and C. Kagya-Agyemang. Growth Performance of Weaner Rabbits Fed on Dried Pito Mash as A Replacement for Maize. ARPN Journal of Agricultural and Biological Science. 2013;8 (4): 267-272.

© 2023 Anongo et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/107909>