



Effect of Integrated Nutrient Management on Productivity and Quality of Aonla

Ankur Sharma ^{a*}, Anusha K. N. ^b, Priyanka Dahiya ^c
and K. N. Nagaich ^d

^a Rajmata Vijayaraje Scindia Krishi Vishwavidyalay, Gwalior, M.P, India.

^b University of Horticultural Sciences, Bagalkot, India.

^c University of Georgia, Athens, U.S.A.

^d School of Agriculture, ITM University, Gwalior, M.P, 474001, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i224162

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/109730>

Original Research Article

Received: 18/09/2023

Accepted: 23/11/2023

Published: 29/11/2023

ABSTRACT

The field experiment was conducted at the Horticulture Research CRC Farm – 1 of the Department of Horticulture, School of Agriculture, ITM University Gwalior (M.P. during the year 2019 and 2020 to study the effect of Integrated Nutrient Management on fruit yield and quality of aonla. The experiment was laid out in the randomized block design with three replications and eleven treatments viz. (control - RDF (1000: 500:1000 g/ tree, 3/4th of RDF + FYM, 3/4th of RDF + FYM + Azotobacter (100g, 3/4th of RDF + FYM+ Azospirillum (100g, 3/4th of RDF + FYM+ PSB (100g, 3/4th of RDF + FYM+ Azotobacter + Azospirillum + PSB(100g/tree each, 1/2 of RDF + FYM, 1/2 of RDF + FYM+ Azotobacter (100g, 1/2 of RDF + FYM + Azospirillum(100g, 1/2 of RDF + FYM + PSB(100g, 1/2 of RDF + FYM + Azotobacter + Azospirillum + PSB (100g/tree, each. The results revealed that among different treatments, application of 1/2 of RDF + FYM + Azotobacter +

*Corresponding author: E-mail: ank.rvskv@gmail.com;

Azospirillum + PSB recorded higher fruit yield, TSS, Total sugar, Ascorbic acid which was followed by application of $\frac{3}{4}$ th of RDF+ FYM + Azotobacter + Azospirillum + PSB except acidity. Thus, application of $\frac{1}{2}$ of RDF + FYM + Azotobacter + Azospirillum + PSB was found to best for improving the soil nutrient status which will in turn help in improving the yield of Aonla.

Keywords: Farmyard manures; biofertilizers; yield; TSS; total sugar.

1. INTRODUCTION

The Indian gooseberry (*Emblica officinalis* Gaertn., belonging to *Euphorbiaceae* family and is called as Aonla, Amlika, Amali, Ambala, Amalakamu is a native to tropical South – East Asia, is grown in India since its origin [1]. It is widely grown as backyard fruit in India, due to its hardness and wasteland compatibility, high productivity (15-20 tonnes/ha, high nutritive and therapeutic value, it has been considered as an important fruit tree in our country [2]. It is also known as 'Amrit Phal' or 'wonder drug' because of its great medicinal and nutritional utilization. In respect of nutritional values, it is rich source of vitamin C (650-900 mg/100g which is more than that of guava, citrus and tomato fruits and also contains carbohydrates (14.10 - 21.89%, minerals (1.2% Iron, phenol, polyphenol and tannins, alkaloid and ellagic acid. It is a necessary component of Triphala, Chavanprash and other aurvedic preparations [3].

The nutritional value of aonla fruits is also influenced by the fertilization process, soil fertility and sources of nutrition [4]. Tropical soils are deficient in phosphorus and when a farmer adds phosphatic fertilizers, nearly 75% of it is converted to a form unavailable for plant growth. Many fungi and bacteria like *Aspergillus*, *Penicillium*, and *Bacillus* etc. solublize these bound phosphates by producing organic acids and convert them to a form available to a plant growth. Indiscriminate use of inorganic sources of nutrient to provide better nutrition to plants and achieve high yield, made the soil and water quality degraded and brought stagnation in productivity of crops [5]. Integrated use of chemical fertilizers with biofertilizers markedly increases fertilizer use efficiency, minimizes their losses and leakage and improves fertility status of soil [6]. Integrated nutrient management system consists of effective and judicious utilization of all available sources of nutrients to the plants viz., chemical fertilizers, plant and animal sources, organic sources and microbial sources for sustaibale soil fertility and productivity. The increase in crop productivity results from their combined effects, the

synergistic effect, that helps to improve chemical, physical and biological properties of soil and consequently the soil organic matter and nutrient status; to a large extent balanced nutrient supply to crops in cropping system and with no or minimal deleterious effect on environment if any. Considering the role of microbial consortium in soil and as component of INM, study was conducted to evaluate the response of integrated nutrient management practices on productivity and Quality of Aonla cv. NA-7.

2. MATERIALS AND METHODS

The field experiment was conducted during 2019 and 2020 by selecting thirty-three plants of uniform size (canopy volume and vigour from ten-year-old Aonla cultivar 'NA-7' planted with a spacing of 8.0 x 8.0 meter in ten-year-old orchard of aonla cv. NA-7 located at the Department of Horticulture, ITM, University Gwalior, MP (India). The experiment was laid out in randomized block design with eleven treatments and three replications. The treatments are: Full dose of NPK (1000:500:1000 g/tree control; three-fourth dose of NPK/tree + 100kg FYM; three-fourth dose of NPK/tree + 100kg FYM + Azotobacter; three-fourth dose of NPK/tree + 100kg FYM + Azospirillum; three-fourth dose of NPK/tree + 100kg FYM + PSB; three-fourth dose of NPK/tree + 100kg FYM + Azotobacter + Azospirillum + PSB; half dose of NPK/tree +100kg FYM; half dose of NPK/tree +100kg FYM + Azotobacter; half dose of NPK/tree +100kg FYM + Azospirillum; half dose of NPK/tree +100kg FYM + PSB; half dose of NPK/tree +100kg FYM + Azotobacter + Azospirillum + PSB. The treatments include recommended dose of fertilizer (RDF as 1.0kg of nitrogen, 0.5kg of phosphorus and 1.0kg of potassium per tree. Farm yard manure (FYM @ 100 kg/plant along with bio fertilizers was applied around each tree in the second week of January. The Bio fertilizers viz. *Azotobacter*, *Azospirillum* and PSB (100g/tree each were applied in the rhizosphere zone of Aonla around the tree at a depth of 15 cm leaving 50 cm from the main trunk. The NPK fertilizers were applied in form of Urea, SSP, and MOP, respectively. Two third of the total nitrogen

and whole of the phosphorus and potassium were applied during last week of February. Rest one third dose of N was applied in the first week of August. The fertilizers were applied in trenches of 20-25 cm width and 10-15 cm depth made beneath the tree canopy leaving 50 cm distance from the main trunk. The fertilizer was well mixed with the soil in the trenches and then levelled. The observation on the fruit yield was estimated after picking in the second week of December during the year, 2019 and 2020. All the fruits from the individual trees were picked manually and were collected in the baskets. The total weight of the fruits per tree was estimated and expressed in kilogram. Further, the total soluble solids (TSS of the fruit pulp were estimated by using Erma hand refractometer (0-32° B as per the procedure of A.O.A.C (1995 and was expressed in Brix (°B. Acidity of the fruit was calculated by titrating the pulp extract with 0.1 N NaOH as per the method described in A.O.A.C., (1970 and total sugar of aonla fruits was estimated by using Lane and Eynon method given by Ranganna (1986. Ascorbic acid content of the fruits was estimated as per the method suggested by A.O.A.C. (1980 using standardized 2, 6- dichlorophenol indophenols dye. The data were analyzed as per the method suggested by Gomez and Gomez [7]. The critical difference at 5 per cent level of probability and standard error of mean was worked out for comparing the significance among treatment means.

3. RESULTS AND DISCUSSION

3.1 Fruit Yield

The data related to fruit yield of aonla per plant presented in Table 1 revealed that application of $\frac{3}{4}$ th of RDF+ FYM + *Azotobacter* + *Azospirillum* + PSB was substantially better over all the treatments except $\frac{1}{2}$ of RDF+ FYM + *Azotobacter* + *Azospirillum* + PSB and exhibiting maximum fruit yield (159.60 and 161.68 kg/tree. This may be due to the fact that the use of FYM and Bio fertilizers (*Azotobacter*, *Azospirillum* and PSB supplemented the use of inorganic fertilizers to a considerable extent. The application of bio fertilizers along with different dose of NPK and FYM was effective to maintain the nitrogen level of the soil as the microbial population under such treatments was much higher and the fertility of

the soil and enhance yield. The absorbed nitrogen combined with carbohydrates in leaves could lead to the synthesis of amino acids, nucleic acid, proteins, chlorophyll, alkaloid and amides (Jones and Embleton, 1982; Spehia et al. 2020; Singh et al. [8]. These metabolites are involved in building up of new tissues and are related to a number of metabolic steps [9]. Biofertilizers are known to enrich the soil by way of biological N-fixation and improving the availability of different nutrients to plants [6].

3.2 Chemical Properties

The chemical parameters of fruit viz. TSS, acidity, total sugar and ascorbic acid as affected by various treatments was recorded and presented in Table 2 reported that TSS and sugar were highest with the application of half dose of RDF, FYM with *Azotobacter*, *Azospirillum* and PSB. According to Childers (1996 nitrogen has been shown to stimulate the activities of biological enzymes involved in the various bio-chemical processes which might have resulted in an increase in TSS content of fruit with increasing levels of nitrogen. Similar results were reported by Yadav [10] and Yadav [11] in aonla cv. 'Neelam' who noted increasing trend of TSS and total sugar with the application of graded dose of nitrogen. The minimum titratable acidity (2.24 and 2.22% was observed with the application of $\frac{1}{2}$ of RDF+ FYM + *Azotobacter* + *Azospirillum* + PSB and highest in control. similar results were also reported by Yadav [10] and Tewari et al. (2015, Vishwakarma et al. (2017 in acid lime and Singh et al. [12] in muskmelon. The observation on ascorbic acid would reveal that application of $\frac{1}{2}$ of RDF+ FYM + *Azotobacter* + *Azospirillum* + PSB gave highest ascorbic acid content (551.10 and 556.321 mg/100g of pulp. The possible reasons for increased ascorbic acid may be due to conversion of soluble sugars into ascorbic acid and can be confirmed by the findings reported by Tarai and Ghosh (2005 in aonla. Finally, it can be concluded that application of half of of RDF (500:250:500g + FYM + *Azotobacter* + *Azospirillum* + PSB per plant proved to be the most suitable treatment for improving physico-chemical characters of NA-7 aonla fruit [13,14,15].

Table 1. Effect of NPK, FYM and biofertilizers on fruit yield of Aonla cv. NA-7

Treatments	Average fruit yield (kg/tree)	
	2019	2020
Recommended Dose of Fertilizers (RDF (NPK @ 1000:500:1000 g/tree as control	132.60	131.84
¾ th of RDF + FYM	132.76	134.64
¾ th of RDF+ FYM + <i>Azotobacter</i>	146.39	147.56
¾ th of RDF+ FYM + <i>Azospirillum</i>	140.47	140.87
¾ th of RDF+ FYM + Phosphate solubilizing bacteria (PSB	148.75	151.49
¾ th of RDF+ FYM + <i>Azotobacter</i> + <i>Azospirillum</i> + PSB	159.60	161.68
½ of RDF+ FYM	133.69	134.86
½ of RDF+ FYM + <i>Azotobacter</i>	136.96	139.12
½ of RDF+ FYM + <i>Azospirillum</i>	140.47	141.39
½ of RDF+ FYM + PSB	137.22	137.30
½ of RDF+ FYM + <i>Azotobacter</i> + <i>Azospirillum</i> + PSB	155.32	156.92
S.E. (diff	4.255	4.958
CD at 5 % level	8.877	10.343

Table 2. Biochemical quality attributes of aonla fruits after application of biofertilizers as a component of INM

Treatments	TSS (° Brix		Acidity (%)		Total sugar (%)		Ascorbic acid (mg/100 pulp wt.	
	2019	2020	2019	2020	2019	2020	2019	2020
T0	11.50	11.40	2.53	2.58	4.21	4.21	533.86	531.86
T1	11.83	11.92	2.48	2.49	4.22	4.23	534.82	535.20
T2	12.20	12.30	2.40	2.37	4.30	4.32	538.40	539.42
T3	12.33	12.38	2.34	2.31	4.31	4.33	540.12	542.82
T4	12.66	12.90	2.29	2.28	4.36	4.37	544.88	550.66
T5	12.80	13.10	2.28	2.25	4.42	4.46	550.22	555.21
T6	11.75	11.80	2.35	2.32	4.24	4.25	535.39	537.86
T7	12.00	12.33	2.30	2.29	4.28	4.29	539.42	540.49
T8	12.40	12.40	2.35	2.33	4.31	4.34	542.86	543.86
T9	12.90	13.30	2.26	2.24	4.32	4.35	546.80	549.82
T10	13.25	13.35	2.24	2.22	4.62	5.42	551.10	556.21
S.E. (diff	0.41	0.272	0.056	0.026	0.023	0.011	1.769	2.542
CD at 5%	0.855	0.567	0.118	0.0557	0.048	0.0239	3.691	5.303

Table 3. Effect of NPK, FYM and Biofertilizers on total bacterial population in rhizosphere of Aonla plants (Cell x 10⁵ /g Soil)

Treatments	2019				2020			
	December*	March	August	October	December**	March	August	October
T ₀	15.32	26.64	39.42	34.64	32.48	28.45	34.82	25.62
T ₁	16.45	29.32	41.26	36.84	33.66	35.42	44.12	30.80
T ₂	17.07	33.78	49.39	43.24	40.42	42.46	52.18	36.43
T ₃	17.00	30.72	48.34	42.24	36.84	38.60	50.14	35.32
T ₄	17.05	25.42	39.64	33.20	34.58	36.42	45.16	38.36
T ₅	17.10	43.65	59.39	50.24	42.64	48.86	60.18	41.36
T ₆	15.80	39.30	63.30	56.34	32.70	40.84	65.15	33.42
T ₇	22.34	58.63	88.15	85.42	56.42	63.42	92.14	81.46
T ₈	18.65	51.62	86.16	83.62	54.32	58.16	90.64	75.36
T ₉	18.42	50.30	83.14	73.26	51.62	56.18	88.68	71.42
T ₁₀	26.43	79.42	96.14	90.14	63.18	76.42	101.14	91.82
S.Em±	0.894	1.909	2.864	2.764	1.768	2.070	2.928	2.999
C.D at 5% level	1.865	3.98	5.975	5.767	3.689	4.318	6.109	6.257

* Before first inoculation

** Before second inoculation

4. CONCLUSIONS

Thus, application of ½ of RDF + FYM + *Azotobacter* + *Azospirillum* + PSB recorded higher fruit yield, TSS (° Brix, Total sugar (%), Ascorbic acid which was followed by application

of ¾th of RDF+ FYM + *Azotobacter* + *Azospirillum* + PSB except acidity. Thus, application of ½ of RDF + FYM + *Azotobacter* + *Azospirillum* + PSB was found to best for improving the soil nutrient status which will in turn help in improving the yield of Aonla.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Chadha KL. Dry land horticulture crops. Indian Hort. 2001;46:18-20.
2. Kalloo K. Research and extension activities on organic agriculture in India. Organic farming in Horticulture for sustainable production, CISH, Lucknow. 2003;1.
3. Korwar GR, Pratibha G, Ravi V, Palanikumar D. Influence of organic and inorganic on growth, yield of aonla (*Embilica officinalis*) and soil quality in semi and tropics. Indian Journal of Agricultural Sciences. 2006;76(8):457-461.
4. Kulkarani P, Pandey H, Sharma AK, Joshi DC. Physicochemical properties of aonla fruit and juice. Chem. Sci. Rev. Lett. 2017;6:1343-1347.
5. Ramamurthy V, Prasad Jadish. Prasad VN, Gajbhiye KS. Effect of inorganic fertilizer and biofertilizer on productivity of soybean. Biofertilizer News Letter, 2005; 13:18-20.
6. Singh D, Singh SK, Damathia LB. Impact of fertigation on Papaya crop under protected conditions. Ecology Environment and Conservation. 2019;25 (1):295-299.
7. Gomez KA, Gomez AA. Statistical procedures of Agricultural Research (2nd edition), John Wiley and Sons. Inc. New York; 1984.
8. Singh S, Thakur A, Singh SK. Physiological studies on the growth of temperate fruit crops (Walnut and Cherry): A review. Plant Cell Biotechnology Mol. Bio. 2021;22:38-45.
9. Singh SK, Sharma M, Reddy KR, Venkatesh T. Integrated application of boron and sulphur to improve quality and economic yield in potato. Journal of Environmental Biology. 2018;39(2):204-210.
10. Yadav R, Singh HK, Yadav AL. Effect of integrataed nutrient management on productivity and quality of aonla (*Embilica officinalis* Gaertn). Plant Archives, 2006;7 (2):881- 883.
11. Yadav S. Integrated Nutrient Management Studies in subtropical peach cv. Saharanpur Prabhat under high density system. Thesis Ph.D., C.S.A. Univ. Agril. Tech. Kanpur; 2010.
12. Singh V, Dashora LK, Rathore RS. Meena CL. Flowering and yield of guava (*Psidium guajava* L.) cv. "Sardar" as influenced by various organic and inorganic sources. Current Agriculture, 2007;31(1-2):67-71.
13. Ranganna S. Hand book of analysis and quality control of fruit and vegetable products. New Delhi, Tata Mcgraw Hill Publishing Company Ltd. 1986:253.
14. Singh SK, Sharma M. Evaluating Significance of Vermicompost and Intercropping *Amorphophallus* for integrated Indian Goose Berry Orchard Management. International Journal of Agriculture Sciences. 2016;8(39):1809-1812.
15. Vedamani GM, Ramavatharam N, Haribabu K. Effect of graded levels of nitrogen through organic and inorganic sources on yield and quality of acid lime (*Citrus aurantifolia* Swingle). The Orissa J. Hort. 2006;34(1): 49-51.

© 2023 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/109730>