



Influence of Nitrogen and Phosphorus Fertilizers on Bulb Yield and Yield-Related Attributes of Onion (*Allium cepa* L.) in Fogera Area, Northwest Ethiopia

Abebe Getu Asfaw ^{a*}, Wubayehu Gebremedhin Woldie ^a
and Demsew Bekele Gelagil ^a

^a Ethiopian Institute of Agricultural Research, Fogera National Rice Research and Training Center, P.O. Box 1937, Bahir Dar, Ethiopia.

Authors' contributions

This work was carried out in collaboration among all authors. Authors AGA, WGW and DBG conceived and designed the research. Author AGA collected and analyzed the data and wrote the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJSSPN/2024/v10i1236

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/112899>

Original Research Article

Received: 07/12/2023
Accepted: 13/02/2024
Published: 02/03/2024

ABSTRACT

Aim: Onion is one of the most important vegetable crops grown under irrigation in the Amhara Region of Ethiopia. However, its productivity is affected by sub-optimal fertilization. The present research was conducted to investigate the response and economic returns of onion to N and P fertilizers application under irrigation.

Study Design: Randomized Complete Block Design in Factorial Arrangement with three replications.

*Corresponding author: Email: abegetu3@gmail.com;

Place and Duration of the Study: The research was conducted in 2020 and 2021 in the Fogera area of Northwest Ethiopia.

Methodology: Four levels of N (0, 46, 92, and 138 kg N ha⁻¹) and three levels of P (0, 46, and 92 kg P₂O₅ ha⁻¹) were factorially combined and laid in a randomized complete block design with three replications.

Results: The results showed that the yield and yield components of onion were significantly affected by the main effects of N. However, the main effects of P and its interaction with N did not significantly influence the growth and yield of onion. The highest plant height, number of leaves per plant, bulb girth and marketable bulb yield were recorded from the application of N₁₃₈ statistically at par with N₉₂. The maximum net economic return was obtained from the application of N₁₃₈.

Conclusion: Application of 138 kg N ha⁻¹ was the economic optimum rate for maximum bulb yield of onion in the study area. Further investigation on the response of onion to P is recommended.

Keywords: Fogera; Irrigation; nitrogen; onion; optimum fertilizer rate; phosphorus.

1. INTRODUCTION

“Onion (*Allium cepa* L.) is a vegetable crop grown for its pungent bulbs and flavorful leaves, and it significantly contributes to the nutritional requirements of human beings” [1]. “It is one of the most important vegetables in Ethiopia. As a bulb crop, it is mainly produced by smallholder farmers as a source of cash income and for flavoring the local stew “wot”. However, the productivity of onion in Ethiopia (8.9 t ha⁻¹) is low compared to the average yield in Africa (14.5 t ha⁻¹) and the world average (19.3 t ha⁻¹)” [2,3]. “Different studies have indicated that onion production in Ethiopia is constrained by poor agronomic practices, including unbalanced fertilization, shortage of seeds of improved varieties, diseases, insect pests, and high costs of agricultural chemicals, including fungicides and insecticides” [4,5]. According to [6] and [7], “depleted soil fertility and poor agronomic practices are among the major factors contributing to low vegetable productivity in Ethiopia”. “Onion is more susceptible to nutrient deficiencies than most other crop plants because of its shallow and unbranched root systems; hence, they require and often respond well to the addition of fertilizers” [8].

“In the Fogera area, which is located in the South Gondar Zone, Northwest Ethiopia, small-scale onion production in the dry season under irrigation is the most common and is the major source of onion for the local market in the surrounding towns. Accordingly, growers, merchants, consumers, intermediaries, and transporters highly benefit from onion production in the area. However, the production and productivity of the crop in this area are constrained by different biotic and abiotic stresses, among which low soil fertility and sub-

optimal fertilization are the most important ones” [6-7]. “Adequate management of mineral nutrition is a determining factor for high yield and profitability of the crop” [9]. There were no location-specific N and P fertilizer recommendations for onion production in the Fogera area. Therefore, this study was conducted to determine the response of onion to the application of N and P fertilizers and determine the optimum economic rate of N and P fertilizers to maximize the productivity of onion under irrigation in the Fogera area.

2. MATERIALS AND METHODS

The study area is known for rainfed lowland rice cultivation. Following the rice harvest, onion is cultivated by smallholder farmers under irrigation. Most of the smallholder farmers use rivers as a source of irrigation water. Some smallholder farmers dug wells manually and use groundwater as a source of irrigation.

2.1 Description of the Study Area

The study was conducted at Fogera National Rice Research and Training Center (FNRRTC) in 2020 and 2021 under irrigation. FNRRTC is located in the Fogera District of Amhara region, Northwest Ethiopia. It is situated at 11° 54' 22.84"N latitude and 37° 41' 9.97"E longitude at an altitude of 1806.4 meters above sea level (masl) (Fig. 1). Fogera district is located between latitude 11°49'55" N and longitude 37° 37' 40" E, with altitudinal ranges of 1774 and 2415 masl. The dominant soil types in the study area are Pellic Vertisols.

2.2 Climate

The study area lies in a tropical monsoon climate, where the rainfall is dominated by the

Intertropical Convergence Zone (ITCZ). The climate of the study area is locally classified as “*woina dega*” (intermediate highland between 1500-2300 masl) and it lies in tepid moist mid highlands agro-ecological zone based on the classification by [10,11]. Rainfall in the area is uni-modal, usually occurring from June to September (Fig. 2), and

its average annual total rainfall is 1363.7 mm (with 90% falling from June to September). The mean minimum and maximum temperature of the study area are 12.7 and 27.4°C, respectively with the lowest occurring in December and January and the highest in Feb to May (Data obtained from Woreta Metrological Station).




Fig. 1. Location map of the study area (Fogera district) and the study site, FNRRTC research station




Fig. 2. Monthly total rainfall, monthly mean minimum (Tmin) and maximum (Tmax) temperatures of the study area

2.3 Soil Sampling and Analysis

Five soil samples were collected before transplanting diagonally following 'X' pattern at five spots at a depth of 0-20 cm using auger. Those five soil samples were composited to form one representative soil sample, air-dried, and ground to pass a 2 mm sieve for laboratory analysis. The soil samples were analyzed for pH, organic carbon (OC), total nitrogen (TN), available P, and Cation Exchange Capacity (CEC) using the standard soil testing procedure at the Amahara Design and Supervision Works Enterprise in Bahir Dar, Ethiopia.

2.4 Treatments and Experimental Procedures

Four levels of N (0, 46, 92, and 138 kg N ha⁻¹) and three levels of P (0, 46, and 92 kg P₂O₅ ha⁻¹) were combined in a factorial arrangement and laid in a randomized complete block design with three replicates. The onion variety Bombay Red, which is a widely grown variety in the study area, was used in the study. Onion seedlings were raised in October and November in the two experimental years (2019 and 2020) in a nursery at the FNRRTC research station on well-prepared seed beds. Fifty (50) days old onion seedlings were transplanted to the experimental plots with furrow and ridge spacings of 40 cm and 20 cm, respectively. The seedlings were transplanted on both sides of the ridges with 7 cm spacing between plants. The gross and net size of the experimental plots were 4.2 m * 3.0 m (12.6 m²) and 3.00 m * 2.94 m (8.82 m²), respectively. Triple Super Phosphate (TSP) was used as a source of P fertilizer and was band-applied as basal during transplanting. Urea was used as a source of N fertilizer and was applied in split, half was applied in a band at transplanting, and the remaining half was side-dressed 45 days after transplanting. The experimental plots were uniformly furrow-irrigated up to saturation with six days (6) days intervals in the initial and development growth stages, and in 5 days intervals in the mid and late growth stages of the crop. All the other agronomic management practices were implemented uniformly in all plots as per the recommendation.

2.5 Data Collection

Data on growth, bulb yield, and yield-related parameters of onion were measured and recorded at physiological maturity from the net

plot area of 8.82 m² using their respective standard procedures described below.

2.5.1 Plant height (cm): was obtained by measuring five randomly selected plants from the ground to the tip of the longest mature leaf using a ruler at physiological maturity.

2.5.2 Number of leaves per plant: number of leaves per plant was measured by counting the total number of leaves of five randomly selected plants from the net plot area, and the mean value was used for the analysis.

2.5.3 Bulb diameter (cm): was measured from five randomly selected matured bulbs at harvest using a vernier caliper.

2.5.4 Marketable bulb yield (kg ha⁻¹): "Bulbs that were free of mechanical, disease, and insect pest damage, uniform in color, and medium to large in size (20-160 g) were considered marketable" [6]. The weight of the bulbs obtained from the net plot area was measured in kilograms using a scaled balance and expressed as kilograms per hectare.

2.5.5 Unmarketable bulb yield (kg ha⁻¹): "Bulbs under (<20 g) and oversized (>160 g), decayed, discolored, diseased, and physiologically disordered bulbs are considered unmarketable" according to [6]. The weight of the bulbs obtained from the net plot area was measured in kilograms using a scaled balance and expressed as kilograms per hectare.

2.5.6 Total bulb yield (kg ha⁻¹): total bulb yield of onion was obtained by adding marketable and unmarketable bulb yields.

2.6 Data Analysis

"The agronomic data collected were subjected to a two-way analysis of variance using the SAS Statistical Software package version 9.2" [12]. Means were separated using the Least Significant Difference (LSD) test method at a 5% level of significance. A combined analysis of the variance of the agronomic data over the two years was performed after testing and confirming the homogeneity of error variances in the two years. In the combined analysis, year was considered a random variable, while treatment was considered a fixed variable.

2.7 Economic Analysis

Partial budget analysis was performed following the procedure described by [13]. Based on the

data obtained from the Fogera District Trade and Industry Bureau, the farm gate price of onion bulb yield was 12.33 birr kg⁻¹ and the cost of urea fertilizer was 14.38 birr kg⁻¹. The mean marketable bulb yield used for the partial budget analysis was adjusted to 90% of the measured yield. The most profitable treatment was selected based on the highest net economic return, with a marginal rate of return above 100%.

3. RESULTS AND DISCUSSION

3.1 Initial Soil Fertility Status of the Study Site

The soil analysis results of the surface soil samples collected from the study site before establishing the experiment showed that the surface soil was clay in texture with a very high [14] cation exchange capacity and high exchangeable K content [15]. The surface soil was slightly acidic (pH-H₂O 6.2) in soil reaction with low [16] organic carbon, low [16] total N, and medium [17] available P contents (Table 1).

3.2 Effect of N and P on the Growth, bulb Yield and Yield Components of Onion

3.2.1 Plant height

The plant height of onion was significantly ($p < 0.01$) affected by the main effect of N in the first and second experimental years (Table 2). Pooled analysis over the two experimental years showed that the tallest plants were recorded from the highest rates of N (N₉₂ and N₁₃₈). Increasing the rate of N from 0 to 92 increased the plant height by 14.6%. The positive effect of N on plant height might be attributed to the role of N in improving the leaf area index and chlorophyll content, thus resulting in a higher photosynthetic rate and vegetative growth [18,19]. The results of the present study are supported by those of [20] and [21], who reported that increasing N levels up to 103.5 kg ha⁻¹ increased plant height, but

the application of N beyond 103.5 kg ha⁻¹ decreased plant height. [22] also revealed that nitrogen fertilizer rates (0, 46, 69, 92, 115, and 138 kg ha⁻¹) significantly affected plant height of onion. However, in the present study, the main effect of P and its interaction with N did not significantly ($p \geq 0.05$) influence the plant height. This result is in agreement with those of [21] and [23], who reported the absence of significant differences in plant height owing to the application of P fertilizer.

3.2.2 Leaf number per plant

The number of leaves per plant was significantly ($p < 0.01$) affected by the main effect of N in the first experimental year (Table 2). Although not statistically significant ($p \geq 0.05$), the pooled analysis over the two experimental years indicated that the application of N₁₃₈ resulted in the highest leaf number per plant (10.2), exceeding the leaf number per plant recorded from the N₀ treatment by 14.6%. In accordance with the present research findings, [23] reported an increased number of leaves due to the application of higher rates of N. [22] reported a significant effect of N fertilization up to a rate of 138 kg ha⁻¹ on the leaf number and plant maturity of onion. This could be attributed to increased vegetative growth associated with the role of N in the production of proteins and amino acids required for leaf growth and development [19]. Nasreen et al. [24] also found a significant improvement in the number of leaves as the rate of N application increased to 120 kg ha⁻¹, which declined as the level of N increased to 160 kg ha⁻¹. However, the main effect of P and its interaction with N did not significantly ($p > 0.05$) affect the leaf number in either experimental year. This result is supported by [23], who reported a non-significant difference in the number of onion leaves due to the application of P. However, in contrast, [21] reported significant differences in the number of onion leaves in response to P application.

Table 1. Some soil physico-chemical properties of surface soil (0-20 cm) of the study site

Parameter	Values	Rating
pH (H ₂ O)	6.2	Slightly acidic
Organic carbon (%)	1.39	Low
Total N (%)	0.12	Low
P _(Olsen) (mg kg ⁻¹)	17.2	Medium
Exchangeable K (Cmol _c /kg)	0.61	High
Cation Exchange Capacity (Cmol _c kg ⁻¹)	46.1	Very high
Soil texture (%) (Sand, Silt and Clay)	18, 28, and 54	Clay

3.2.3 Bulb diameter

The influence of N on bulb diameter was highly significant ($p < 0.01$) in the two experimental years (Table 2). The highest bulb diameter (5.4 cm) was recorded from N_{138} , at par with N_{92} . The lowest bulb diameter was recorded in N_0 . The pooled analysis over the two years indicated that the application of N_{138} increased the bulb diameter by 5.9% compared to the N_0 treatment. In line with the present result, an increase in bulb diameter by about 12 and 16% due to application of 138 and 150 kg N

ha^{-1} over the control was reported by [23]. Similarly, [21] reported a significant difference in bulb diameter due to application of N and P. However, the main effect of P and its interaction with N had no significant ($p \geq 0.05$) effect on bulb diameter. In agreement with our results, [23] also reported a non-significant difference in bulb girth of onion due to the application of P fertilizer. In accordance with the present results, highest (6.6 cm) bulb diameter of onion was recorded from application of N fertilizer at the rate of 103.5 kg ha^{-1} with 30 t ha^{-1} farm yard manure [25].

Table 2. Main effects of N and P fertilizers on growth and yield-related attributes of onion as affected by the main effects of N and P fertilizers in the 2020 and 2021 experimental years

N level* (kg N ha^{-1})	2020			2021			Pooled		
	PH	LN	BD	PH	LN	BD	PH	LN	BD
N_0	40.9 ^b	9.3 ^b	5.6 ^c	39.0 ^c	8.4	4.6 ^c	39.7 ^c	8.9	5.1 ^c
N_{46}	41.0 ^b	10.7 ^a	5.7 ^b	41.8 ^b	9.1	4.8 ^b	41.4 ^b	10.1	5.3 ^b
N_{92}	45.3 ^a	10.7 ^a	5.9 ^a	45.6 ^a	8.6	5.0 ^a	45.5 ^a	10.1	5.4 ^a
N_{138}	42.1 ^b	11.0 ^a	5.8 ^{ab}	47.4 ^a	8.8	5.1 ^a	45.1 ^a	10.2	5.4 ^a
LSD (5%)	1.5	1.0	0.14	2.2	NS	0.11	1.4	NS	0.08
P level* (kg P_2O_5 ha^{-1})									
P_0	41.4 ^a	10.9	5.7	44.2	8.7	4.8 ^b	42.9 ^{ab}	10.1	5.3
P_{46}	44.5 ^a	10.5	5.7	43.8	9.4	5.0 ^a	44.1 ^a	10.1	5.3
P_{92}	40.8 ^b	10.1	5.8	42.4	8.3	4.4 ^b	41.7 ^b	9.3	5.3
Mean	42.3	10.5	5.7	43.5	8.7	4.9	43	9.8	5.3
CV (%)	3.1	9.8	2.2	5.2	11	2.3	4.6	10.1	2.2
LSD (5%)	1.3	NS	NS	NS	NS	0.1	1.2	NS	0.1
N level * P level	NS	NS	NS	NS	NS	NS	NS	NS	NS

Note: PH: Plant height (cm); LN: Leaf number per plant; BD: Bulb diameter (cm); CV: Coefficient of variation; LSD: Least significant difference. *Treatments within a column followed by the same letter are not significantly different at 5% probability level. NS: Non-significant at the 5% significance level

Table 3. The marketable bulb yield of onion as affected by the main effects of application of N and P fertilizers

N level* (kg N ha^{-1})	Marketable bulb yield (kg ha^{-1})		
	2020	2021	Pooled
N_0	14519.6 ^c	11713 ^d	13033.6 ^d
N_{46}	15779.9 ^{bc}	16071 ^c	15934.0 ^c
N_{92}	17346.9 ^{ab}	20698 ^b	19232.1 ^b
N_{138}	18185.1 ^a	23763 ^a	21137.9 ^a
LSD (5%)	1795.8	2509.5	1521.4
P level* (kg P_2O_5 ha^{-1})			
P_0	17002.4	17649	17339.5
P_{46}	15721.4	18900	17379.9
P_{92}	16593.8	17635	17188.7
Mean	16429.2	18061.2	17306.1
CV (%)	10.2	14.2	12.5
LSD (5%)	NS	NS	NS
N level * P level	NS	NS	NS

Note: CV: Coefficient of variation; LSD: Least Significant Difference. *Treatments within a column followed by the same letter are not significantly different at 5% probability level. NS: Non-significant at the 5% significance level

3.2.4 Bulb yield

The marketable bulb yield of onion was significantly ($p < 0.01$) affected by the main effect of N in the two experimental years (Table 3). However, the main effect of P and its interaction with N did not significantly influence ($p \geq 0.05$) the bulb yield of onion. Pooled analysis over the two experimental years indicated that the highest marketable bulb yield (21.1 t ha^{-1}) was obtained from N_{138} . The lowest marketable bulb yield (13.0 t ha^{-1}) was obtained in the N_0 treatment. The marketable bulb yield increased by 62.2% owing to the application of N_{138} compared to the marketable bulb yield obtained from the N_0 treatment.

In agreement with the present research results, [26] revealed that application of 138 kg N ha^{-1} was the best treatment in terms of yield and yield components of onion. The present research finding was also supported by [27] who reported a significant marketable bulb yield increase from the application of 138 kg N ha^{-1} with the highest level of irrigation. A study conducted around the Rib area in Fogera district by [28] revealed application of 150 kg N ha^{-1} and $46 \text{ kg P}_2\text{O}_5 \text{ ha}^{-1}$ as an economically optimum rate for onion production. A research report on integrated nutrient management for onion by [29] showed an increased bulb yield of onion as a result of applications of higher levels of inorganic and

organic fertilizers. The significant yield improvement in response to higher rates of N could be due to the enhancement of growth and production of assimilates and their partitioning into the bulbs [30]. Overall, the results from the present study and research findings from various studies revealed that good growth performance of onion was obtained from N fertilization with 138 kg N ha^{-1} . However, there was no significant ($p \geq 0.05$) yield difference from the zero P (P_0) due to the application of P_{46} and P_{92} , which might be because of the medium level available P content of the surface soil of the study site.

3.3 Yield Response to N

Marketable bulb yield increased with increasing levels of N fertilizer and it was best expressed by a quadratic response curve ($r^2 = 0.993$ and 0.998 in 2020 and 2021, respectively), as shown in Fig. 3. Excessive N application decreases economic returns and increases the risk of groundwater nitrate contamination. Furthermore, a high fertilization rate leads to saline soil problems, and onion is sensitive to soil salinity. Salt damage to young onion plants ultimately has an impact on yield [31,32]. Different studies have revealed that excessive N fertilization reduces bulb dry matter content and storability by enhancing sprouting and rotting percentages [33].




Fig. 3. Bulb yield response of onion to N fertilizer in the 2020 and 2021, where Y is marketable bulb yield and x is the level of N applied in kg N ha^{-1}

3.4 Yield Response to P

The bulb yield response to P in the two experimental years was not significant, regardless of the rate of P application (0, 46, and 92 kg P₂O₅ ha⁻¹) (Fig. 4). This might be attributed to the medium level of available soil P in the surface soil of the study site. Moreover, the non-significant yield response to P might be due to the fact that, with regard to P, onion crop is inefficient in the extraction of P from the soil because the root hairs are mostly shorter than the length of phosphate diffusion. This result is similar with findings of [28], where the effect of P and its interaction with N was found statistically non-significant on the yield of onion. According to [34], special care from growers is required with respect to the levels of this nutrient in the soil, as well as the sources and forms of application of phosphate-based fertilizers. Furthermore, the response of onion to P

fertilization depends on the genotype used, P level in the soil, P source, soil, and weather conditions [35]. Therefore, further studies with higher rates of P need to be conducted for the onion in the study area.

3.5 Partial Budget Analysis

The partial budget analysis indicated that the highest net economic return of Ethiopian currency (Birr) 230,252.4 was obtained from the application of N₁₃₈ followed by the net economic return of Birr 210,542.1, which was obtained from the application of N₉₂ (Table 4). There was a rate of return of 137.0 and 244.5 birr for every birr invested on onion production by application of 138 and 92 kg N ha⁻¹, respectively. The result is supported by the findings of [28] who found out application of 150 kg N ha⁻¹ as an economic optimum rate of N fertilizer for onion production.




Fig. 4. Yield (Marketable bulb yield) response of onion to application of P fertilizer in 2020 and 2021

Table 4. Partial budget analysis for the mean effect of N fertilizer on the marketable bulb yield of onion in Fogera area of Northwest Ethiopia

N (kg ha ⁻¹)	MBY (kg ha ⁻¹)	Adj. MBY (kg ha ⁻¹)	Gross return	Production cost	Net return	MRR (%)
N ₀	13033.6	11730.2	144633.9	0	144633.9	-
N ₄₆	15934.0	14340.6	176819.6	1438	175381.3	2137.8
N ₉₂	19232.1	17308.9	213418.6	2877	210542.1	2444.6
N ₁₃₈	21137.9	19024.1	234567.3	4315	230252.4	1370.4

Note: MBY: Marketable bulb yield, Adj. MBY: Adjusted Marketable bulb yield, MRR: Marginal Rate of Return (%). All costs are in Ethiopian currency (Birr)

4. CONCLUSION

The results of the study revealed that the growth, yield, and yield components of onion were significantly influenced by the main effects of N. However, the application of P fertilizer and its interaction with N did not significantly affect the growth and yield of onion. Application of N₁₃₈ was found superior in the plant height, number of leaves per plant, bulb diameter, and marketable bulb yield recorded. Based on the partial budget analysis, the highest net economic return was obtained from N₁₃₈ followed by N₉₂. Therefore, the application of 138 kg N ha⁻¹ can be recommended for maximum bulb yield and maximum net economic return of onion production in the study area and similar agro-ecologies. However, further investigation on the response of onion to P should be done.

ACKNOWLEDGEMENTS

The authors acknowledge the Ethiopian Institute of Agricultural Research for financing the research. We would like to extend our gratitude to the Fogera National Rice Research and Training Center (FNRRTC) for providing the facilities to undertake this research. We are also grateful to the Amhara Design and Supervision Works Enterprise for conducting the soil analysis.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ketter CAT, Randle WM. Pungency assessment in onions. In: S. J. Karcher (Ed.). Tested studies for laboratory teaching. Association for Biology Laboratory Education (ABLE). 1998;19: 177-196.
2. FAOSTAT (Food and Agriculture Organization). Agriculture in the world. Top country of onion producer area and production of major tables of Central Statistical Agency. Available: www.potato.org.
3. CSA (Central Statistical Agency). Report on area and production of major crops (Private peasant holdings, meher season). The federal democratic republic of Ethiopia central statistical agency agricultural sample survey 2020/21. Vol. I. Statistical bulletin 589; 2015.
4. Kumilachew AK, Mengistu G Fekadu. Risks in vegetable production from the perspective of smallholder farmers: the case of Kombolcha Woreda, Oromia region, Ethiopia. Agric. For. Fish. 2014;3(6-1):1-5.
5. Alemayehu M, Minwuyelet J. Optimum rates of NPS fertilizer application for economically profitable production of potato varieties at Koga Irrigation Scheme, Northwestern Ethiopia. Cogent. Food and Agric. 2018;4:1.
6. Lemma D, Shimelis A. Research experience in onion production. Research Report No 55, Ethiopian Agricultural Research Organization. Addis Ababa, Ethiopia; 2003.
7. Gebretsadik K, Dechassa N. Agronomic and economic evaluation of nitrogen fertilizer rates and intra row spacing on growth and bulb yield of onion (*Allium cepa* L.) under rainfall condition. Environment. 2016;6(21):1–10.
8. El-hamady, M. M. Growth and Yield of Onion *Allium cepa* L. as Influenced by Nitrogen and Phosphorus Fertilizers Levels. Canadian Journal of Agriculture and Crops. 2017;2(1):34-41.
9. Marschner H. Mineral Nutrition of Higher Plants, 2nd ed. Academic press. London. 2015;196.
10. MoARD (Ministry of Agriculture and Rural Development). Major Agro-ecological Zones of Ethiopia. Forestry, Land Use, and Soil Conservation Department. Ministry of Agriculture and Rural Development, Addis Ababa; 2005.
11. Belete B, Assefa M, Yilma S. GIS-based hydrological zones and Soil Geo-Database of Ethiopia. Catena. 2013;104: 21-31.
12. SAS/STAT Software, Version 9.2 of the SAS System for Windows. SAS Institute, Cary, NC; 2008.
13. CIMMYT (International Maize and Wheat Improvement Center). From Agronomic data to Farmers Recommendations: An economic training manual. Revised Edition. 1988;200. Mexico DF.
14. Hazelton P, Murphy B. Interpreting soil test results: What do all the numbers

- mean? 2nd Edition. CSIRO Publishing. 2007;152.
15. FAO (Food and Agriculture Organization). Plant nutrition for food security: A guide for integrated nutrient management. FAO, Fertilizer and Plant Nutrition Bulletin 16, Rome, Italy; 2006.
 16. Tekalign T. Soil, plant, water, fertilizer, animal manure and compost analysis. Working Document No. 13. International Livestock Research Center for Africa, Addis Ababa, Ethiopia; 1991.
 17. Cottenie A. Soil and plant testing as a basis of fertilizer recommendations. FAO Soil Bulletin 38/2. Food and Agriculture Organization of the United Nations. Rome, Italy; 1980.
 18. Eastin JD, Haskins FA, Sullivan CY, Van Bavel CHM, Eds. Physiological aspects of crop yields. American Society of Agronomy. Madison, USA: Crop Society of America. 1969;253.
 19. Havlin JL, Beaton JD, Tisdale SL, Nelson WL. Soil Fertility and fertilizers. An introduction to nutrient management, PHI Learning Private Limited, New Delhi, India, 7th edition; 2010.
 20. Kumar H, Singh JV, Ajay K, Mahak S, Kumar A, Singh M. Studies on the influence of nitrogen on growth and yield of onion Cv. Patna Red Indian J. Agric. Res. 1998;32:88-92.
 21. Negasi T, Nigussie D, Kebede W, Lemma D, Abuhay T. Influence of Nitrogen and Phosphorus Application on Bulb Yield and Yield Components of Onion (*Allium cepa* L.). The Open Agric. J. 2018;12:194-206.
 22. Weldemariam SG, Woldetsadik K, Mohammed W. Growth parameters of onion (*Allium cepa* L. var. Cepa) as affected by nitrogen fertilizer rates and intra-row spacing under irrigation in Gode, South-Eastern Ethiopia. Agric. Forestry and Fisheries. 2015;4(6):239-245.
 23. Abdissa Y, Tekalegn T, LM. Pant. Growth, bulb yield and quality of onion (*Allium cepa* L.) as influenced by nitrogen and phosphorus fertilization on Vertisol I: Growth attribute biomass production and bulb yield. Afr. J. Agric. Res. 2011;6(14): 3252-8.
 24. Nasreen S, Haque MM, Hossain MA, Farid ATM. Nutrient uptake and yield of onion as influenced by nitrogen and sulphur fertilization. Bangladesh J. Agric. Res. 2007;32(3):413-420.
 25. Lchi AG, Negasi TA, Tesfay BR. Growth and yield of onion (*Allium cepa* L.) as affected by farmyard manure and nitrogen fertilizer application in Tahtay Koraro District, Northwestern Zone of Tigray, Ethiopia. Int. J. Plant Res. 2020;33:617-627.
 26. Gessesew WS, Woldetsadik K, Mohammed W. Effect of nitrogen fertilizer rates and intra-row spacing on yield and yield components of onion (*Allium cepa* L. Var. cepa) under irrigation in Gode, South-Eastern Ethiopia. Int. J. Plant Breeding Crop Sci. 2015;2(2): 46-54.
 27. Gebregwergis FH. Effect of irrigation depth and nitrogen levels on growth and bulb yield of onion (*Allium cepa* L.) at Algae," M.Sc Thesis, University of Haramaya, Dire Dawa, Ethiopia; 2014.
 28. Agumas B, Abewa A, Abebe D. Response of Irrigated Onion (*Allium cepa* L.) to Nitrogen and Phosphorus Fertilizers at Ribb and Koga Irrigation Schemes in Amhara Region, North western Ethiopia. Int. Res. J. Agric. Sci. Soil Sci. 2014;4(5):95-100.
 29. Bagali AN, Patil HB, Chimmad VP, Patil PI, Patil RV. Effect of Inorganics and Organics on Growth and Yield of Onion (*Allium Cepa* L.). Karnataka J. Agric Sci. 2012;25(1): 112–115.
 30. Girma Z. Response of onion (*Allium cepa* L. Var. Cepa) to organic and inorganic fertilizers at Gode, South-Eastern Ethiopia. Doctoral Dissertation, School of Graduate Studies, Haramaya University, Ethiopia; 2011.
 31. Allen RG, Pereira LS, Raes D, Smith M. Crop Evapotranspiration Guidelines for computing crop water requirements, 56, FAO Irrigation and Drainage Paper, Rome; 1998.
 32. Daniel G, Roberto SO, Jairo D. Nitrogen nutrition and fertilization of onion (*Allium cepa* L.). A literature review. Scientia Horticulturæ. 2022;291:3.
 33. Tekalign T, Abdissa Y, Pant LM. Growth, bulb yield and quality of onion (*Allium cepa* L.) as influenced by nitrogen and phosphorus fertilization on vertisol. II: Bulb

- quality and storability. African J. of Agric. Res. 2012;7(45):5980-5985.
34. Brewster JL. Onion and other vegetable Alliums. CAB International, Wallingford, UK; 1994;236.
35. Grant C, Bittman S, Montreal M, Plenchette C, Morel C. Soil and fertilizer phosphorus: Effects on plant P supply and mycorrhizal development. Canadian J. Plant Sci. 2005;85:3-14.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/112899>