



Evaluation of Growing Media and Chemical Treatment on the Seedling Growth of High Chill Peach

**Ayushi Jain ^{a*}, D.C. Dimri ^a, Monu Kumari ^b, Visha Jain ^c,
Harshit Kumar ^b, Sunil ^d, Mangi Lal ^e
and Ram Lakhan Meena ^b**

^a Department of Horticulture, College of Agriculture, GBPUA&T, Pantnagar, Uttarakhand, 263145, India.

^b Department of Horticulture, Rajasthan College of Agriculture, MPUAT, Udaipur, Rajasthan, 313001, India.

^c Department of Forestry, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana-125004, India.

^d Department of Vegetable Science, Chaudhary Charan Singh Haryana Agricultural University, Hisar, Haryana-125004, India.

^e Department of Vegetable Science, College of Horticulture, Sardarkrushinagar Dantiwada Agricultural University, Jagudan, Mehasana, Gujarat-382710, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i113031>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/126364>

Original Research Article

Received: 01/09/2024

Accepted: 02/11/2024

Published: 08/11/2024

*Corresponding author: E-mail: jayushi797@gmail.com;

ABSTRACT

The present study entitled “Evaluation of Growing Media and Dormancy Breaking Chemical on the Seedling Growth of High Chill Peach” was conducted at Dept. of Horticulture, College of Agriculture and Horticulture Research Centre, G.B.P.U.A&T, Pantnagar, Uttarakhand during the year 2022-23. The experiments were carried out in Factorial Complete Randomized Design (FCRD) with 3 replications per treatment. Under seedling growth studies, the total numbers of treatment were 16 with 2 factors comprising of four growing media (soil: sand: FYM, soil: sand: vermicompost, soil: sand: cocopeat and soil: FYM: vermicompost: cocopeat) and four chemical treatments (control, GA₃ @ 500 ppm, kinetin @ 50 ppm and thiourea @ 5000 ppm). In the experiment the maximum seedling length, shoot fresh weight, shoot dry weight, root length, root fresh weight and root dry weight was recorded in soil: FYM: vermicompost: cocopeat, GA₃ @ 500 ppm and interaction M₄C₃ (soil: FYM: vermicompost: cocopeat + GA₃ @ 500ppm) while the minimum values were recorded in soil: sand: cocopeat, control and M₃C₁. Result from this study discovered that GA₃ @ 500 ppm along with the growing media soil: FYM: vermicompost: cocopeat recorded the maximum seedling length, diameter, number of leaves, leaf area, shoot fresh weight and shoot dry weight which can be recommended for raising the rootstocks for the production of quality planting material of peach for seedling rootstock.

Keywords: Thiourea; vermicompost; kinetin and cocopeat.

1. INTRODUCTION

Prunus persica (L.) Batsch, a member of the Rosaceae family, is the third most important fruit crop grown worldwide in temperate and subtropical regions, after apples and pears. These are extremely nutrient-dense fruits that are abundant in sugar, protein, vitamins and minerals. Peach cultivation is most successful in latitudes between 30° and 45°. Cultivars are classified as high or low chilling according on the rate of chilling they require. For normal blooming and fruiting, high chill peaches typically require 500–1000 chilling hours below 7.2°C. They are cultivated in the Himalayan regions of Uttarakhand, Jammu and Kashmir.

Many temperate fruit seeds are dormant when they are harvested, this means that even under ideal conditions, the embryo will not germinate, possibly because of a hard seed coat or the presence of growth inhibitors. Dormant seeds contain a lot of growth inhibitors, during stratification, their concentration, nature, or both are altered Lipe and Crane, Lipe and Crane, 1966 and Diaz and Martin, 1972. Abad et al. 2002, Agbo and Omaliko, 2006 and Wilson et al., 2001 reported that ABA levels in embryonic axes declined during stratification and use of growth regulators like GA₃, kinetin and thiourea decrease ABA levels, enhancing germination and seedling growth. GA₃ stimulates enzyme synthesis, weakening the endosperm and mobilizing enzyme reserves (Hota et al., 2018,

Imani et al. 2011, Abou et al. 2021 and Parvin et al. 2015). Cytokinins aid in nutrient mobilization and plant growth (El-Badawy and El-Aal 2013). Thiourea breaks dormancy and enhances stress tolerance, gene expression and signaling processes, making it effective under environmental stress (Chang and Sung, 2000; El-Keblawy, 2013; Pallavi et al., 2022; Cetinbus and Koyuncu 2006 and Thakur and Singh 2015).

To produce high-quality nursery plants, a suitable growth medium must be used. It has an immediate impact on the germination, growth and maintenance of the large, well-functioning roots system. According to Abad et al. Abad et al. (2002), a robust growth medium would provide the plant with enough anchoring or support, serve as a reservoir for nutrients and water let oxygen reach the roots and provide gas exchange between the roots and the surrounding atmosphere. The medium composition affects the quality of the seedlings (Wilson et al., 2001). To improve seedling growth, media such as soil, cocopeat (Netam et al., 2020), vermicompost (Vandana 2019), FYM (Khot et al., 2022) and sand are employed.

To produce a suitable plant material by using dormancy breaking hormone and appropriate growing media for cultivating peach seedlings. This study aims to seedling growth attributes of high-chill peach, focusing on the effects of different growing media and growth regulators on promoting seedling growth.

2. MATERIALS AND METHODS

In 2022–2023, the experiment was carried out in the Department of Horticulture, College of Agriculture and Horticulture Research Centre, G.B.P.U.A & T, Pantnagar, Uttarakhand. Pantnagar is situated in the Himalayan Mountains at 29° North latitude and 79.3° East longitude. In summer, the temperature varies from 30 to 43°C, while in winter, it ranges from 0 to 9°C. Peach seeds gathered from KVK in Pithoragarh were employed as experimental material in these studies. In September 2022, when the peach was fully mature, the seeds were harvested and de-fleshed in preparation for stone extraction. The fruit stones were thoroughly cleaned with water after three to four days. The shade dried the pristine stones. The seeds were carefully removed by carefully breaking the stone (endocarp) from the ventral suture with a tiny hammer after they had dried for five to ten days in the shade. Along with use of media like Sand, soil, FYM, vermicompost and cocopeat were used as growing media in various proportions such as; Soil: sand: FYM (2:1:1), Soil: sand: cocopeat (2:1:1), Soil: sand: vermicompost (2:1:1) and Soil: cocopeat: vermicompost: FYM (2:1:1:1) were autoclaved under 15 psi pressure for 15 minutes followed by drenching in 0.2% Bavistin solution and filled in poly bags (Size 15 x 30 cm).

The following chemical solutions were made for research purposes in accordance with the protocol:

- i. Gibberellic acid (GA₃ 500 ppm): 500 mg of the chemical was dissolved in N/10 NaOH solution and the pH was adjusted to 7.0 using N/10 HCl. Distilled water was added to the mixture, increasing the final capacity to one liter.
- ii. Kinetin (50 ppm): 50 mg of the chemical was dissolved in distilled water and the volume was increased to one liter using distilled water.
- iii. Thiourea (5000 ppm): 5 g of the chemical was dissolved in distilled water and the amount was increased to 1 liter using distilled water.

In the experiment, focused on seedling growth studies, there were a total of 16 treatments involving two factors as well. The first factor was the growing medium, which included four combinations: soil: sand: farmyard manure (FYM), soil: sand: vermicompost, soil: sand: cocopeat and soil: FYM: vermicompost: cocopeat. The second factor was the chemical treatment, which remained consistent with the first experiment control, GA₃ at 500 ppm, kinetin at 50 ppm and thiourea at 5000 ppm.

List 1. The treatments combinations were as follows

Sl. No.	Symbol Used	Treatments Combination	Growing Media with Seed Treatment
1.	T ₁	M ₁ C ₁	Soil: Sand: FYM + Control
2.	T ₂	M ₁ C ₂	Soil: Sand: FYM + GA ₃ @ 500 ppm
3.	T ₃	M ₁ C ₃	Soil: Sand: FYM + Kinetin @ 50 ppm
4.	T ₄	M ₁ C ₄	Soil: Sand: FYM + Thiourea @ 5000 ppm
5.	T ₅	M ₂ C ₁	Soil: Sand: Vermicompost + Control
6.	T ₆	M ₂ C ₂	Soil: Sand: Vermicompost + GA ₃ @ 500 ppm
7.	T ₇	M ₂ C ₃	Soil: Sand: Vermicompost + Kinetin @ 50 ppm
8.	T ₈	M ₂ C ₄	Soil: Sand: Vermicompost + Thiourea @ 5000 ppm
9.	T ₉	M ₃ C ₁	Soil: Sand: Cocopeat + Control
10.	T ₁₀	M ₃ C ₂	Soil: Sand: Cocopeat + GA ₃ @ 500 ppm
11.	T ₁₁	M ₃ C ₃	Soil: Sand: Cocopeat + Kinetin @ 50 ppm
12.	T ₁₂	M ₃ C ₄	Soil: Sand: Cocopeat + Thiourea @ 5000 ppm
13.	T ₁₃	M ₄ C ₁	Soil: FYM: Vermicompost: Cocopeat + Control
14.	T ₁₄	M ₄ C ₂	Soil: FYM: Vermicompost: Cocopeat + GA ₃ @ 500 ppm
15.	T ₁₅	M ₄ C ₃	Soil: FYM: Vermicompost: Cocopeat + Kinetin @ 50 ppm
16.	T ₁₆	M ₄ C ₄	Soil: FYM: Vermicompost: Cocopeat + Thiourea @ 5000 ppm

3. RESULTS AND DISCUSSION

It is also evident from the data presented in Tables 1 and 2 and Graph 1 the effect of different growing media and seed treatment on seedling length and diameter was found to be significant. The maximum seedling length and diameter was reported in M₄ (81.81 cm and 7.6 mm), C₂ treated seeds was also help to enhance seedling length and diameter *i.e.*, 94.92 cm, 9.2 mm and combination effect M₄C₂ was reported maximum seedling length was 98.70 cm but the diameter was found to be non-significant. While the minimum seedling length and diameter were recorded under M₃ (73.13 cm and 7.0 mm), C₁ (56.03 cm and 4.1 mm) and M₃C₁ (49.18 cm), respectively. The combined use of cocopeat, vermicompost and FYM was more effective than individual because each have their own properties like; vermicompost improves physical properties and water holding capacity, while supplying adequate nutrients to the plants (Vandana, 2019). Similarly, FYM also contain organic matter, which binds soil particles and contain good source of soil nutrients, having a beneficial impact on plant growth (Khot et al., 2022) and cocopeat having good porosity, water holding capacity and aeration which increases the photosynthetic activity of functional leaves and thus improves the girth of the seedlings (Mishra et al., 2017 and Netam et al., 2020). Shrivastava et al. (2021) concluded in papaya that suitable growing media *i.e.*, soil: sand: FYM: vermicompost recorded the highest seedling length. Patel et al. (2019) also reported the significant increase in seedling length with the use of suitable growing media like soil: FYM: vermicompost. The results of our investigation from similar inspections were also reported by Haribhau et al. (2022) showing the positive impact of growing media, enhancing plant diameter rangpur lime seedlings. Similar perceptions are also proclaimed by gibberellic acid treated seeds shows higher plant height as compare to control (untreated seeds) might be due to GA₃ seems to activate metabolic processes or neutralize the impact of a growth inhibitor by causing cell multiplication and elongation in the cambium tissue of the internodal region (Hota et al., 2018). Singh (2020) observed that it activated the amylase and protease enzymes, which helped to convert stored starch into simple sugars and transfer them to the developing embryo, where they provided energy for seedling growth. These findings are also in agreement with Imani et al. (2011), who observed maximum seedling length

in peach with application of GA₃ 500 ppm. Similar, observations were also reported by Abou et al. (2021) in bitter almond, Negi et al. (2017) in walnut and Parvin et al. (2015) in black walnut. The above findings are also in line with Wani et al. (2014) and Rai et al. (2018) reported the positive effect of GA₃ @ 500 ppm on increasing the seedling diameter of apple and Khirni respectively.

It is evident from the data presented in Tables 3 and 4 and Graph 2 that there is significant effect of growing media and chemicals on number of leaves and leaf area. The maximum number of leaves and leaf area (198.92 and 6 cm² respectively,) were counted under the M₄ whereas the GA₃ treated seeds was also found maximum leaves and leaf area were 240.25 and 6.47 cm² and treatment combination the maximum leaves was found in M₄C₂ (298) while the leaf area was found non-significant under interaction. While the minimum number of leaves and leaf area were recorded under M₃ (140.83 and 5.32 cm²), C₁ (77.83 and 4.57 cm²) and M₃C₁ (67.67), respectively. The physical and chemical properties of soil is improved by application of organic manure which may supply adequate nutrients to the plants and encouraged the vigorous vegetative growth *i.e.*, more number of leaves per plant and also increasing the leaf area is better availability of moisture and nutrients to the plants thereby increased plant growth (Nainar et al. 2021). Our present finding are in line with the earlier finding of Vandana (2019) who studied that media containing soil: sand: FYM: vermicompost produces maximum number of leaves per plant in mango cv. Totapuri. They also found the positive influence of growth promoters on increasing the number of leaves might be due to GA₃, which enhance plant's physiological processes, thereby encouraging the rapid development of new leaves (Mishra et al., 2017). The increase in leaf area with GA₃ application may be because it promotes cell multiplication and cell elongation. These findings also support the result of Negi et al. (2017), Wani et al. (2014) and Sanaullah et al. (2020) were observed that GA₃ to be most effective in increasing number of leaves in walnut, apple and rough lemon.

The critical inspection of the data on shoot fresh and dry weight of high chill peach seedlings as influenced by growing media (Table 5 and 6) and the highest shoot fresh and dry weight (21.65 g and 12.49 g) in M₄, GA₃ treated seed found maximum (31.08 g and 18.16 g) and treatment combination M₄C₂ (36.90 g 22.07 g) while, the

least was recorded in M3 (15.19 g and 8.76 g), C1 (7.21 g and 3.61 g) and M3C1 (5.47 g and 2.90 g). They promote higher rate of water and nutrient mobilization leads to increased photosynthetic activity and translocation to various plant parts (Rameteke et al., 2015). Similarly, combination of vermicompost and FYM enhances the production of leaves and

chlorophyll content, which could improve the photosynthetic rate and favours more production of fresh and dry weight of plant (Patel et al., 2019). These finding are in agreement with the prior finding of Nainar et al. (2021) Netam et al. (2020) in acid lime and pomegranate cultivar Super Bhagwa, reported the maximum fresh weight and dry weight of shoots with

Table 1. Effect of growing media and seed treatment on seedling length (cm) of high chill peach at 150 days after seed germination

Sl. No	Growing Media	Chemical				Mean
		Control (C ₁)	GA ₃ (C ₂)	Kinetin (C ₃)	Thiourea (C ₄)	
1.	Soil: Sand: FYM (M ₁)	54.06	93.95	73.20	84.80	76.50
2.	Soil: Sand: Vermicompost (M ₂)	58.36	96.57	76.65	86.81	79.60
3.	Soil: Sand: Cocopeat (M ₃)	49.18	90.47	70.36	82.50	73.13
4.	Soil: FYM: Vermicompost: Cocopeat (M ₄)	62.53	98.70	78.59	87.43	81.81
	Mean	56.03	94.92	74.70	85.39	
		Media(M)	Chemical (C)	Interaction (MxC)		
	C.D. (5%)	1.39	1.39	2.78		
	S.Em±	0.48	0.48	0.96		

Table 2. Effect of growing media and seed treatment on diameter (mm) of high chill peach seedlings at 150 days after seed germination


Sl. No	Growing Media	Chemical				Mean
		Control (C ₁)	GA ₃ (C ₂)	Kinetin (C ₃)	Thiourea (C ₄)	
1.	Soil: Sand: FYM (M ₁)	4.0	9.0	7.5	8.2	7.2
2.	Soil: Sand: Vermicompost (M ₂)	4.2	9.3	7.6	8.3	7.4
3.	Soil: Sand: Cocopeat (M ₃)	3.9	8.9	7.4	8.0	7.0
4.	Soil: FYM: Vermicompost: Cocopeat (M ₄)	4.3	9.6	7.8	8.5	7.6
	Mean	4.1	9.2	7.6	8.3	
		Media(M)	Chemical (C)	Interaction (MxC)		
	C.D. (5%)	0.1	0.1	NS		
	S.Em±	0.14	0.14	0.07		

Table 3. Effect of growing media and seed treatment on number of leaves per plant of high chill peach seedlings at 150 days after seed germination


Sl. No	Growing Media	Chemical				Mean
		Control (C ₁)	GA ₃ (C ₂)	Kinetin (C ₃)	Thiourea (C ₄)	
1.	Soil: Sand: FYM (M ₁)	67.67	222.67	123.33	192.33	151.50
2.	Soil: Sand: Vermicompost (M ₂)	85.33	228.33	128.67	198.00	160.08
3.	Soil: Sand: Cocopeat (M ₃)	55.00	212.00	119.00	177.33	140.83
4.	Soil: FYM: Vermicompost: Cocopeat (M ₄)	103.33	298.00	142.00	252.33	198.92
	Mean	77.83	240.25	128.25	205.00	
		Media(M)	Chemical (C)	Interaction (MxC)		
	C.D. (5%)	1.85	1.85	3.70		
	S.Em±	0.90	0.90	1.81		

Table 4. Effect of growing media and seed treatment on leaf area (cm²) per leaf of high chill peach seedlings at 150 days after seed germination

Sl. No	Growing Media	Chemical				Mean
		Control (C ₁)	GA ₃ (C ₂)	Kinetin (C ₃)	Thiourea (C ₄)	
1.	Soil: Sand: FYM (M ₁)	4.39	6.45	5.50	6.00	5.58
2.	Soil: Sand: Vermicompost (M ₂)	4.75	6.50	5.65	6.29	5.79
3.	Soil: Sand: Cocopeat (M ₃)	4.03	6.23	5.27	5.73	5.32
4.	Soil: FYM: Vermicompost: Cocopeat (M ₄)	5.12	6.68	5.76	6.46	6.00
	Mean	4.57	6.47	5.54	6.11	
		Media(M)	Chemical (C)	Interaction (MxC)		
	C.D. (5%)	0.39	0.39	NS		
	S.Em±	0.13	0.13	0.27		



Graph 1. Effect of growing media and seed treatment on seedling length (cm) and seedling diameter (mm) of high chill peach



Graph 2. Effect of growing media and seed treatment on number of leaves and leaf area (cm²) of high chill peach seedlings

soil: sand: vermicompost. Anjanawe et al. (2013) also observed that soil: sand: FYM enhance shoot fresh weight and shoot dry weight in papaya. Similarly, Sanaullah et al. (2020) also found that GA₃ helps in enhancing absorption and redistribution of nutrients in plants which improves vegetative growth. These results also

elucidate the finding of Rameteke et al. (2015) and Rai et al. (2018) who proved that application of GA₃ increases shoot fresh weight and dry weight in papaya cv. Coorg Honey Dew and khirni, respectively. Similarly, Singh (2020) observed the maximum fresh weight and dry weight of shoot under GA₃ in aonla.

Table 5. Effect of growing media and seed treatment on shoot fresh weight (g) of high chill peach seedlings at 150 days after seed germination

Sl. No	Growing Media	Chemical				Mean
		Control (C ₁)	GA ₃ (C ₂)	Kinetin (C ₃)	Thiourea (C ₄)	
1.	Soil: Sand: FYM (M ₁)	7.10	29.87	12.05	22.10	17.78
2.	Soil: Sand: Vermicompost (M ₂)	7.50	32.20	13.20	24.68	19.39
3.	Soil: Sand: Cocopeat (M ₃)	5.47	25.33	9.95	20.04	15.19
4.	Soil: FYM: Vermicompost: Cocopeat (M ₄)	8.75	36.90	15.78	25.17	21.65
	Mean	7.21	31.08	12.74	23.00	
		Media(M)	Chemical (C)	Interaction (MxC)		
	C.D. (5%)	1.24	1.24	2.48		
	S.Em±	0.43	0.43	0.86		

Table 6. Effect of growing media and seed treatment on shoot dry weight (g) of high chill peach seedlings at 150 days after seed germination

Sl. No	Growing Media	Chemical				Mean
		Control (C ₁)	GA ₃ (C ₂)	Kinetin (C ₃)	Thiourea (C ₄)	
1.	Soil: Sand: FYM (M ₁)	3.23	16.81	6.32	12.87	9.81
2.	Soil: Sand: Vermicompost (M ₂)	3.51	18.28	7.78	13.49	10.76
3.	Soil: Sand: Cocopeat (M ₃)	2.90	15.48	5.00	11.64	8.76
4.	Soil: FYM: Vermicompost: Cocopeat (M ₄)	4.82	22.07	8.10	14.98	12.49
	Mean	3.61	16.01	7.77	11.17	
		Media(M)	Chemical (C)	Interaction (MxC)		
	C.D. (5%)	0.72	0.72	1.44		
	S.Em±	0.25	0.25	0.50		

4. CONCLUSION

Thus, the finding in this experiment *i.e.*, seedling growth studies carried with the high chill peach, the seed treatment with GA₃ @ 500 ppm along with the growing media soil: FYM: vermicompost: cocopeat (2:1:1:1) recorded the maximum seedling length, diameter, number of leaves, leaf area, shoot fresh weight and shoot dry weight which can be recommended for raising the rootstocks for the production of quality planting material of peach on seedling rootstock.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Abad, M., Noguera, P., Puchades, R., Maquieira, A., & Noguera, V. (2002). Physico-chemical and chemical properties of some coconut dusts for use as a peat substitute for containerized ornamental plants. *Bioresource Technology*, 82, 241-245.
- Abou Rayya, M. S., Nabila, E. K., Malaka, A. S., & Thanaash, M. (2021). Effect of cold stratification and GA₃ on desheled seeds germination and seedlings growth of bitter almond. *Middle East Journal of Agriculture Research*, 10(4), 1173-1181.
- Agbo, C. U., & Omaliko, C. M. (2006). Initiation and growth of shoots of *Gongronema latifolia* Benth stem cuttings in different rooting media. *African Journal of Biotechnology*, 5(5), 425-428.
- Anjanawe, S. R., Kanpure, R. N., Kachouli, B. K., & Mandloi, D. S. (2013). Effect of plant growth regulators and growth media on seed germination and growth vigour of

- papaya. *Annals of Plant and Soil Research*, 15(1), 31-34.
- Cetinbas, M., & Koyuncu, F. (2006). Improving germination of *Prunus avium* L. seeds by gibberellic acid, potassium nitrate, and thiourea. *Horticultural Science*, 33(3), 119-123.
- Chang, Y. S., & Sung, F. H. (2000). Effects of gibberellic acid and dormancy-breaking chemicals on flower development of *Rhododendron pulchrum* Sweet and *Rhododendron scabrum* Don. *Scientia Horticulturae*, 83, 331-337.
- Diaz, D. H., & Martin, G. C. (1972). Peach seed dormancy in relation to endogenous inhibitors and applied growth substances. *Journal of the American Society for Horticultural Science*, 97, 651-654.
- El-Badawy, H. E. M., & El-Aal, M. M. M. A. (2013). Physiological response of Keitt mango (*Mangifera indica* L.) to kinetin and tryptophan. *Research Journal of Applied Sciences*, 9(8), 4617-4626.
- El-Keblawy, A. (2013). Impacts of dormancy-regulating chemicals on innate and salinity-induced dormancy of four forage grasses native to Arabian deserts. *Grass and Forage Science*, 68, 288-296.
- Haribhau, K. A., Paikra, M. S., Deshmukh, U. B., Ramteke, L. K., Nishad, D., Taram, M., & Rathore, A. (2022). Effect of GA₃ and growing media on seed germination, growth, and vigour of Rangpur lime (*Citrus limonia* Osbeck). *Pharma Innovation*, 11(12), 3056-3061.
- Hota, S. N., Karna, A. K., Jain, P. K., & Dakhad, B. (2018). Effect of gibberellic acid on germination, growth, and survival of jamun (*Syzygium cumini* L. Skeels). *Journal of Pharmaceutical Innovation*, 7(8), 323-326.
- Imani, A., Rasouli, M., Tavakoli, R., Zarifi, E., Fatahi, R., Barba-Espin, G., & Martinez-Gomez, P. (2011). Optimization of seed germination in *Prunus* species combining hydrogen peroxide or gibberellic acid pre-treatment with stratification. *Seed Science and Technology*, 39(1), 204-207.
- Khot, A. A., Jadhav, R. A., Khot, F. A., & Pathan, S. F. (2022). Effect of different potting media on the growth of Bullock's heart (*Annona reticulata* L.). *International Journal of Phytology Research*, 2(2), 14-17.
- Lipe, W. N., & Crane, J. C. (1966). Dormancy regulation in peach seeds. *Science*, 153, 541-542.
- Mishra, U., Bahadur, V., Prasad, V. M., Verty, P., Singh, A. K., Mishra, S., Swaroop, N. (2017). Influence of GA₃ and growing media on growth and seedling establishment of papaya (*Carica papaya* L.) cv. Pusa Nanha. *International Journal of Current Microbiology and Applied Sciences*, 6(11), 415-422.
- Nainar, P., Muthulakashmi, S., & Manivannan, L. M. (2021). Effect of growing media on seed germination and seedling growth of acid lime (*Citrus aurantifolia* Swingle). *International Journal of Current Microbiology and Applied Sciences*, 10(8), 110-122.
- Negi, P., Nautiyal, B., Thakur, N., Negi, M., & Kumari, V. M. (2017). Effect of different pre-sowing treatments on seed germination and seedling growth of walnut (*Juglans regia* L.). *International Journal of Current Microbiology and Applied Sciences*, 6(7), 3844-3849.
- Netam, S. R., Sahu, G. D., Markam, P. S., & Minz, A. P. (2020). Effect of different growing media on rooting and survival percentage of pomegranate (*Punica granatum* L.) cuttings cv. Super Bhagwa under Chhattisgarh plains condition. *International Journal of Chemical Studies*, 8, 1517-1519.
- Pallavi, S. P., Masuthi, D. A., Sabarad, A. I., Naik, N. H., Gollagi, S. G., & Nataraj, K. H. (2022). Influence of pre-germination treatments on germination, growth, and vigour of passion fruit (*Passiflora edulis* var. *flavicarpa*) seeds. *Journal of Pharmaceutical Innovation*, 11(2), 479-482.
- Parvin, P., Khezri, M., Tavasolian, I., & Hosseini, H. (2015). The effect of gibberellic acid and chilling stratification on seed germination of eastern black walnut (*Juglans nigra* L.). *Journal of Nuts*, 6(1), 67-76.
- Patel, M. V., Parmar, B. R., Halpati, A. P., Parmar, A. B., & Pandey, A. K. (2019). Effect of growing media and foliar spray of organics on seedling growth and vigour of acid lime. *International Journal of Chemical Studies*, 7(1), 1-4.
- Rai, R., Samir, M., Srivastava, R., & Uniyal, S. (2018). Improving seed germination and seedling traits by pre-sowing treatments in khirni (*Manilkara hexandra*). *Bulletin of Environmental Pharmacology and Life Sciences*, 7(4), 77-81.
- Ramteke, V., Paithankar, D. H., Kamatyanatti, M., Baghel, M. M., Chauhan, J., & Khichi,

- P. (2015). Seed germination and seedling growth of papaya as influenced by GA₃ and potting media. *Journal of Progress in Agriculture*, 6(1), 129-133.
- Sanaulah, A., Hazarika, B. N., Wangchu, L., & Sarma, P. (2020). Effect of plant growth regulators and chemicals on seedling growth of rough lemon (*Citrus jambhiri* L.) under hydroponic conditions. *International Journal of Current Microbiology and Applied Sciences*, 9(9), 2353- 2358.
- Shrivastava, P., Prasad, V. M., Panigrahi, H. K., Bahdur, V., & Singh, Y. K. (2021). Effect of different types of media and containers on germination, survival, growth, and establishment of papaya (*Carica papaya* L.) cv. Red Lady under protected conditions. *Biological Forum*, 13(2), 670-675.
- Singh, P. (2020). Effect of potassium nitrate (KNO₃) and gibberellic acid (GA₃) on root and shoot growth of aonla (*Emblica officinalis* L.). *Journal of Pharmacognosy and Phytochemistry*, 9(1), 584-586.
- Thakur, B., & Singh, H. (2015). Studies on seed germination in peach (*Prunus persica* L. Batsch) rootstock 'Flordaguard'. *Bioscan*, 10(2), 651-654.
- Vandana, S., Murthy, P. V., & Goudappanavar, B. (2019). Effect of organic mixtures on stone germination and seedling growth of mango (*Mangifera indica* L.) cv. Totapuri under net house and polyhouse conditions. *International Journal of Current Microbiology and Applied Sciences*, 9(4), 1643-1655.
- Wani, R. A., Malik, T. H., Malik, A. R., Baba, J. A., & Dar, N. A. (2014). Studies on apple seed germination and survival of seedlings as affected by gibberellic acid under cold arid conditions. *International Journal of Science and Technology Research*, 3(3), 2010-2016.
- Wilson, S. B., Stoffella, P. J., & Graetz, D. A. (2001). Use of compost as a media amendment for containerized production of two subtropical perennials. *Journal of Environmental Horticulture*, 19(1), 37-42.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/126364>