

Gorlin-Goltz Syndrome: Report of 4 Cases

Ouassime Kerdoud^{1*} and Faical Slimani^{1,2}

¹*Department of Stomatology, Oral and Maxillofacial Surgery, Hospital of August 20th, University Hospital, Casablanca, Morocco.*

²*Faculty of Medicine and Pharmacy, Hassan 2 University of Casablanca, Casablanca, Morocco.*

Authors' contributions

This work was carried out in collaboration between both authors. Author OK wrote the first draft of the manuscript, managed the analyses of the study and the literature searches. Author FS revised the scientific content of the paper. Both authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Armando Montesinos Flores, National Autonomous University of Mexico, Mexico.

Reviewers:

(1) Yousif Osman Yousif Ali, University of Khartoum, Sudan.

(2) Archana Buch, Dr. D. Y. Patil Vidyapeeth, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/61980>

Case Study

Received 24 August 2020

Accepted 29 October 2020

Published 07 November 2020

ABSTRACT

Gorlin-Goltz syndrome, also known as basal cell nevomatosis (CBN), is a rare inherited disorder belonging to the family of neurocristopathies or diseases caused by abnormalities of the neural ridges. We report the case of 4 patients, suffering from this syndrome, followed up in the stomatology and maxillofacial surgery service in Casablanca, including a familial form in two cases: a mother and her son and 2 isolated cases, followed and treated in our service of maxillofacial surgery of the August 20 hospital in Casablanca. A clinical, biological and radiological assessment was made followed by surgical treatment for the 3 patients, an anatomo-pathological examination and the establishment of a strict monitoring plan.

Keywords: Gorlin-Goltz syndrome; cell nevomatosis; stomatology; maxillofacial surgery.

1. INTRODUCTION

Gorlin-Goltz syndrome, also known as basal cell nevomatosis (CBN), is a rare

inherited disorder belonging to the family of neurocristopathies or diseases caused by abnormalities of the neural ridges.

*Corresponding author: Email: wassim.kdd@gmail.com;

The mode of transmission is autosomal dominant with full penetrance and variable expressivity.

The diagnosis of disease is clinico-radiological, which can be confirmed by genetic study. The risk of malignant transformation of the skin lesions requires ad vitam monitoring of patients.

2. CASE PRESENTATION

We report the case of 4 patients, suffering from this syndrome, followed up in the stomatology and maxillofacial surgery service in Casablanca, including a familial form in two cases: a mother and her son and 2 isolated cases, followed and treated in our service of maxillofacial surgery of the August 20 hospital in Casablanca.

A clinical, biological and radiological assessment was made followed by surgical treatment for the 3 patients, an anatomic-pathological examination and the establishment of a strict monitoring plan.

3. OBSERVATIONS

We report the case of a 60-year-old woman and her 39-year-old son as well as a 55-year-old man, and another 36-year-old. 3 of the patients presented with multiple maxillo-mandibular cysts.

Observation 1: A 55-year-old man, operated on for mandibular cyst in October 2016, presented for swelling next to the ascending branch of the left mandible, and the left jug, with 3 suspicious lesions in the nasal wing, the root of the left nose and the head of the left eyebrow, and facial and cervical nevi.



Fig. 1. Cervical naevomatosis




Fig. 2. CT of the face showing maxillomandibular cysts

Conservative surgical treatment was performed in the patient consisting of enucleation of the cysts. Anatomical pathological examination concluded with odontogenic keratocysts.

Observation 2: A 60-year-old woman with a history of: High blood pressure under hypertensive treatment, excision of multiple maxillo-mandibular cysts in 2007, in whom clinical examination shows: a budding lesion with raised edges at the level of the lower right eyelid with extension to the free edge of the lower eyelid, a lesion in the upper left eyelid, frontal bump, café-au-lait spots on the trunk and upper limbs.

A biopsy of the right palpebral lesion, the pathological examination was in favor of a nodular basal cell carcinoma and focal adenoid ulcerated. After the patient benefited from an excision of the lesions: (right internal canthale, left cheekbone, upper left eyelid, right eyelid-conjunctival), and a 2nd stage reconstruction by an advancement flap, and a graft of the oral mucosa for the conjunctival plane.

Observation 3: A 39-year-old man (the son), having as an antecedent: 1st degree consanguinity with the above-mentioned patient, on clinical examination presents: a large frontal bump, macrocephaly, diffuse pigmentary lesions at the level of the face and hypertelorism.



Fig. 3. Nodular lesion of the right lower eyelid and the left upper eyelid



Fig. 4. Picture after eyelid reconstruction (result after 1 month)




Fig. 5. Facial nevi and protruding frontal bumps




Fig. 6. Mandibular cystic images

An excision of the mandibular cyst was performed, the pathological examination was in favor of an odontogenic keratocyst

Observation 4: A 36-year-old man, operated at the age of seventeen for a mandibular cyst (no documents), on clinical examination presented: two swellings in the mandibular level; angular right and at the level of the left horizontal branch, an enlarged frontal bump, and two pigmented lesions at the level of the two lower eyelids

The diagnosis of Gorlin Goltz syndrome was made in all four patients based on various clinical and radiological criteria.

The monitoring plan is made up of regular clinical and radiological monitoring.

4. DISCUSSION

Gorlin and Goltz syndrome is an autosomal dominant genetic disorder. The first description was made in 1894 by Jarisch but it was Gorlin and Goltz who in 1960 defined the affection. And since then, more than 700 cases have been reported in the literature [1].

It is a rare syndrome, its prevalence is estimated at 1 / 57,000 in the general population, males and females are affected in the same proportions, and the majority of cases have been reported in the Caucasian race [2].

This condition is linked to a mutation in the PTCH tumor suppressor gene having chromosomal location 9q22.3-q31 [3].



Fig. 7. Right cheek swelling



Fig. 8. Tri-focal cystic image

Clinically, this condition is characterized by a spectrum of developmental abnormalities and a predisposition to different cancers [4].

The first manifestations seem to be maxillary cysts, during the first decade. In a third of cases, the cutaneous signs start at puberty [4].

The clinical manifestations of GORLIN-GOLTZ syndrome are very diverse, we note mainly:

4.1 Maxillofacial Signs

4.1.1 Odontogenic keratocysts

Approximately 5% of keratocysts are associated with this syndrome. They are found in 80% of

patients with Gorlin syndrome while they are present in only 5 to 7% of the general population, frequently discovered incidentally during a radiological assessment, for orthodontic purposes [5].

4.1.2 Dental abnormalities

Malocclusions, impacted teeth, ectopia or heterotopia, and dental agenesis [2].

4.2 Cutaneous and Mucous Signs

4.2.1 Skin basal cell nevi

Usually appear between puberty and the 3rd decade. They appear as lenticular papules a few millimeters in diameter, painless, polymorphic,

single or multiple (up to several thousand), these lesions preferentially sit on the cervico-facial area as well as on the thorax [6].

The nevi can degenerate into typical basal cell epitheliomas, favored by solar exposure especially to UVB, these transformations are evidenced by an increase in size of the ridges which become crusty and ulcerated. Evolution remains local [7].

4.2.2 Palm plantar wells

The wells found on the palms of the hands and the soles of the feet appear to be pathognomonic and are one of the major criteria for the disease [6].

4.2.3 Epidermoid cysts

Large (1-2 cm) and often multiple epidermoid cysts resembling odontogenic keratocysts occur on the limbs and trunk [8].

4.3 Skeletal Manifestations

4.3.1 Calcifications of the brain envelopes

Calcifications of the scythe of the brain, the interclinoid ligaments or the tent of the cerebellum [6].

4.3.2 Anomalies in the shape of the skull

These anomalies, although belonging to the minor criteria of the disease, are of great diagnostic interest since they are visible from birth.

The skull is unusually large with prominent frontal bumps, the eyebrow arches are prominent, hypertelorism is common [6].

4.3.3 Costal abnormalities

Costal abnormalities are present in approximately 42% of patients with CBN, and are mainly represented by costal bifidities, agenesis, and synostosis [2].

4.3.4 Vertebral abnormalities

Abnormalities of the cervical or thoracic vertebrae are useful diagnostic signs, being found in about 60% of affected people. The C6,

C7, T1 and T2 vertebrae are most frequently involved [5].

4.4 Gynecological Signs

4.4.1 Ovarian fibroids

These fibroids are present in 25% of patients with Gorlin syndrome and are often bilateral [9,10].

4.5 Neurological and Psychiatric Signs

- Mental retardation
- Agenesis of the corpus callosum,
- Medulloblastoma

5. DIAGNOSTIC

The complexity of the clinical signs found in this syndrome has led to the establishment of specific criteria to facilitate the diagnosis. It involves finding in a patient at least two of the four major criteria or one major criterion and two minor criteria.

5.1 Management of Patients with Basal Cell Nevomatosis

Therapeutically, there is currently no specific treatment. Their management is symptomatic with special oncology monitoring, involving the intervention of various specialists [6].

Patients with the syndrome need information about the syndrome. The results of several epidemiological studies have shown that the risks of CBC have a strong correlation with exposure to UV radiation. Thus, these patients are able to avoid excessive exposure to the sun. It is important that they wear 100% UV blocking sunglasses since the skin around the eyes (similar to the nose / ears) is vulnerable to CBCs [7].

Sunscreens with a high protection factor (SPF 30 +) should be applied before going out and reapplied every 2 to 3 hours, and more frequently when swimming or sweating [8].

Genetic investigation is essential for the detection of new cases in the family. A neurological exam is required annually in children up to the age of seven to detect a medulloblastoma [3].

Table 1. Diagnostic criteria for CBN

Major criteria:
<ul style="list-style-type: none"> • Multiple basal cell carcinomas (> 2), or basal cell carcinoma before the age of 20 years • Histologically proven odontogenic cysts of the jaw • Palmoplantar pits ≥ 3 • Bifid, merged or particularly flared dimensions • Affected by a first degree relative)
Minor criteria:
<ul style="list-style-type: none"> • Macrocephaly • Orofacial congenital malformations: cleft lip or palate, frontal lump • Hypertelorism • Radiological abnormalities: closed saddle turcica, vertebral anomalies: hemi-vertebrae, fusion or lengthening of the vertebral bodies, bone defects of the hands or feet, small flame-shaped bony gaps in the hands and feet • Ovarian fibroma • Medulloblastoma

6. CONCLUSION

Gorlin and Goltz syndrome is a genetic disorder, rare, multisystem, with autosomal dominant transmission. It is classically defined by the triad composed of basal cell nevi, maxillary keratocysts and skeletal malformations. This syndrome is due to mutations in a tumor suppressor gene having 9q22.3-q31 as chromosomal location.

The carcinological potential of this syndrome is serious, justifying early detection and regular and prolonged monitoring of patients and their descendants. The treatment is simply symptomatic.

CONSENT

As per international standard or university standard, participant's written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ghailan MR, Benhalima H, Rabeh G, et al. Syndrome de Gorlin-Goltz: A propos d'un cas Med. buccale chir. Buccale. 2007; 13(2):97-101.

2. Manfredi M, et al. Nevoid basal cell carcinoma syndrome: A review of the literature. *Int. J. Oral. Maxillofac. Surg.* 2004;33:117-124.
3. Dereure O. Gorlin syndrome and SUFU germline mutations: *Annales de dermatologie et de vénéréologie.* 2015; 142:231-232.
4. Mouhsine, et al. Syndrome de nævomatose baso-cellulaire: A propos de 2 cas, feuillets de radiologie. 2009;49(2): 111-115.
5. Salmon B, et al. Odontogenic keratocyst related to a Gorlin syndrome: A case report *Archives de Pédiatrie.* 2008;15: 406-409.
6. Lo Muzio I. Gorlin syndrome. *Encyclopédie Orphanet*, Janvier; 2002.
7. Lomuzziol. Nevoid basal cell carcinoma syndrome (Gorlin syndrome). *Orphanet Journal of Rare Diseases.* 2008;3(32).
8. Kimonis VE, et al. Clinical manifestations in 105 persons with nevoid basal cell carcinoma syndrome. *Am. J. Med. Genet.* 1997;69: 229-308.
9. Omrani H, et al. Recurrent ovarian fibromas in condition of gorlin syndrome; *Journal de Gynécologie Obstétrique et Biologie de la Reproduction.* 2010;39:584-587.

10. Kadlub N, et al. Pediatric keratocystic odontogenic tumor and nevoid basal cell carcinoma syndrome. Predictive factors for recurrence and aggressiveness: *Rev Stomatol Chir Maxillofac.* 2012;113:148-154.

© 2020 Kerdoud and Slimani; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/61980>