



Antifungal Activity of Copper, Zinc and Potassium Compounds on Mycelial Growth and Conidial Germination of *Fusarium solani* f. sp. *piperis*

**Renata Aparecida Ahnert dos Santos¹, Verônica D'Addazio^{1*},
João Vitor Garcia Silva¹, Antelmo Ralph Falqueto¹, Marcelo Barreto da Silva¹,
Edilson Romais Schimldt¹ and Adriano Alves Fernandes¹**

¹*Departamento de Ciências Agrárias e Biológicas, Universidade Federal do Espírito Santo (UFES),
Centro Universitário Norte do Espírito Santo, Rodovia Governador Mario Covas, Km 60, CEP 29932-
540, São Mateus, Espírito Santo, Brazil.*

Authors' contributions

All authors contributed equally to this work. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/MRJI/2019/v29i630179

Editor(s):

(1) Dr. Ana Cláudia Coelho, Department of Veterinary Sciences, University of Trás-os-Montes and Alto Douro, Portugal.

Reviewers:

(1) Seweta Srivastava, Lovely Professional University, India.

(2) Ferreira, Luiz Leonardo, Universidade Estadual da Paraíba, Brazil.

(3) Douira Allal, University Ibn Tofail, Kenitra, Morocco.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/52926>

Original Research Article

Received 03 September 2019

Accepted 10 December 2019

Published 16 December 2019

ABSTRACT

Fusariosis is a disease that causes economic damage to black pepper (*Piper nigrum* L.) producers. Despite being a major disease, there is no record of efficient chemical control. Thus, the objective was to evaluate the antifungal activity of copper, zinc and potassium compounds in mycelial growth and conidial germination of *Fusarium solani* f. sp. *piperis* *in vitro*. For inoculation in PDA (Potato Dextrose Agar) medium, 7 mm discs from the pure culture were transferred to Petri dishes. The plates were incubated at 25°C in a biochemical oxygen demand (BOD) chamber, with photoperiod of 12 h, for 15 days. Micronutrients were supplied as sulfates, CuSO₄ (copper sulfate) and ZnSO₄ (zinc sulfate), at concentrations of 1, 5, 10, 15 and 20 mmol/L. Potassium macronutrient (K) was supplied as KCl (potassium chloride) at concentrations of 30, 60, 90, 120 and 150 mmol/L. The experiment was performed using a completely randomized design with 6 treatments and ten replications. CuSO₄ showed fungicidal effect at concentrations of 10, 15 and 20 mmol/L. For ZnSO₄ mycelial growth was completely inhibited at concentrations of 15 and 20 mmol/L. There was no

*Corresponding author: E-mail: veronicadaddazio@yahoo.com;

inhibition or reduction of fungal growth in the presence of K. Copper and zinc at minimal concentrations were efficient in controlling mycelial growth and inhibition of spore germination of *F. solani* f. sp. *piperis*. In contrast, potassium did not exert fungicidal or fungistatic effect on the fungus.

Keywords: *Fusarium solani*; antigungal activity; fungal growth inhibition; nutrients.

1. INTRODUCTION

Pathogenic fungi include a large and heterogeneous group of organisms that occupy an important position in both agriculture and natural populations [1]. The genus *Fusarium* spp. is classified as Ascomycete, characterized by a hyaline mycelium, branched and septate, with phyllide-shaped sporophores and conidia of varying size and shape. It has spindle-shaped macroconidia with several septa [2]. *Fusarium* species are known as phytopathogens, saprophytes with worldwide distribution [3].

Fusarium solani (Mart.) Appel & Wr. emend. Snyd. & Hans. f. sp. *piperis*, Albuquerque (Teleomorph *Nectria haematococca* Berk. & Br. f. sp. *piperis* Albuq.) is the causal agent of fusariosis and has brought great economic damage to black pepper (*Piper nigrum* L.) producers, decimating whole crops [4,5]. The disease may start in the root system or in the shoot [6,7]. When initiated by the roots, the root system is reduced and becomes necrotic, causing yellowish and flaccid leaves and premature fall. On the other hand, once started in the aerial part, it is characterized by the presence of yellowish branches in a very vigorous plant [6]. With the evolution of the disease is observed drying in the upper and lower part of the plant [6-8]. According to Pandey et al. [9], production losses due to plant diseases are a considerable challenge for the current agricultural production system worldwide, representing at least 25% of the total. Although fusariosis is a major disease, there is no record of efficient chemical control.

The methods used in the control of pathogenic isolates of *Fusarium* include the use of resistant varieties and soil disinfection with the chemical fungicide and crop rotation using non-host plants [10]. The use of resistant cultivars would be an alternative, but difficulties such as the identification of resistance genes or the pathogen's ability to adapt to new genotypes may make resistance a temporary solution [11]. In addition, the chemical control of *Fusarium* spp. is not fully efficient since the pathogen penetrates the vascular tissue of the plant [12].

Crop rotation would be of little efficiency since this pathogen is a soil fungus, capable of surviving for long periods in crop debris and presents several plant species as hosts [11]. Conventional synthetic fungicides are largely considered the most effective and economical means for treating the disease. However, the intensity of use and specific mode of action of most synthetic fungicides ultimately lead to resistance problems and an increased environmental cost [9,13,14]. Thus, alternative forms of control are of increasing interest, leading to the investigation and development of effective and sustainable products for the control of plant pathogens [15]. Zambolim et al. [16] reported that some micro and macronutrients have been identified as one of the main mineral elements associated with the induction of disease resistance in plants.

Due to their low cost, protective activity and reduced risk of resistance development controlled by the broad mode of action against pathogens, Cu compounds have been exploited to protect crops from many pests, including those that cause numerous bacterial and fungal infections [17]. According to Zambolim et al. [18], Cu ions in contact with spores or the pathogen's germ tube may accumulate in the membrane or penetrate and concentrate inside of the spores or mycelium, where they act by inhibiting enzymes essential to the metabolic process of microorganisms. Once accumulated in the cells, their effects become irreversible.

Interest in research on zinc (Zn) derivatives is increasing [19,20], due to its strong antimicrobial activity at low concentrations and its non-toxic characteristics in adequate quantities. Zn as an essential micronutrient plays an important role in many integral metabolic processes [21]. It can also help increase chlorophyll and carotenoid biosynthesis and improve plant photosynthetic apparatus [22]. Significant optoelectric, physical and antimicrobial properties of Zn offer great potential for increasing agricultural productivity [23]. Its mode of action is not completely understood, but it is known to act directly on the pathogen [18].

Among macronutrients of great importance to the plant, potassium (K) is one of the elements that has very positive results in reducing the incidence of pests and diseases [24,25], being able to reduce the severity of more of 100 fungi [26]. Taiz and Zeiger [27] reported that K is an essential plant nutrient required as a cofactor for over 40 enzymes, many of which are involved in respiration and photosynthesis. As such, it is an important nutrient in plant disease prevention as it is involved in many cellular processes that influence disease severity. Its effect on the prevention of diseases caused by bacteria, fungi and nematodes has been reported [18,28]. Increased resistance to disease from K fertilization has been attributed to several mechanisms, such as cell permeability and decreased susceptibility of tissues to pathogen maceration and penetration [18,29]. K influences the reduction of plant diseases due to the activation of enzymes involved in respiration and photosynthesis, carbon chain supply processes for defense substance synthesis, as well as stomatal regulation influencing mass flow solute transport [30].

The use of nutrients with antifungal action may be a strategy for controlling pathogens that cause invaluable economic losses. Nutrients such as Cu, Zn and K are easily accessible, inexpensive and still contribute to plant nutrition. Thus, the objective of this study was to evaluate the antifungal activity of Cu, Zn and K compounds against *Fusarium solani* f. sp. *piperis* growth.

2. MATERIALS AND METHODS

2.1 Microorganism and Cultivation

The isolate of *Fusarium solani* f. sp. *piperis* CML 2466, from the Coleção Micológica de Lavras, Federal University of Lavras - MG was used. The fungus was maintained on Petri dishes containing PDA (Potato Dextrose Agar) at 4°C. For inoculation, 7 mm pure culture discs were transferred to Petri dishes containing the same medium. The plates were incubated at 25°C in BOD (Biochemical Oxygen Demand), photoperiod of 12 h for 15 days.

2.2 Copper, Zinc and Potassium Concentrations

Micronutrients were supplied in PDA medium as copper sulfate (CuSO₄) and zinc sulfate (ZnSO₄)

at concentrations of 1, 5, 10, 15 and 20 mmol/L. Potassium macronutrient (K) was supplied as potassium chloride (KCl) at concentrations of 30, 60, 90, 120 and 150 mmol/L. The nutrients were diluted in sterile distilled water and at the time of plating were added to the PDA culture medium in laminar flow hood. After solidification, a 7 mm diameter fungal mycelium disc with 15-day-old was transferred to the center of each Petri dish (68 mm diameter). The PDA medium with the fungus disc was used as control. The plates were sealed with Parafilm and maintained in BOD at 25°C with 12 h photoperiod.

2.3 Mycelial Growth

The evaluation of *F. solani* mycelial growth in the control plates and treatments was determined every 2 days by measuring the diameter of the colonies in orthogonal directions with the digital pachymeter until the control treatment colony reached edge of the plate, 12 days after inoculation (DAI). The growth inhibition percentage was calculated according to Guo et al. [31], where the antifungal index (%) = $(1 - Da / Db) \times 100$, where: Da gives the diameter of the growth zone in the test plate and Db the diameter of the growth zone in the control plate.

2.4 Spore Count

The spore suspension was prepared by adding 20 mL of sterile distilled water to each plate containing the fungus, which was scraped with a Drigalsky handle for efficient spore extraction. The spore count was performed in a Neubauer Chamber and the suspension was adjusted to a concentration of 10^6 spores/mL⁻¹.

2.5 Statistical Analysis

The experiment was performed using a completely randomized design with 6 treatments and 10 replications for each treatment (Cu, Zn and K). Each repetition consisted of a petri dish. Mycelial growth data were analyzed by linear regression. The values obtained in spore production and colony growth were submitted to analysis of variance by the F test and the means compared by the Tukey test at 5% of probability using Genes software [32].

3. RESULTS

3.1 Mycelial Growth

The mycelial growth of *F. solani* was dependent on the nutrient and the dose used. Fungal growth

was completely inhibited in some treatments. The CuSO_4 showed fungicidal effect at concentrations of 10, 15 and 20 mmol/L, completely inhibiting growth of the colonies (Fig. 1a). However, at a concentration of 5 mmol/L, a fungistatic effect was observed since mycelial growth was initiated 6 DAI (Fig. 2a), despite the projection of the line. For ZnSO_4 mycelial growth was completely inhibited at concentrations of 15 and 20 mmol/L, showing fungicidal effect and

significantly reduced ($P \leq 0.05$) at concentrations of 5 and 10 mmol/L, exerting fungistatic effect (Fig. 1b). This result is ratified after eight days of incubation by observing mycelial growth at a concentration of 10 mmol/L (Fig. 2b). There was a significant difference for KCl treatment ($P \leq 0.05$) between the tested concentrations. However, there was no inhibition or reduction of fungal growth in the presence of this nutrient (Figs. 1c and 2c).





Fig. 1. Antifungal activities of CuSO_4 (a), ZnSO_4 (b) and KCl (c) against *F. solani* f. sp. *piperis* on PDA at different concentrations

Data are shown as average values. Columns followed by the identical letter are not statistically different according to by Tukey test, at 5% probability ($P \leq 0.05$). Bars represent the standard error of the mean



**Fig. 2. Effect of CuSO₄ (a), ZnSO₄ (b) and KCl (c) on the mycelial growth of *F. solani* f. sp. *piperis*, 12 days after inoculation. Control: only PDA medium
*Signifivative at 5% probabily by F test**

3.2 Percent Growth Inhibition (P.I.).

The mycelial growth inhibition index confirmed the efficiency of the antifungal activity of CuSO₄

and ZnSO₄ (Table 1). For CuSO₄, at a concentration of 5 mmol/L, there was inhibition greater than 50% 12 DAI. The other concentrations inhibited 100% fungal growth.

Similar results were observed for ZnSO₄ (Table 1). However, for KCl, in none of the evaluated concentrations was observed P.I. below 50%. At 2 DAI there was growth induction (Table 1) with no fungistatic or fungicidal effect for this nutrient.

3.3 Conidia Number

Twelve days after inoculation (12 DAI), the conidia number of *F. solani* was inhibited in the presence of CuSO₄, ZnSO₄ and KCl. The Cu micronutrient reduced by 84% the conidial germination at 1mmol/L concentration in relation to the control. The same was not observed for Zn at the same concentration (Fig. 3a, Table 2). In the other Cu and Zn concentrations, conidial germination was significantly inhibited ($P \leq 0.05$),

with values greater than 80%. For K treatment, there was a 20.6% reduction in the number of conidia at 30 mmol/L. The other concentrations presented a reduction greater than 50% when compared to the control (Fig. 3b, Table 2).

4. DISCUSSION

The widespread and persistent nature of *Fusarium* spp. may be due to its ability to maintain and multiply in a wide variety of complex carbohydrates and proteins, thus resisting adverse climates and high levels of toxic substances such as many antibiotics and fungicides. *F. solani* seems to incorporate some of the most difficult members of the genus [33].




Fig. 3. Conidia production of *F. solani* f. sp. *piperis* on CuSO₄, ZnSO₄ (a) and KCl (b) in different concentrations

Data are shown as average values. Columns followed by the identical letter are not statistically different according to by Tukey test, at 5% probability ($P \leq 0.05$). Bars represent the standard error of the mean

Table 1. Percent growth inhibition (P.I.) of *F. solani* f. sp. *piperis* in CuSO₄, ZnSO₄ and KCl

Time (days)	P.I. (%)					
	2	4	6	8	10	12
CuSO₄ mmol/L						
Control	0	0	0	0	0	0
1	12.0	19.4	29.4	33.0	34.5	37.2
5	100	100	77.8	70.7	66.4	61.0
10	100	100	100	100	100	100
15	100	100	100	100	100	100
20	100	100	100	100	100	100
ZnSO₄ mmol/L						
Control	0	0	0	0	0	0
1	14.4	18.8	22.0	20.3	17.0	19.8
5	65.7	70.4	73.6	71.5	68.9	67.7
10	100	100	89.9	81.5	79.5	79.9
15	100	100	100	100	100	100
20	100	100	100	100	100	100
KCl mmol/L						
Control	0	0	0	0	0	0
30	-0.9	7.1	16.6	18.7	19.2	21.4
60	-3.8	8.0	16.6	18.8	16.2	16.5
90	2.8	9.8	15.8	17.4	14.0	13.3
120	-3.0	2.7	8.2	6.1	4.4	7.0
150	-1.4	3.7	4.8	3.8	5.0	8.3

Table 2. Percent growth inhibition (P.I.) of *F. solani* f. sp. *piperis* in CuSO₄, ZnSO₄ and KCl

Reduction in conidia number (%)	
mmol/L	CuSO ₄
1	84.2
5	97.2
10	100.0
15	100.0
20	100.0
ZnSO ₄	
1	0.0
5	82.2
10	98.4
15	100.0
20	100.0
KCl	
30	20.6
60	52.9
90	48.3
120	53.6
150	73.5

Inhibition of mycelial growth of *F. solani* f. sp. *piperis* *in vitro* revealed significant variations in fungus sensitivity to different nutrients tested. In the present study, CuSO₄, ZnSO₄ were more effective in inhibiting fungal growth, while KCl showed relatively weak effect comparing it to other nutrients. However, it was observed that

Cu was the most effective compound against mycelial growth and *F. solani* conidia germination, showing significant inhibition at relatively low concentrations of the compound.

According to Resende et al. [34] and Melo et al. [35], Cu, Zn and K compounds are used for the control of microorganisms. They can have a direct effect on the pathogen (fungicidal or fungistatic effect) or activate the natural defense of plants, resulting in induced resistance [36,37].

The Cu is currently used due to its antifungal properties. In particular, Cu is responsible for interference with homeostatic processes and cell membrane functions, protein synthesis damage, reactive oxygen species production, and DNA disruption [38,39]. Civardi et al. [40] observed that Cu exerted toxic effect on *Rhodonía placenta* fungal cell by breakdown of different basic metabolic processes. Significant antifungal activity of Cu has been revealed in a number of pathogenic species including *Fusarium* sp., *Aspergillus niger*, *Rhizoctonia solani*, *Alternaria solani*, *Alternaria alternata* and *Phoma destructiva* [9,41,42].

Regarding Zn, several studies have shown its antibacterial activity [43-46]. However, there are few studies reporting the suggested mechanism

for antifungal activity of Zn compounds [47,48]. Some authors suggest that such a mechanism may be based on the formation of reactive oxygen species that disrupt the integrity of the cell membrane, preventing pathogen growth [49,50,46,48]. According to He et al. [48], Król et al. [51], and Ashajyothi et al. [52], Zn compounds showed fungistatic potential against *Fusarium* sp., *Botrytis cinerea*, *Penicillium expansum*, *Aspergillus niger* and *Rhizopus stolonifer*. Chand et al. [53] observed that among the micronutrients tested Zn presented the greatest inhibition of mycelial growth of *Fusarium oxysporum* f. sp. *cuban*.

The marked toxic effect of Cu and Zn against fungal spores compared to mycelial growth can be attributed to the structural differences between the spore wall and the fungal vegetative phase. Bartnicki-Garcia [54] observed that the chitin content of many fungal species is significantly higher in the hyphae wall compared to the spore wall, making the latter more susceptible to some compounds. In addition, during the spore germination process, the presence of enzymes such as disulfide reductase and glucanases result in weakening of the cell wall, facilitating germ tube lengthening and thus creating sites of greater sensitivity to toxic substances in contact with the cell fungal [54]. In general, conidial germination may reflect reproductive capacity and fungal development. Savi et al. [20] suggest that the effect of Zn compounds on fungal growth may be related to their property, altering reproductive capacity in terms of conidia viability. Malandrakis et al. [55], studying the effect of copper and zinc on various microorganisms, found that Cu was effective against *Alternaria alternata*, *Botrytis cinerea*, *Monilia fructicola*, *Verticillium dahliae*, *Colletotrichum gloeosporioides*, *Fusarium oxysporum* f. sp. *radicis lycopersici* and *Fusarium solani* while Zn exerted a fungicidal effect against *M. fructicola*, *F. solani* and *V. dahliae*.

Although K has not exerted inhibition or reduction of *F. solani* mycelial growth and spore production, there are reports in the literature that K acts as an inducer of resistance to plant diseases [56-58]. The use of K as a plant fertilizer may decrease the incidence of fungal and bacterial as well as insect diseases [59], mainly due to changes in primary metabolism and plant hormonal responses [29,57]. Dordas [60] observed that the application of KCl on foliage can prevent the attack of mildew on wheat.

5. CONCLUSION

The results obtained in this work provide evidence that copper and zinc exhibited antifungal activity against *F. solani* f. sp. *piperis* under laboratory conditions. High antifungal activity was observed at low concentrations of copper (10 mmol/L) and zinc (15 mmol/L), favoring the use of these compounds to control mycelial growth and conidia production. On the other hand, no potassium concentration was active against the fungus. Based on these results, field experiments can be performed to verify the performance of these nutrients as *Fusarium* growth inhibitors, as well as the physiological and nutritional behavior of the plant.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Burdon JJ, Silk J. Sources and patterns of diversity in plant-pathogenic fungi. *Phytopathology*. 1997;87:664-669.
2. Leal PC, Cantanhede KL, Silva LM, Bezerra GFB. *Micotoxinas do Fusarium e seu potencial carcinogênico*. 70^a Ed. São Paulo: NewsLab Press; 2005. Brazil.
3. Hennequin C, Abachin E, Symoens F, Lavarde V. Identification of *Fusarium* species involved in human infections by 28S rRNA gene sequencing. *J Clin Microbiol*. 1999;37:3586-3589.
4. Duarte MLR, Albuquerque FC. *Sistema de Produção da Pimenta-do-reino*, Embrapa Amazônia Oriental; 2005. [Accessed 24 July 2017] Available:<http://sistemasdeproducao.cnptia.embrapa.br/FontesHTML/Pimenta/PimenteiradoReino/paginas/importancia.htm>
5. D'Addazio V. *Crescimento micelial de Fusarium solani* f. sp. *piperis* e respostas de cultivares de pimenta-do-reino (*Piper nigrum* L.) ao estresse abiótico e biótico: *Biometria, fotossíntese, resistência e avaliação de produtos alternativos de controle da fusariose*. Vitória: Universidade Federal do Espírito Santo; 2017. Brazil.
6. Ventura JA, Costa H. *Manejo da fusariose da pimenta-do-reino no estado do Espírito Santo*. Vitória: Instituto Capixaba de Pesquisa, Assistência Técnica e Extensão Rural; 2004. Brazil.

7. Silva SO, Neto APD, Silva MB. Pimenta-do-reino: importância da defesa fitossanitária para a sustentabilidade da atividade na região norte do Espírito Santo. Rev Bras Agropecu Sustent. 2011;1:88-92. Brazil.
8. Tremacoldi CR. Principais doenças fúngicas da pimenteira-do-reino no estado do Pará e recomendações de controle. Belém: Embrapa Amazônia Oriental; 2010. Brazil.
9. Pandey S, Giri K, Kumar R, Mishra G. Nanopesticides: Opportunities in crop protection and associated environmental risks. Proc Natl Acad Sci Sect B Biol Sci. 2018;88:1287–1308.
10. Agrios GN. Plant pathology. 5th Ed. New York: Academic Press; 2005.
11. Sutton JC. Strategies for biological control of necrotrophic pathogens in perennial crops. Fitopatol Bras. 2000;25:235-238.
12. Tokeshi H. Doenças e pragas agrícolas geradas e multiplicadas pelos agrotóxicos. Fitopatol Bras. 2000;25:264-271. Brazil.
13. Kah M, Kookana RS, Gogos A, Bucheli TD. A critical evaluation of nanopesticides and nanofertilizers against their conventional analogues. Nat Nanotechnol. 2018;13:677–684.
14. Sun Q, Li J, Le, T. Zinc oxide nanoparticle as a novel class of antifungal agents: current advances and future perspectives. J Agric Food Chem. 2018;66:11209–11220.
15. Deliopoulos T, Kettlewell PS, Hare MC. Fungal disease suppression by inorganic salts: A review. Crop Protec. 2010;29: 1059-1075.
16. Zambolim L, Rodrigues FA, Capucho AS. Resistência a doenças de plantas induzida pela nutrição mineral. In: Controle alternativo de pragas e doenças. Venzon M, Paula Junior TJ, Pallini A, Editors. Viçosa: Epamig/CTZM. 2005;275-318.
17. Keller AA, Adeleye AS, Conway JR. Comparative environmental fate and toxicity of copper nanomaterials. Nanolmpact. 2017;7:28–40.
18. Zambolim L, Ventura AJ, Zañón LA. Efeito da nutrição mineral no controle de doenças de plantas. Viçosa: Editora Independente; 2012. Brazil.
19. Savi GD, Vitorino V, Bortoluzzi AJ, Scussel VM. Effect of zinc compounds on *Fusarium verticillioides* growth, hyphae alterations, conidia, and fumonisin production. J Sci Food Agric. 2013;93:3395–3402.
20. Savi GD, Vitorino V, Bortoluzzi AJ, Scussel VM. Antifungal properties of zinc-compounds against toxigenic fungi and mycotoxin. Int J Food Sci Technol. 2013;48:1834–1840.
21. Rout GR, Das P. Effect of metal toxicity on plant growth and metabolism. Agronomie. 2003;23:3–11.
22. Aravind P, Prasad MNV. Zinc protects chloroplasts and associated photochemical functions in cadmium exposed *Ceratophyllum demersum* L., a fresh water macrophyte. Plant Sci. 2004;166:1321–1327.
23. Hussain I, Singh NB, Singh A, Singh H. Green synthesis of nanoparticles and its potential application. Biotechnol Lett. 2016;38:545–560.
24. Graham DR. Effects of nutrients stress on susceptibility of plants to disease with particular reference to the trace elements. Adv Bot Res. 1983;10:221–276.
25. Sharma S, Duveiller E, Basnet R, Karki CB. Effect of potash fertilization on *Helminthosporium* leaf blight severity in wheat, and associated increases in grain yield and kernel weight. Field Crop Res. 2005;93:142–150.
26. Raij, van B. Potássio: necessidade e uso na agricultura moderna. Piracicaba: Potafós;1990. Brazil.
27. Taiz L, Zeiger E. Fisiologia e Desenvolvimento Vegetal. 6^a Ed. Porto Alegre: Artmed; 2017. Brazil.
28. Rawat N, Pumphrey MO, Li S, Zhang X. Wheat Fhb1 encodes a chimeric lectin with agglutinin domains and a pore-forming toxin-like domain conferring resistance to *Fusarium* head blight. Nat Genet. 2016;48: 1576–1580.
29. Wang M, Zheng Q, Shen Q, Guo S. The critical role of potassium in plant stress response. Int J Mol Sci. 2013;14:7370–7390.
30. Bloom AJ. Nutrição mineral. In: Fisiologia Vegetal. Taiz L, Zeiger E, Editors. Porto Alegre: Artmed. 2004;96-103.
31. Gu Z, Chen R, Xing R, Liu S. Novel derivatives of chitosan and their antifungal activities *in vitro*. Carbohydr Res. 2006;341:351–354.
32. Cruz CD. Genes Software – extended and integrated with the R, Matlab and Selegen. Acta Sci. 2016;38:547-552.
33. Smith SN. An overview of ecological and habitat aspects in the genus *Fusarium* with

- special emphasis on the soil-borne pathogenic forms. *Plant Pathol Bull.* 2007;16:97–120.
34. Resende MLV. Indução de resistência na cafeicultura: Perspectivas de uso. In: Manejo fitossanitário da cultura do cafeeiro. Blun, LEB, Editor. Lavras: Universidade Federal de Lavras. 2008;25-35. Brazil.
 35. Melo LGDL, Silva EKC, Campos Neto JRM, Lins SRO. Indutores de resistência abióticos no controle da fusariose do abacaxi. *Pesq Agropec Bras.* 2016;51: 1703-1709. Brazil.
 36. Spolti P, Valdebenito-Sanhueza RM, Campos AD, Del Ponte EM. Modo de ação de fosfitos de potássio no controle da podridão olho de boi em maçã. *Summa Phytopathol.* 2015;41:42-48. Brazil.
 37. Costa BHG, Resende MLV, Ribeiro Júnior PM, Mathioni SM. Suppression of rust and brown eye spot diseases on coffee by phosphites and by-products of coffee and citrus industries. *J Phytopathol.* 2014;162: 1–8.
 38. Krumova EZ, Pashova SB, Dolashka-Angelova PA, Stefanova T. Biomarkers of oxidative stress in the fungal strain *Humicola lutea* under copper exposure. *Process Biochem.* 2009;44:288–295.
 39. Rai M, Ingle PA, Pandit R, Paralikar P. Copper and copper nanoparticles: role in management of insect-pests and pathogenic microbes. *Nanotechnol Rev.* 2018;1-14.
 40. Civardi C, Schwarze FWMR, Wick P. Micronized copper wood preservatives: An efficiency and potential health risk assessment for copper-based nanoparticles. *Env Poll.* 2015;200:126–132.
 41. Nemati A, Shadpour S, Khalafbeygi H, Ashraf S. Efficiency of hydrothermal synthesis of nano/microsized copper and study on *in vitro* antifungal activity. *Mater Manuf Process.* 2015;30:63–69.
 42. Aleksandrowicz-Trzcinska M, Szaniawski A, Olchowik J, Drozdowski S. Effects of copper and silver nanoparticles on growth of selected species of pathogenic and wood-decay fungi *in vitro*. *For Chron.* 2018;94:109–116.
 43. Yamamoto O. Influence of particle size on the antibacterial activity of zinc oxide. *Int J Inorg Mater.* 2001;3:643–646.
 44. Stoimenov PK, Klingler RL, Marchin GL, Klabunde JS. Metal oxide nanoparticles as bactericidal agents. *Langmuir.* 2001;18: 6679–6686.
 45. Zhang L, Jiang Y, Ding Y, Povey M. Investigation into the antibacterial behaviour of suspensions of ZnO nanoparticles (ZnO nanofluids). *J Nanopart Res.* 2007;9:479–489.
 46. Liu Y, He L, Mustapha A, Li H. Lin. Antibacterial activities of zinc oxide nanoparticles against *Escherichia coli* O157:H7Y. *J Appl Microbiol.* 2009;107: 1193–1201.
 47. Sawai J, Yoshikawa T. Quantitative evaluation of antifungal activity of metallic oxide powders (MgO, CaO and ZnO) by an indirect conductimetric assay. *J Appl Microbiol.* 2004;96:803-9.
 48. He L, Liu Y, Mustapha A, Lin M. Antifungal activity of zinc oxide nanoparticles against *Botrytis cinerea* and *Penicillium expansum*. *Microbiol Res.* 2011;166:207–215.
 49. Feng QL, Wu J, Chen GQ, Cui FZ. A mechanistic study of the antibacterial effect of silver ions on *Escherichia coli* and *Staphylococcus aureus*. *J Biomed Mat Res.* 2000;52:662–668.
 50. Applerot G, Lipovsky A, Dror R. Enhanced antibacterial activity of nanocrystalline ZnO due to increased ROS mediated cell injury. *Adv Funct Mater.* 2009;19:842–852.
 51. Król A, Pomastowski P, Rafińska K, Railean-Plugaru V. Zinc oxide nanoparticles: synthesis, antiseptic activity and toxicity mechanism. *Adv Colloid Interf Sci.* 2017;249:37–52.
 52. Ashajyothi C, Prabhurajeshwar C, Handral HK, Kelmani CR. Investigation of antifungal and anti-mycelium activities using biogenic nanoparticles: An eco-friendly approach. *Environ Nanotechnol Monit Manag.* 2016;5:81–87.
 53. Chand G, Jaiswal US, Maru AK. Effect of micronutrients on Panama wilt of banana (*Fusarium oxysporum* f. sp. *cubense*) and its synergistic action with *Trichoderma viride*. *Conference Proceedings of International Conference on Innovative Approaches in Applied Sciences and Technologies (iCiAsT-2016)*; 2016.
 54. Bartnicki-Garcia S. Cell wall chemistry, morphogenesis, and taxonomy of fungi. *Annu Rev Microbiol.* 1968;22:87–108.
 55. Malandrakis AA, Kavroulakis N, Chrysikopoulos CV. Use of copper, silver and zinc nanoparticles against foliar and soil-borne plant pathogens. *Sci Total Environ.* 2019;670:292–299.

56. Holzmueller EJ, Jose S, Jenkins MA. Influence of calcium, potassium, and magnesium on *Cornus florida* L. density and resistance to dogwood anthracnose. *Plant Soil*. 2007;290:189–199.
57. Zorb C, Senbayram M, Peiter E. Potassium in agriculture—status and perspectives. *J Plant Physiol*. 2014;171: 656–669.
58. Gao X, Zhang S, Zhao X, Wu O. Potassium-induced plant resistance against soybean cyst nematode via root exudation of phenolic acids and plant pathogen-related genes. *PLoS ONE*. 2018;13:1-13.
59. Amtmann A, Troufflard S, Armengaud P. The effect of potassium nutrition on pest and disease resistance in plants. *Physiol Plantarum*. 2008;133:682–691.
60. Dordas C. Role of nutrients in controlling plant diseases in sustainable agriculture. A review. *Agro Sustain Dev*. 2008;28:33–46.

© 2019 Santos et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/52926>