



The Variation in Anaemic Parameters in the Ethnic Groups of Pakistan

Ayesha Nageen^{1*}, Alina Arshad² and Jamal Ara¹

¹Department of Medicine, UMDC/Creek General Hospital, Karachi, Pakistan.

²Lyceum School, Karachi, Pakistan.

Authors' contributions

This work was carried out in collaboration between all authors. Author AN designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Author AA managed the statistical analyses of the study. Author JA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IBRR/2018/42073

Editor(s):

(1) Dharmesh Chandra Sharma, Incharge Blood Component & Aphaeresis Unit, G. R. Medical College, Gwalior, India.

Reviewers:

(1) Flavia Miranda Bandeira, Universidade do Estado do Rio de Janeiro, Brazil.

(2) Iragbogie Al-Mustapha Imoudu, Federal Medical Centre Azare, Nigeria.

Complete Peer review History: <http://www.sciencedomain.org/review-history/24877>

Received 22nd March 2018

Accepted 27th May 2018

Published 30th May 2018

Original Research Article

ABSTRACT

Aims: To observe any diversity of pattern in the anaemic parameters in the major ethnic groups of Pakistan.

Study Design: Observational, cross-sectional.

Place and Duration of Study: Department of Medicine, Creek general Hospital, Korangi, Karachi, Pakistan from October 2017 to March 2018.

Methodology: 1200 adults above 18 years were included. Pregnant females, age less than 18 years, any severe organ insufficiency, stroke, neurodegenerative disease, known blood dyscrasias or terminal illness at baseline evaluation were excluded. Their ethnic background was grouped into Urdu speaking, Punjabi, Balochi Sindhi, Pashtun, Bengali, and Gilgit according to their place of origin, mother tongue or parents' place of origin. The complete blood picture was performed to identify anaemia, grade the severity, classify the red cell indices and exclude any blood dyscrasias in each of the ethnic group.

Results: Out of 1200 adults, 626(52%) were anaemic. In decreasing order, the most anaemic were Bengalis (57.6%), Sindhis (55%), Urdu speaking (54%), Pashtuns (54%), Balochis (52%), Gilgits

*Corresponding author: E-mail: ayeshanageen@gmail.com;

(49%) and Punjabis (45%). In males, Sindhis (60%), Pashtuns (57%), Bengalis (56%), and Urdu speaking (51%) were most anaemic. In females, Balochis (71%), Bengalis (64%), Urdu speaking (55%) were most anaemic. Anaemia was prevalent in the younger age group of the Urdu speaking, Pashtun and Sindhis community, while in the rest of the ethnic groups anaemia was most common in the middle age group. In males, anaemia was highest in the young age group of Urdu Speaking, Sindhis, and Pashtuns while in females in all the ethnic groups the middle age group was most anaemic and had least anaemia in elderly females except in Bengalis where the least was in young females. Mild anaemia was most common overall and in both genders. Regarding red cell indices, Punjabis and Bengalis were normocytic while in the rest the most were microcytic. In the males, Urdu speakers, Punjabis and Bengalis were significantly more normocytic anaemic, while Sindhis were more microcytic. In females, in all the ethnic groups' microcytic anaemia was prevalent. Anaemia was three times more in the Sindhi group of smokers compared to the Sindhi nonsmokers while in the rest of the ethnic groups there was no specific correlation.

Conclusion: This survey specifies that in Pakistan the key measures of health status are albeit dependent on the financial status, literacy rate, sanitation, access to primary care facilities, effective role of health providers but the genetic composition and cultural norms also play their part.

Keywords: Anaemia; ethnicity; hemoglobin; red cell indices.

1. INTRODUCTION

Anaemia persists to be a primary universal health concern with a global prevalence of 33% as stated in 2010 [1]. The World Health Organization (WHO) estimated approximately 1.62 billion people of the world to be anaemic [2]. Pakistan is foremost in this concern as more than half of its population is anaemic leading to a deterioration of mental and physical growth, subsequently affecting its economic progress. Its identification, correction and prevention being an urgent priority should be the key focus in the domain of health of this country [3].

Anaemia is a multidimensional disease with both nutritional and non-nutritional etiologies being responsible such as iron, folate or cobalamin deficiencies, chronic or acute blood loss, inherited genetic defects, chronic diseases and/or inflammatory disorder, parasitic infestations, etc [4].

Genomics is linked to public health science as population genetics and the epidemiology of a disease is found to be associated under a certain pattern which then affects the clinical practice of that population [5,6] this health inequality between ethnic groups are seen due to a number of reasons like reluctance in accessing healthcare, language and cultural barriers, certain myths and religious views, gender discrimination, poverty or underprivileged sectors [7]. Understanding the health physiognomies of each group can enable us to prioritize screening programs, education and resource allocation according to their need.

Pakistan has several ethnic communities personalized by their culture and language according to their place of origin [8]. This Karachi based study is to observe any difference in the pattern of anaemia and its extended parameters with respect to the ethnic groups in adults. It is to probe whether any genetic inborn structure or any cultural norms of a certain ethnic group would stand out as a specific culprit leading to anaemia. An insight into these health inequalities with respect to ethnicity is needed for effective healthcare to be delivered appropriately.

2. MATERIALS AND METHODS

The study was a cross sectional, observational one conducted in the outpatient clinic of a tertiary care hospital, Creek General Hospital, Korangi, Karachi from October 2017 to March 2018. The hospital is located in a multi ethnic area where strata from different ethnic backgrounds present to the clinics. It was done on 1200 adults (age 18 years and above) selected by convenient sampling. Pregnant females, age less than 18 years and any severe organ insufficiency, stroke, neurodegenerative disease, known blood dyscrasias or terminal illness at baseline evaluation were excluded. The ethnic background was noted and grouped into Urdu speaking, Punjabi, Balochi Sindhi, Pashtun, Bengali, and Gilgit according to place of origin, mother tongue or parents' place of origin. Other relevant information was obtained. The complete blood picture (CBC) was performed by ABX Micros 60 Hematology Analyzer to identify anaemia, grade the severity, classify the red cell indices by Mean Corpuscular Volume (MCV) and

exclude any blood dyscrasias or acute hemolytic conditions. The data was analyzed on SPSS version 22 with the universal and dependent variables, quantitative and descriptive, analyzed in each group. Mean hemoglobin with standard deviation and range was tabulated for each ethnic group. Linear regression was done for hemoglobin and parity.

2.1 Objective Definitions

Anaemia (WHO Criteria) [9]

Hemoglobin levels; Males: <13 g/dl Females: <12 g/dl

Severity of anaemia

	Mild	Moderate	Severe
Females	10-11.9 g/dl	8 – 9.9 g/dl	<8.0 g/dl
Males	10-12.9 g/dl	8 – 9.9 g/dl	<8.0 g/dl

MCV Classification of anaemia: Normocytic; 84-96fl Microcytic ;< 84fl Macrocytic >96fl

Age groups

Young; 18-30 years Middle age group;31-50 years Elderly group; >50 years

3. RESULTS

Among the 1200 adults that presented 631(52.6%) were Urdu Speaking, 250(20.8%) were Punjabis, 92(7.7%) were Sindhis, 80(6.7%) were Pashtuns, 71(6%) were Gilgits, 52(4.3%) were Bengalis, and 24(2%) were Balochis. Out of the 626(52%) were anaemic. The frequency of anaemia in each ethnic group and the gender distribution of anaemia in each group are demonstrated in Table 1. The mean hemoglobin (with std deviation and range) of each ethnic group is shown in Table 2. The percentage of anaemia in the ethnic groups according to age groups is displayed in Fig. 1. The severity of anaemia in the various ethnic groups is presented in Fig. 2. The frequency of anaemia with respect to age groups in males is displayed in Fig. 3. In females, the anaemia percentage was highest in the middle age group in all the ethnic groups and least in the elderly group of all ethnicities except the Bengalis where the young females had least anaemia ($P= 0.12$). Mild anaemia was more common in male gender of all groups ($P=.743$) and in the female gender ($P=.090$) when severity was checked separately in each gender. The red cell indices in the different ethnic groups are tabularized in Table 3.

When analyzed gender wise; in the males - Urdu speakers, Punjabis and Bengalis were significantly more normocytic anaemic, while Sindhis were more microcytic. The Balochis, Pashtuns, and Gilgits were equally more normocytic and microcytic than macrocytic. ($P=.884$) in females, in all the ethnic groups' microcytic anaemia was more prevalent. There is no specific correlation or pattern of variation of anaemia with the number of children in the females in any of the ethnic groups. Anaemia was three times more in the Sindhi group of smokers compared to the Sindhi nonsmokers while in the rest of the ethnic groups there was no specific correlation on linear regression. There was no specific ethnic group in which diabetics had significantly more anaemia compared to non-diabetics of that group.

4. DISCUSSION

Ethnicity influences the parameters of a disease by and large factors such as differences in service availability, geographical origin, communication and language obstacles, cultural attitudes/beliefs, socio-economic status, and genetic differences in disease prevalence. We analyzed the anaemic parameters of each ethnic group residing in Karachi and then compared the statistics with the adults of that group living at their place of origin. The severity of anaemia was by large mild in all groups and genders giving a hope that the risk factors if effectively attended to would minimize the prevalence of anaemia in Pakistan.

Bengalis had the highest frequency of anaemia in this cohort and the equal prevalence in both genders points to certain common factors that are contributing to anaemia in the whole population. The Bengalis of Karachi are mainly involved in the fishing industry living near the coastal areas in unhygienic and poor conditions [10]. Their diet mainly consists of rice, lentils and fish which lack iron and other micronutrients. This lack of red meat in their diet was also stressed upon by Faruk in his Bangladeshi study [11]. Males had mostly normocytic anaemia probably because of the mixed nutritional deficiencies (of iron and cobalamin) and prevalence of chronic diseases. The females are microcytic understandably due to their hormonal cycles, multiple pregnancies and malnutrition with inadequate iron intake compared to the loss which is why the middle age groups were worst affected. Bharti [12] studied anaemia in West Bengal and noted that it was prevalent in all

socioeconomic group of females and Merrill insisted that females are subject to gender biasness in diet availability [13]. Factors, present in Bengalis of Karachi, common to both genders are a parasitic infestation, daily diet lacking folate and iron, unhygienic conditions, poverty and food contamination by sea pollutants which was seen more in the middle and elderly age group.

A conspicuous result in Sindhis which was an eye opener was the higher frequency of anaemia in males compared to females with both the genders being microcytic. A probable conclusion is that dietary iron deficiency, hookworm manifestation, acid peptic disease and smoking are the main anaemic culprits here (after we

exclude blood dyscrasias). The addiction profile was very high in Sindhi males and anaemia being three times more in smokers than nonsmokers' glares at the consequences of gutka and tobacco and explains why the young males are more anaemic than other age groups of Sindhis. The growing use of gutka and other forms of tobacco results in severe malnutrition as these noxious products suppress appetite and prevent absorption of micronutrients in the body. This low hemoglobin is present despite the secondary physiological erythropoietin response caused by tobacco. The statistics of Sindhi women health has also worsened. The results of the NNS 2011 for anaemia showed that the prevalence of

Table 1. Anaemia in different ethnic groups and with respect to gender in that ethnic group (P=0.282)

Ethnicity group	Total presented	The no. of Anaemic/ % anaemic in that group	The no. of males of that group	The no. and % of anaemic males in that group	The no. of females of that group	The no. and % of anaemic females in that group
Bengali	52	30(57.6%)	27	14(56%)	25	16(64%)
Sindhi	92	51(55%)	40	24(60%)	52	27(52%)
Urdu speaking	631	341(54%)	252	130(51%)	379	211(55%)
Pashtun	80	43(54 %)	35	19(54%)	45	24(53%)
Balochis	24	12(52%)	10	2(22%)	14	10(71%)
Gilgits	71	35(49%)	26	12(46%)	45	23(51%)
Punjabis	250	114(45%)	104	37(35%)	146	77(53%)

Table 2. Mean hemoglobin found in all the groups

Ethnicity	Mean	Std. Deviation	Range
Urdu Speaking	11.93	3.84	89.00
Punjabi	12.59	6.47	104.50
Sindhi	11.62	2.09	10.30
Bengali	12.04	2.16	11.00
Gilgit	12.02	1.87	9.20
Balouchi	11.86	2.45	8.80
Pashtun	12.22	2.42	14.60
Total	12.07	4.21	107.00

Table 3. The frequency of red cell indices in ethnic groups (P= .197)

Ethnicity	MCV group			Total anaemic in that group
	Normocytic (% of total anaemic of that group)	microcytic<84(% of total anaemic of that group)	macrocytic>96(% of total anaemic of that group)	
Urdu Speaking	140(41)	182(53)	19(5)	341
Punjabi	55(48)	53(46)	6(5)	114
Sindhi	17(33)	31(60)	3(5)	51
Bengali	16(53)	13(43)	1(3)	30
Gilgit	14(46)	17(48.5)	4(11)	35
Balouchi	2(16)	10(83)	0	12
Pushto	12(27)	29(67)	2(4)	43
Total	256	335	35	626




Fig. 1. The anaemic distribution in the ethnic groups according to age group




Fig. 2. The severity of anaemia in different ethnic groups ($P= .856$)

anaemia in non-pregnant females had risen from 28% in 2001 to 50% in 2011 [14,15]. Cultural myths, lack of approach to healthcare, ineffective counseling by health care providers, multiple

pregnancies and addiction to 'beeri'(raw tobacco) is causing a serious setback in decreasing the rising ratio of women with malnutrition which takes its toll in the middle age group females.




Fig. 3. The pattern of anaemia frequency according to age groups in the males of the different ethnic groups (P= .164)

Laghari [16] studied anaemia in students of Sindh University, Jamshoro and his studies showed females to be more anaemic compared to males which is unlike our ratio of the Sindhis residing in Karachi.

The Urdu speaking community is primarily an urban population and their exposure and tendency towards Fad diets and unhealthy junk food which then causes acid peptic disease as a sequel is comparatively more compared to other communities. Their high percentage of addiction to gutka and betel nut has led to high prevalence of anaemia in young males. Bharati [12] has already emphasized that there is a high anaemic frequency in India with factors more or less common to both sides of the country borders. The females suffer more due to the gender discrimination, unequal food sharing and child bearing leading to the middle age group being most anaemic and the elderly females having attained menopause are the least anaemic.

The Pashtuns of Karachi although are physically strong and involved in strenuous physical occupational work yet more than half of them were found to be anaemic. One of the prevailing offenders of the high anaemic prevalence in

young males is the rampant Niswar addiction. There is also suspicion of high frequency of constipation leading to hemorrhoids in Pashtuns (but studies are needed to confirm this) which could be the cause of microcytic anaemia in them. Shahab's study [17] conducted in Peshawar agreed to our results which shows that there are common correctable propensities in both genders of the community.

The Balochis that presented here were mostly those living in the coastal areas and are fisherman by profession. They catch trade and eat fish. This then leads to another important factor of sea pollution which is the toxic effect of the minerals consumed by the fish which eventually is transferred to the fish consumers. This sea pollution is taking a heavy toll on human health. There is also a high addiction rate of 'tiryak' (finer opioid). The women were triple times more anaemic than men and when we compared this to the Balochis living in Balochistan the statistics were similar for women. In Balochistan, poverty and unemployment coupled with worsening law and order situation have deepened woes and severely affected the nutritional status of women [18]. However, a common factor in most Balochi women, whether

urban or rural, was the issue of multiple pregnancies, low adult literacy rate and conservative restricted living [19]. According to National Nutrition Survey (NNS) conducted in 2011, the anaemic statistics of Balochistan jumped from 26.9 to 49 per cent in 2011 with higher prevalence of anaemia in females and the knowledge of micronutrients being least in Balochis [14].

Half of the Gilgits of Karachi were anaemic which is a higher rate compared to the natives living in the Gilgit Baltistan. One obvious reason of the natives to be less anaemic is the high altitude in the Northern region of Baltistan which causes secondary polycythemia masking anaemia if it is internally present. Karachi is a low lying area which would alienate the erythropoietin effect. Uzma [20] stressed in her study on anaemia in Gilgit that the anaemia frequency is lower because of higher literacy rate in the region due to a well-established system of basic education. Inference can be drawn from this fact that educational status of women has a major impact in solving this problem. Khalil [21] showed that there is a high prevalence of intestinal infestations especially *Ascaris Lumbricoides* among Gilgits which can then be a factor causing anaemia even if they migrated to Karachi. The highest percentage of macrocytic anaemia was in Gilgit group showing cobalamin/folic acid deficiency to be a contributing factor which is assured by Tanweer in his Karachi based study on macrocytic anaemia [22]. This needs to be explored in further studies to rule out any genetic intrinsic factor anomalies.

Punjabis residing in Karachi had the lowest percentage of anaemia in this study cohort showing them to be nutritionally balanced. Females were considerably more anaemic, and a microcytic one, due to their physiological processes along with being in a male privileged society. According to the NNS 2011 report the knowledge of micronutrients was the highest in Punjab area [14] which again emphasizes the deep impact education has in the development of healthcare. Middle age group of males are the working class and the revenue generator and they being normocytic anaemic means that we need to reinforce awareness of diet and health checkups in them.

5. LIMITATIONS OF STUDY

It was a hospital-based study and a larger scale research should be done in the community.

There could have been an overestimation of the frequency of anaemia since it was a hospital sampling. There was also an inability to know for certain whether patients had underlying occult hemoglobinopathy or X-linked G6PDH disorder which is known causes for anaemia here. Some the ethnic group had a low sample size. Further studies are warranted in each ethnic group to investigate each of the variables separately.

6. CONCLUSION

This survey suggests that in Pakistan the key measures of health status are albeit dependent on the financial status, literacy rate, sanitation, access to primary care facilities, effective role of health providers but the genetic composition and cultural norms also play their part. It is important to try to understand where and how ethnic differences impacts on healthcare delivery if health inequalities are to be reduced across the whole population. Evidence suggests that the poorer socio-economic position of some ethnic groups is the main driver and aggravator of ethnic health inequalities.

Pakistan needs a more comprehensive strategy to prevent losing 'working years' from anaemia by focusing on practically implemented, home based interventions that include diet counselling, micronutrient supplementation, food fortification to attend to nutritional anaemia while concomitantly responding to other etiologies of anaemia. Hygiene and sanitation, deworming. Malaria prevention is other areas of necessary interposition. Health policies are required to tackle health inequalities keeping the ethnic variations in mind. Callen's review [23] on healthcare facility of Punjab points to a necessity for improvement in the role and availability of healthcare officers at Basic Health Unit level in the whole country who can then tackle issues at a local ground level. Specific health messages through electronic media can promote the importance of a healthy, prestigious woman in the family [24,25]. Woman employment and family planning programs should be stressed [26] to relieve the misery of poverty. Further extended studies should be conducted to probe into each of the variables (cultural, social, genetic composition, education) in detail in each individual ethnic group.

CONSENT

All authors declare that written informed consent was obtained from the patient.

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki. The ethics committee is that of the Research department of the Creek General Hospital and UMDC.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Nicholas JK, Rashmi J, Mohsen N, Sarah KW, Nicole J, Rafael L, et al. A systematic analysis of global anaemia burden from 1990 to 2010. *Blood*. 2014;123:615-24.
- World Health Organisation. The global prevalence of anaemia in 2011. *Nutrition*. Available:http://www.who.int/nutrition/publications/micronutrients/global_prevalence_anaemia_2011/en/ (Accessed February 20, 2018)
- Saeed A, Anwaar A, Asif A, Zulfiqar A, Muhammad R, Tariq I. Iron status of the Pakistani population-current issues and strategies. *Asia Pac J Clin Nutr*. 2013;22(3):340-47.
- Nageen A, Bashir F, Suhail S, Ki, Munir M, Ara J. Anaemia.... An ongoing challenge - a focus on the anaemic parameters of adults of Asia. *WJPMR*. 2018;4(1):05-11.
- National Research Council (US) Panel on Race, Ethnicity, and Health in Later Life; Anderson NB, Bulatao RA, Cohen B, editors. *Critical Perspectives on Racial and Ethnic Differences in Health in Late Life*. Washington (DC): National Academies Press (US). Genetic Factors in Ethnic Disparities in Health. 2004;8. Available:<https://www.ncbi.nlm.nih.gov/books/NBK25517> (Accessed March 16, 2018)
- Dr Mary Lowth. Diseases and different ethnic groups. *Patient*; 2015. Available:<https://patient.info/doctor/diseases-and-different-ethnic-groups> (Accessed on April 23, 2018)
- Darlington F, Norman P, Ballas D, Exeter J. Exploring ethnic inequalities in health: Evidence from the Health Survey for England, 1998-2011. *Diversity and Equality in Health and Care*. Available:<http://diversityhealthcare.imedpub.com/exploring-ethnic-inequalities-in-health-evidencefrom-the-health-survey-for-england-19982011.php?aid=7680> (Accessed March 18, 2018)
- Wikipedia contributors, "Ethnic groups in Pakistan," Wikipedia. The Free Encyclopedia. Available:https://en.wikipedia.org/w/index.php?title=Ethnic_groups_in_Pakistan&oldid=830456658 (Accessed March 26, 2018)
- Hemoglobin concentrations for the diagnosis of anaemia and assessment of severity. *VMNIS | Vitamin and Mineral Nutrition Information System*. WHO/NMH/NHD/MNM/11.1.
- Sameer Mandhro. Too many mouths to feed in Karachi's slum fishermen's colony. Published: July 11; 2015. Available:<https://tribune.com.pk/story/918808/too-many-mouths-to-feed-in-karachis-slum-fishermens-colony>
- Faruk A. Anaemia in Bangladesh: A review of prevalence and etiology. *Public Health Nutrition*. 2000;3(4):385-93.
- Bharati P, Ghosh R, Gupta R. Socioeconomic condition and anaemia among the Mahishya population of Southern West Bengal, India. *Mal J Nutr*. 2004;10(1):23-30.
- Merrill RD, Shamim AA, Ali H, Labrique AB, Schulze K, Christian P, West KP. High prevalence of anaemia with lack of iron deficiency among women in rural Bangladesh: A role for thalassemia and iron in groundwater. *Asia Pac J Clin Nutr*. 2012;21(3):416-24.
- Bhutta Z, Soofi S, Zaidi S, Habib A, Hussain M. *Pakistan National Nutrition Survey*; 2011. Available:http://ecommons.aku.edu/pakistan_fhs_mc_women_childhealth_paediatr/262 (Accessed at March 3, 2018)
- Mudaser Kazi. On the rise: No significant improvement in fighting anaemia in Sindh. Published: December 23; 2017. Available:<https://tribune.com.pk/story/1581600/1-rise-no-significant-improvement-fighting-anaemia-sindh>
- Laghari Z, Baig MNM, Memon F, Panhwar F, Qambarani M, Palh Z. Correlation of BMI and MUAC with

- anaemia among Sindh University Students, Jamshoro, Pakistan. Sindh Univ. Res. Jour. (Sci. Ser.) 2017;49(3): 553-56.
17. Shahab F, Sikandar S, Shahab A, Raziq F. Frequency of anaemia in patients presenting to tertiary care hospital in Peshawar, Pakistan. Khyber Med Univ J. 2015;7(1):30-3.
18. Syed Ali Shah. Women in Balochistan severely malnourished, experts warn; 2014. Available:<https://www.dawn.com/news/1147717/women-in-balochistan-severely-malnourished-experts-warn> (Accessed at April 14, 2018)
19. 61% people facing food insecurity in Balochistan Source: Daily Balochistan Express. Available:<http://www.bexpress.com.pk/2017/07/httpwp-mep5f0gk-8v0> (Accessed at April 14, 2018)
20. Uzma H. Frequency of anaemia in married women in Jutial, Gilgit. PAFMJ. 2013;63:3.
21. Ahmed K, Mehtab S, Imran R, Shuja N, Shah G. Prevalence of intestinal parasitic pathogens among gastroenteritis patients in District Gilgit, Gilgit-Baltistan, Pakistan. Pakistan J. Zool. 2012;44(4):1059-63.
22. Ahmed T, Rahman S, Ahmed S, Siddiqui A, Javed A, Kamal J, Ahmed L. Frequency of vitamin B12 and red cell folate deficiency in macrocytic anaemia. Journal of Basic & Applied Sciences. 2012;8:706-13.
23. Callen M, Saad Gulzar S, Ali Hasanain A, Abdul Rehman Khan AR, Yasir Khan Y, Zia M. Improving public health delivery in Punjab, Pakistan: Issues and opportunities. The Lahore Journal of Economics. 2013;18:249-69.
24. Hafeez M. Poverty and poor health in Pakistan: Exploring the effects of privatizing healthcare. Harvard International Review. Available:hir.harvard.edu/article/?a=5768 (Accessed at May 15, 2018)
25. Harding KL, Victor MA, Grace N, Patrick W. Determinants of anaemia among women and children in Nepal and Pakistan: An analysis of recent national survey data. Matern Child Nutr. 2017;e12478 Available:<https://doi.org/10.1111/mcn.12478>
26. Research and Development Solutions. Policy briefs series no. 41, December 2013 key findings of the national nutrition survey of 2011. Available:http://www.resdev.org/files/policy_brief/41/Policy%20Brief%2041%20-%20Nutritional%20Status.pdf (Accessed at April 24, 2018)

© 2018 Nageen et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/24877>